

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

Leave blank

General Certificate of Education
 June 2003
 Advanced Level Examination



PHYSICS (SPECIFICATION A) PHA8/W
Unit 8 Nuclear Instability: Turning Points in Physics Option

Friday 20 June 2003 Afternoon Session

In addition to this paper you will require:

- a calculator;
- a pencil and a ruler.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Time allowed: 1 hour 15 minutes

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

Information

- The maximum mark for this paper is 40.
- Mark allocations are shown in brackets.
- The paper carries 10% of the total marks for Physics Advanced.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- In questions requiring description and explanation you will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary where appropriate. The degree of legibility of your handwriting and the level of accuracy of your spelling, punctuation and grammar will also be taken into account.

Data Sheet

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Fundamental constants and values				Mechanics and Applied Physics	Fields, Waves, Quantum Phenomena
Quantity	Symbol	Value	Units		
speed of light in vacuo	c	3.00×10^8	m s^{-1}	$v = u + at$	$g = \frac{F}{m}$
permeability of free space	μ_0	$4\pi \times 10^{-7}$	H m^{-1}	$s = \left(\frac{u+v}{2}\right)t$	$g = -\frac{GM}{r^2}$
permittivity of free space	ϵ_0	8.85×10^{-12}	F m^{-1}	$s = ut + \frac{at^2}{2}$	$g = -\frac{\Delta V}{\Delta x}$
charge of electron	e	1.60×10^{-19}	C	$v^2 = u^2 + 2as$	$V = -\frac{GM}{r}$
the Planck constant	h	6.63×10^{-34}	J s	$P = Fv$	$a = -(2\pi f)^2 x$
gravitational constant	G	6.67×10^{-11}	$\text{N m}^2 \text{kg}^{-2}$	$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$	$v = \pm 2\pi f \sqrt{A^2 - x^2}$
the Avogadro constant	N_A	6.02×10^{23}	mol^{-1}	$\omega = \frac{v}{r} = 2\pi f$	$x = A \cos 2\pi ft$
molar gas constant	R	8.31	$\text{J K}^{-1} \text{mol}^{-1}$	$a = \frac{v^2}{r} = r\omega^2$	$T = 2\pi\sqrt{\frac{m}{k}}$
the Boltzmann constant	k	1.38×10^{-23}	J K^{-1}	$I = \sum mr^2$	$T = 2\pi\sqrt{\frac{l}{g}}$
the Stefan constant	σ	5.67×10^{-8}	$\text{W m}^{-2} \text{K}^{-4}$	$E_k = \frac{1}{2} I\omega^2$	$\lambda = \frac{\omega s}{D}$
the Wien constant	α	2.90×10^{-3}	m K	$\omega_2 = \omega_1 + \alpha t$	$d \sin \theta = n\lambda$
electron rest mass	m_e	9.11×10^{-31}	kg	$\theta = \omega_1 t + \frac{1}{2} \alpha t^2$	$\theta = \frac{\lambda}{D}$
(equivalent to $5.5 \times 10^{-4} \text{u}$)				$\omega_2^2 = \omega_1^2 + 2\alpha\theta$	$1^n n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$
electron charge/mass ratio	e/m_e	1.76×10^{11}	C kg^{-1}	$\theta = \frac{1}{2} (\omega_1 + \omega_2)t$	$1^n n_2 = \frac{n_2}{n_1}$
proton rest mass	m_p	1.67×10^{-27}	kg	$T = I\alpha$	$\sin \theta_c = \frac{1}{n}$
(equivalent to 1.00728u)				<i>angular momentum</i> = $I\omega$	$E = hf$
proton charge/mass ratio	e/m_p	9.58×10^7	C kg^{-1}	$W = T\theta$	$hf = \phi + E_k$
neutron rest mass	m_n	1.67×10^{-27}	kg	$P = T\omega$	$hf = E_1 - E_2$
(equivalent to 1.00867u)				<i>angular impulse</i> = change of <i>angular momentum</i> = Tt	$\lambda = \frac{h}{p} = \frac{h}{mv}$
gravitational field strength	g	9.81	N kg^{-1}	$\Delta Q = \Delta U + \Delta W$	$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$
acceleration due to gravity	g	9.81	m s^{-2}	$\Delta W = p\Delta V$	Electricity
atomic mass unit	u	1.661×10^{-27}	kg	$pV^\gamma = \text{constant}$	$\epsilon = \frac{E}{Q}$
(1u is equivalent to 931.3 MeV)				<i>work done per cycle</i> = area of loop	$\epsilon = I(R + r)$
Fundamental particles				<i>input power</i> = calorific value \times fuel flow rate	$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$
<i>Class</i>	<i>Name</i>	<i>Symbol</i>	<i>Rest energy</i>	<i>indicated power</i> as (area of $p-V$ loop) \times (no. of cycles/s) \times (no. of cylinders)	$R_T = R_1 + R_2 + R_3 + \dots$
			/MeV	<i>friction power</i> = indicated power - brake power	$P = I^2 R$
photon	photon	γ	0	$\text{efficiency} = \frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$	$E = \frac{F}{Q} = \frac{V}{d}$
lepton	neutrino	ν_e	0	<i>maximum possible efficiency</i> = $\frac{T_H - T_C}{T_H}$	$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$
		ν_μ	0		$E = \frac{1}{2} QV$
	electron	e^\pm	0.510999		$F = BI$
	muon	μ^\pm	105.659		$F = BQv$
mesons	pion	π^\pm	139.576		$Q = Q_0 e^{-t/RC}$
		π^0	134.972		$\Phi = BA$
	kaon	K^\pm	493.821		
		K^0	497.762		
baryons	proton	p	938.257		
	neutron	n	939.551		
Properties of quarks					
<i>Type</i>	<i>Charge</i>	<i>Baryon number</i>	<i>Strangeness</i>		
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0		
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0		
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1		
Geometrical equations					
<i>arc length</i> = $r\theta$					
<i>circumference of circle</i> = $2\pi r$					
<i>area of circle</i> = πr^2					
<i>area of cylinder</i> = $2\pi rh$					
<i>volume of cylinder</i> = $\pi r^2 h$					
<i>area of sphere</i> = $4\pi r^2$					
<i>volume of sphere</i> = $\frac{4}{3} \pi r^3$					

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	2.00×10^{30}	7.00×10^8
Earth	6.00×10^{24}	6.40×10^6

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

Electronics

Resistors

Preferred values for resistors (E24)
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2
2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2
6.8 7.5 8.2 9.1 ohms
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

Alternating Currents

$$f = \frac{1}{T}$$

Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$


$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left(\frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

SECTION A: NUCLEAR INSTABILITY

Answer **all** parts of this question.

1 **Figure 1** shows a grid of neutron number against proton number. A nucleus ${}^A_Z\text{X}$ is marked.



- (a) Draw arrows on **Figure 1**, each starting on ${}^A_Z\text{X}$ and ending on a daughter nucleus after the following transitions:
- (i) β^- emission (label this arrow A)
neutron emission (label this arrow B)
electron capture (label this arrow C).
 - (ii) Give the equation for electron capture by the nucleus ${}^A_Z\text{X}$.

.....
(4 marks)

Turn over ►

- (b) When ${}_{12}^{27}\text{Mg}$ decays to ${}_{13}^{27}\text{Al}$ by β^- decay, the daughter nucleus is produced in one of two possible excited states. These two states are shown in **Figure 2** together with their corresponding energies.

${}_{12}^{27}\text{Mg}$	$E/10^{-13}\text{J}$
	4.18
	1.63
	1.33
${}_{13}^{27}\text{Al}$	0.00 (ground state)

Figure 2

- (i) Calculate the maximum possible kinetic energy, in J, which an emitted β^- particle can have.

.....

.....

- (ii) The excited aluminium nuclei emit γ photons. Calculate each of the three possible γ photon energies in J.

.....

.....

.....

- (iii) Calculate the frequency of the most energetic γ photon emitted.

.....

.....

.....

(3 marks)

- (c) (i) State and explain **two** precautions that should be taken when working with a sample of $^{27}_{12}\text{Mg}$ in a school laboratory.

You may be awarded marks for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Discuss which of the two types of radiation, β^- or γ , emitted from a sample of $^{27}_{12}\text{Mg}$ would be the more hazardous.

.....

.....

.....

(3 marks)

10

TURN OVER FOR THE NEXT QUESTION

Turn over ▶

SECTION B: TURNING POINTS IN PHYSICS

Answer **all** questions.

2 In an experiment to measure the charge of an oil droplet, a positively charged oil droplet was held stationary by means of a uniform electric field of strength $4.9 \times 10^5 \text{ V m}^{-1}$.

(a) (i) What was the direction of the electric field?

.....

(ii) Show that the specific charge of the oil droplet was $2.0 \times 10^{-5} \text{ C kg}^{-1}$.

.....
.....
.....
.....

(3 marks)


(b) When the electric field was switched off the oil droplet fell and quickly reached constant speed. Explain why the oil droplet reached constant speed.

.....
.....
.....
.....
.....
.....

(3 marks)

6

3 The Michelson-Morley experiment represented in the diagram was designed to find out if the speed of light depended on its direction relative to the Earth's motion through space. Interference fringes were seen by the observer.



(a) (i) Explain why interference fringes were seen.

.....

.....

.....

.....

.....

.....

.....

(ii) The interference fringe pattern did not shift when the apparatus was rotated by 90° . Explain the significance of this null observation.

.....

.....

.....

.....

.....

.....

.....

(5 marks)

Turn over ▶

(b) Einstein postulated that the speed of light in free space is invariant. Explain what is meant by this postulate.

.....
.....
.....
.....
.....
.....

(2 marks)



4 (a) Describe **one** piece of evidence that shows that matter has

(i) a wave-like nature,

.....
.....
.....
.....
.....

(ii) a particle-like nature.

.....
.....
.....
.....
.....

(3 marks)

(b) For a proton of kinetic energy 5.0 MeV,

(i) show that its speed is $3.1 \times 10^7 \text{ m s}^{-1}$,

.....

.....

.....

.....

.....

.....

(ii) calculate its de Broglie wavelength.

.....

.....

.....

.....

.....

.....


(4 marks)

○
—
7

TURN OVER FOR THE NEXT QUESTION

Turn over ▶

- 5 A narrow beam of electrons at a speed of $3.2 \times 10^7 \text{ m s}^{-1}$ travels along a circular path in a uniform magnetic field of flux density, B , as shown in the diagram.



- (a) Explain why the path of the beam in the field is circular.

You may be awarded marks for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

(3 marks)

- (b) (i) Show that the speed, v , of the electrons in the field is given by

$$v = \frac{Ber}{m},$$

where r is the radius of the circular path of the beam in the field.

.....

.....

.....

.....

.....

- (ii) When the flux density was 7.3 mT, the radius of the circular path of the beam in the field was 25 mm. Use the data to calculate the specific charge of the electron.

.....

.....

.....

.....

.....

.....

.....

.....

(5 marks)

8

QUALITY OF WRITTEN COMMUNICATION (2 marks)

2

END OF QUESTIONS