

Surname		Other Names	
Centre Number		Candidate Number	
Candidate Signature			

Leave blank

General Certificate of Education
January 2004
Advanced Subsidiary Examination



PHYSICS (SPECIFICATION A)
Unit 1 Particles, Radiation and Quantum Phenomena

PA01

Monday 12 January 2004 Morning Session

In addition to this paper you will require:

- a calculator;
- a pencil and a ruler.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Time allowed: 1 hour

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

Information

- The maximum mark for this paper is 50.
- Mark allocations are shown in brackets.
- The paper carries 30% of the total marks for Physics Advanced Subsidiary and carries 15% of the total marks for Physics Advanced.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- In questions requiring description and explanation you will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary where appropriate. The degree of legibility of your handwriting and the level of accuracy of your spelling, punctuation and grammar will also be taken into account.

Data Sheet

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Fundamental constants and values				Mechanics and Applied Physics	Fields, Waves, Quantum Phenomena
Quantity	Symbol	Value	Units		
speed of light in vacuo	c	3.00×10^8	m s^{-1}	$v = u + at$	$g = \frac{F}{m}$
permeability of free space	μ_0	$4\pi \times 10^{-7}$	H m^{-1}	$s = \left(\frac{u+v}{2}\right)t$	$g = -\frac{GM}{r^2}$
permittivity of free space	ϵ_0	8.85×10^{-12}	F m^{-1}	$s = ut + \frac{at^2}{2}$	$g = -\frac{\Delta V}{\Delta x}$
charge of electron	e	1.60×10^{-19}	C	$v^2 = u^2 + 2as$	$V = -\frac{GM}{r}$
the Planck constant	h	6.63×10^{-34}	J s	$F = \frac{\Delta(mv)}{\Delta t}$	$a = -(2\pi f)^2 x$
gravitational constant	G	6.67×10^{-11}	$\text{N m}^2 \text{kg}^{-2}$	$P = Fv$	$v = \pm 2\pi f \sqrt{A^2 - x^2}$
the Avogadro constant	N_A	6.02×10^{23}	mol^{-1}	$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$	$x = A \cos 2\pi ft$
molar gas constant	R	8.31	$\text{J K}^{-1} \text{mol}^{-1}$	$\omega = \frac{v}{r} = 2\pi f$	$T = 2\pi \sqrt{\frac{m}{k}}$
the Boltzmann constant	k	1.38×10^{-23}	J K^{-1}	$a = \frac{v^2}{r} = r\omega^2$	$T = 2\pi \sqrt{\frac{l}{g}}$
the Stefan constant	σ	5.67×10^{-8}	$\text{W m}^{-2} \text{K}^{-4}$	$I = \sum mr^2$	$\lambda = \frac{ws}{D}$
the Wien constant	α	2.90×10^{-3}	m K	$E_k = \frac{1}{2} I\omega^2$	$d \sin \theta = n\lambda$
electron rest mass	m_e	9.11×10^{-31}	kg	$\omega_2 = \omega_1 + at$	$\theta \approx \frac{\lambda}{D}$
(equivalent to $5.5 \times 10^{-4}u$)				$\theta = \omega_1 t + \frac{1}{2} at^2$	$1/n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$
electron charge/mass ratio	e/m_e	1.76×10^{11}	C kg^{-1}	$\omega_2^2 = \omega_1^2 + 2a\theta$	$1/n_2 = \frac{n_2}{n_1}$
proton rest mass	m_p	1.67×10^{-27}	kg	$\theta = \frac{1}{2}(\omega_1 + \omega_2)t$	$\sin \theta_c = \frac{1}{n}$
(equivalent to 1.00728u)				$T = I\alpha$	$E = hf$
proton charge/mass ratio	e/m_p	9.58×10^7	C kg^{-1}	<i>angular momentum</i> = $I\omega$	$hf = \phi + E_k$
neutron rest mass	m_n	1.67×10^{-27}	kg	$W = T\theta$	$hf = E_1 - E_2$
(equivalent to 1.00867u)				$P = T\omega$	$\lambda = \frac{h}{p} = \frac{h}{mv}$
gravitational field strength	g	9.81	N kg^{-1}	<i>angular impulse</i> = change of <i>angular momentum</i> = Tt	$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$
acceleration due to gravity	g	9.81	m s^{-2}	$\Delta Q = \Delta U + \Delta W$	Electricity
atomic mass unit	u	1.661×10^{-27}	kg	$\Delta W = p\Delta V$	$\epsilon = \frac{E}{Q}$
(1u is equivalent to 931.3 MeV)				$pV^\gamma = \text{constant}$	$\epsilon = I(R+r)$
Fundamental particles				<i>work done per cycle</i> = area of loop	$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$
<i>Class</i>	<i>Name</i>	<i>Symbol</i>	<i>Rest energy</i>	<i>input power</i> = calorific value \times fuel flow rate	$R_T = R_1 + R_2 + R_3 + \dots$
			/MeV	<i>indicated power</i> as (area of $p-V$ loop) \times (no. of cycles/s) \times (no. of cylinders)	$P = I^2 R$
photon	photon	γ	0	<i>friction power</i> = indicated power - brake power	$E = \frac{F}{Q} = \frac{V}{d}$
lepton	neutrino	ν_e	0	<i>efficiency</i> = $\frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$	$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$
		ν_μ	0	<i>maximum possible efficiency</i> = $\frac{T_H - T_C}{T_H}$	$E = \frac{1}{2} QV$
mesons	electron	e^\pm	0.510999		$F = BI$
		muon	μ^\pm	105.659	
	pion	π^\pm	139.576		$Q = Q_0 e^{-t/RC}$
		π^0	134.972		$\Phi = BA$
kaon	K^\pm	493.821			
	K^0	497.762			
baryons	proton	p	938.257		
	neutron	n	939.551		
Properties of quarks					
<i>Type</i>	<i>Charge</i>	<i>Baryon number</i>	<i>Strangeness</i>		
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0		
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0		
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1		
Geometrical equations					
<i>arc length</i> = $r\theta$					
<i>circumference of circle</i> = $2\pi r$					
<i>area of circle</i> = πr^2					
<i>area of cylinder</i> = $2\pi rh$					
<i>volume of cylinder</i> = $\pi r^2 h$					
<i>area of sphere</i> = $4\pi r^2$					
<i>volume of sphere</i> = $\frac{4}{3}\pi r^3$					

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F l}{A e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	2.00×10^{30}	7.00×10^8
Earth	6.00×10^{24}	6.40×10^6

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ kms}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

Electronics

Resistors

Preferred values for resistors (E24)
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2
2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2
6.8 7.5 8.2 9.1 ohms
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

Alternating Currents

$$f = \frac{1}{T}$$

Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left(\frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

Answer **all** questions in the spaces provided.

1 (a) Give the number of nucleons and the number of electrons in an atom of $^{22}_{11}\text{Na}$.

nucleons.....

electrons

(2 marks)

(b) The isotope $^{22}_{11}\text{Na}$ is a positron emitter. In positron emission an up quark undergoes the following change,

$$u \rightarrow d + \beta^+ + \nu_e.$$

Show that charge, lepton number and baryon number are conserved in this decay.

charge

lepton number

baryon number

(3 marks)

(c) Describe what happens when a positron collides with an electron.

.....

.....

.....

(2 marks)

7

TURN OVER FOR THE NEXT QUESTION

Turn over ▶

- 2 (a) Quarks may be combined together in a number of ways to form sub-groups of hadrons. Name **two** of these sub-groups and for each, state its quark composition.

sub-group 1

.....

sub-group 2

.....

(3 marks)

- (b) A free neutron is an unstable particle.

- (i) Complete the following to give an equation that represents the decay of a neutron.



- (ii) Describe the change that occurs to the quark structure when a neutron decays.

.....

.....

.....

.....


.....

.....

(4 marks)

7

3 The diagram shows a ray of monochromatic light, in the plane of the paper, incident on the end face of an optical fibre.



- (a) (i) Draw on the diagram the complete path followed by the incident ray, showing it entering into the fibre and emerging from the fibre at the far end.
- (ii) State any changes that occur in the speed of the ray as it follows this path from the source. Calculations are not required.

.....

 (4 marks)

- (b) (i) Calculate the critical angle for the optical fibre at the air boundary.

refractive index of the optical fibre glass = 1.57

.....

- (ii) The optical fibre is now surrounded by cladding of refractive index 1.47. Calculate the critical angle at the core-cladding boundary.


.....

- (iii) State **one** advantage of cladding an optical fibre.

.....

(6 marks)

- 4 In the apparatus shown, monochromatic ultraviolet radiation is incident on the surface of metal **X**. Photoelectrons are emitted from **X** and are collected at electrode **Y**.



- (a) Calculate the work function of **X**, given that each photon in the incident radiation has $3.2 \times 10^{-19} \text{ J}$ of energy.
The maximum kinetic energy possessed by a single photoelectron is $2.1 \times 10^{-19} \text{ J}$.

.....

(3 marks)

- (b) The source of the incident radiation is replaced with a new source. The wavelength of the radiation from the new source is half the wavelength of the original radiation.

Calculate the maximum kinetic energy of the emitted photoelectrons.

.....

(3 marks)

5 The diagram shows some energy levels, in eV, of an atom.

energy/eV	level
-0.1	n = 4
-3.1	n = 3
-12.4	n = 2
-18.6	n = 1 (ground state)

Photons of specific wavelengths are emitted from these atoms when they are *excited* by collisions with electrons.

You may be awarded marks for the quality of written communication in your answer.

(a) Explain

(i) what is meant by the process of excitation,

.....

.....

.....

.....

(ii) why the emitted photons have specific wavelengths.

.....

.....

.....

.....

.....

(5 marks)

(b) One of the emitted photons has an energy of $9.92 \times 10^{-19} \text{ J}$.

(i) Calculate the wavelength of this photon.

.....
.....
.....
.....

(ii) Determine which transition is responsible for this emitted photon.

.....
.....
.....

(iii) Draw an arrow on the energy level diagram on **page 10** to show the transition responsible for the emission of a photon with the shortest wavelength.

(7 marks)

12

TURN OVER FOR THE NEXT QUESTION

Turn over ►

