



**GCE AS/A level**

**1092/01**

**CHEMISTRY – CH2**

**P.M. TUESDAY, 3 June 2014**

**1 hour 30 minutes plus your additional time allowance**

**Surname** \_\_\_\_\_

**Other Names** \_\_\_\_\_

**Centre Number** \_\_\_\_\_

**Candidate Number** 2 \_\_\_\_\_

<b>For Examiner's use only</b>		
<b>Question</b>	<b>Maximum Mark</b>	<b>Mark Awarded</b>
<b>Section A</b> <b>1. to 6.</b>	<b>10</b>	
<b>Section B</b> <b>7.</b>	<b>16</b>	
<b>8.</b>	<b>16</b>	
<b>9.</b>	<b>15</b>	
<b>10.</b>	<b>11</b>	
<b>11.</b>	<b>12</b>	
<b>Total</b>	<b>80</b>	

## **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need a:  
  
calculator;

**DATA SHEET** containing a **PERIODIC TABLE** supplied by **WJEC**. Refer to it for any **RELATIVE ATOMIC MASSES** you require.

## **INSTRUCTIONS TO CANDIDATES**

Use black ink, black ball-point pen or your usual method.

Write your name, centre number and candidate number in the spaces provided on the front cover.

**SECTION A** Answer ALL questions in the spaces provided.

**SECTION B** Answer ALL questions in the spaces provided.

Candidates are advised to allocate their time appropriately between **SECTION A (10 MARKS)** and **SECTION B (70 MARKS)**.

## **INFORMATION FOR CANDIDATES**

**The number of marks is given in brackets at the end of each question or part-question.**

**The maximum mark for this paper is 80.**

**Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.**

**The QWC label alongside particular part-questions indicates those where the Quality of Written Communication is assessed.**

**If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.**

**SECTION A**

**Answer ALL questions in the spaces provided.**

- 1. Put the following in order of increasing strength.**

**[1]**

**covalent bonds**


**hydrogen bonds**

**van der Waals' forces**

**weakest** \_\_\_\_\_

\_\_\_\_\_ **strongest**

2. Give the **SYSTEMATIC** name of the compound whose structure is shown below. [1]



3. Draw dot-and-cross diagrams to show the formation of calcium chloride from atoms of chlorine and calcium. [2]

4. The table below gives the electronegativity values of some elements.

Atom	H	N	O	Al	Cl
Electronegativity value	2.1	3.0	3.5	1.6	3.0

- (a) Use the data in the table to identify any dipoles present in the following bonds. Mark their polarity clearly. [1]



- (b) Use the data to give a reason why aluminium chloride is considered to be a covalent compound, while aluminium oxide is an ionic compound. [1]

---

---

---

---


---

---


---



hex-2-ene



cyclohexane



5. Cyclohexane and hex-2-ene, shown opposite, are isomers. Give a chemical test to distinguish between these two compounds. [2]

Reagent(s) \_\_\_\_\_

\_\_\_\_\_

Observations \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

6. Select ALL the molecules from the list below that have bond angles of less than  $109^\circ$ .



---

[2]

SECTION A TOTAL [10]

	<b>magnesium nitrate</b>	<b>barium chloride</b>	<b>sodium hydroxide</b>
<b>potassium carbonate</b>	<b>white precipitate</b>	<b>white precipitate</b>	<b>no visible change</b>
<b>sodium hydroxide</b>			
<b>barium chloride</b>			

**SECTION B**

Answer ALL questions in the spaces provided.

7. Ewan and Gwyneth are given four unlabelled bottles. They know that these contain the following four solutions:

potassium carbonate

sodium hydroxide

barium chloride

magnesium nitrate

- (a) Ewan predicted what will happen when each of the four solutions is added to the others, and presented this information in the grid opposite.

(i) Complete the three empty boxes opposite with the observations expected in each of these cases. [2]

(ii) Name the white precipitate formed when magnesium nitrate is mixed with potassium carbonate, and write an IONIC equation for its formation. [2]

Name of precipitate \_\_\_\_\_

Ionic equation \_\_\_\_\_

**7(b) Gwyneth uses different tests to identify the four solutions. Each test allows her to distinguish between some of the solutions. For each test state the solution(s) that would give a visible change and the observation(s) that would be made.**

**(i) Addition of litmus solution [1]**

---

---

---

---

---

---

---

---

---

---

**7(b) (ii) Flame test [2]**

---

---

---

---

---

---

---

---

---

---

**7(b) (iii) Addition of sodium sulfate solution [2]**

---

---

---

---

---

---

---

---

---

---



**7(c) Ewan and Gwyneth are provided with a white solid that they believe to be sodium bromide or sodium iodide.**

**(i) They dissolve the solid in water to make a solution. Explain what occurs when an ionic solid such as sodium bromide dissolves in water. [2]**

---

---

---

---

---

---

---

---

---

---

- 7(c) (ii) **Gwyneth uses aqueous silver nitrate to identify the solution. Give the observations expected when silver nitrate is added separately to solutions of sodium bromide and sodium iodide. [2]**

**Observation with sodium bromide**

---

---

---

---

**Observation with sodium iodide**

---

---

---

---

**7(c) (iii) Ewan thinks that a further test is needed after addition of the silver nitrate to distinguish between sodium bromide and sodium iodide. Give the reagent and observations for this further test. [2]**

**Reagent** \_\_\_\_\_

**Observation with sodium bromide**

**Observation with sodium iodide**

- 7(c) (iv) When bromine water is added to a solution of sodium iodide, a reaction occurs. Write an equation for this reaction. [1]

---

---

Total [16]

8. Crude oil is a complex mixture of hydrocarbons, with samples from different locations in the world having different compositions. The table below gives the composition of crude oil from two locations.

Fraction	Percentage by mass	
	Brent Crude	Gulf of Suez
petroleum gases	2.4	1.2
naphtha	19.1	13.6
kerosene	14.2	12.7
gas oil	20.9	18.7
residue	43.4	53.8

**8(a) The different fractions are separated by fractional distillation. Explain why the different fractions have different boiling temperatures. [2]**

---

---

---

---

---

---

---

---

---

---

**8(b) The petroleum gases produced from crude oil can contain both propane and butane.**

- (i) A barrel of Gulf of Suez crude oil has a mass of 145 kg. Assuming all the petroleum gas released from the oil is butane, calculate the volume that this gas would occupy at 1 atmosphere pressure. [3]**

**[1 mol of gas occupies  $24.0 \text{ dm}^3$  under these conditions]**

**Volume = \_\_\_\_\_  $\text{dm}^3$**

**8(b) (ii) Propane can be chlorinated by a similar method to methane.**

**I Give the condition(s) required for the chlorination of propane. [1]**

---

---

**II Write an equation for the initiation stage of the chlorination of propane. [1]**

---

---

**III The chlorination of propane also produces hexane as a minor product.  
Explain how this compound forms.**

**[2]**

---

---

---

---

---

---

---



**8(c) Naphtha is used as a starting material for the production of alkenes, and these are then used to produce polymers such as poly(ethene). Discuss how poly(ethene) is produced, starting from naphtha. Your answer should include:**

- An explanation of which of the two types of crude oil given would be more useful for producing alkenes.**
- How the naphtha is converted into alkenes.**
- An equation for the production of ethene from decane, an alkane with 10 carbon atoms.**
- An explanation of what is meant by polymerisation.**
- An equation for the polymerisation of ethene, clearly stating the type of polymerisation that is occurring.**
- A different polymer in common use, with the structure of the monomer used in its production.**

**[6]**

**QWC [1]**

---

---

---

---





9. Haematite is an ore of iron that contains  $\text{Fe}_2\text{O}_3$ . Iron is extracted from this ore in a blast furnace.

(a) Balance the equation opposite for the extraction of iron from  $\text{Fe}_2\text{O}_3$ . [1]

(b) Use oxidation states to show that the reaction in (a) is a redox reaction. [2]

---

---

---

---

---

---

---

---

---

---

**9(c) A different oxide of iron is iron(II) oxide, FeO. The ions in this compound adopt an arrangement similar to that of sodium chloride.**

**(i) Give the crystal co-ordination numbers for the ions in FeO. [1]**

---

---

**(ii) Draw the arrangement of oxide ions around each iron ion. [1]**

- 9(d) Iron can be extracted from FeO according to the equation below.



Calculate the mass of iron that could be extracted from 20.0 kg of iron(II) oxide, FeO. [3]

Mass of iron = \_\_\_\_\_ kg

**9(e) Carbon monoxide contains two covalent bonds and one co-ordinate bond. Explain what is meant by the terms COVALENT BOND and CO-ORDINATE BOND, indicating the difference between them. [2]**

---

---

---

---

---

---

---

---

---

---

**9(f) Iron is a typical metal. Describe the bonding present in iron. Explain how it can conduct electricity and why it has a high melting temperature.**

**[4]**

**QWC [1]**

---

---

---

---

---

---

---

---

---

---



---

---

---

---

---

---

---

---

---

---

---

---

---



---

---

---

---

**Total [15]**



**10(a) 1-bromobutane is a liquid that is insoluble in water. It can be converted to butan-1-ol in a one-step reaction as shown opposite.**

**(i) Give the reagent(s) and condition(s) required for this reaction. [2]**

---

---

---

---

**(ii) Explain why butan-1-ol is soluble in water whilst 1-bromobutane is not. [3]**

---

---

---

---

---

---

---



---

---

---

---

---



**10(b) Butan-1-ol can be converted into liquid butanoic acid in a one-step reaction as shown opposite.**

- (i) Give the reagent(s) and condition(s) required for this reaction. [2]**

---

---

---

---

- (ii) Explain why butanoic acid has a much higher boiling temperature than 1-bromobutane. [3]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**10(b) (iii)      The reaction as shown opposite page 33 frequently produces a mixture containing unreacted butan-1-ol and butanoic acid. State how these two liquids could be separated. [1]**

---

---

---

---

**Total [11]**

**11(a) Propene reacts with hydrogen bromide to give 2-bromopropane.**

**(i) Draw the mechanism for this reaction. [3]**



**11(a) (ii) Explain why the product of this reaction is mainly 2-bromopropane rather than 1-bromopropane. [2]**

---

---

---

---

---

---

---

11(b) Compound **C** is a compound of carbon, hydrogen and bromine only. Bromine has two isotopes,  $^{79}\text{Br}$  and  $^{81}\text{Br}$ , in equal abundance. Use all the information below to deduce the structure of compound **C**, giving your reasoning. [6]

QWC [1]

- Compound **C** contains 29.8% carbon, 4.2% hydrogen and 66.0% bromine by mass.
- The mass spectrum of compound **C** contains peaks at  $m/z$  of 15, 41 and a pair of peaks at 120 and 122.
- The infrared spectrum of compound **C** has absorptions at  $550\text{ cm}^{-1}$ ,  $1630\text{ cm}^{-1}$  and  $3030\text{ cm}^{-1}$ .
- Compound **C** is a Z-isomer.

---

---

---

---

---

---

---

---



---

---

---

---

**Total [12]**

**SECTION B TOTAL [70]**

**END OF PAPER**













