



**General Certificate of Education (A-level)  
June 2012**

**Statistics**

**SS02**

**(Specification 6380)**

**Statistics 2**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2012 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

## SS02

Q	Solution	Marks	Total	Comments
1 (a)	0.2, 0.3	B1, B1	2	CAO nms B2
(b)	$E(X) = 0 \times 0.2 + 1 \times 0.3 + 3 \times 0.5$ $= 1.8$ $E(X^2) = 0^2 \times 0.2 + 1^2 \times 0.3 + 3^2 \times 0.5$ $\text{Var}(X) = E(X^2) - E(X)^2 = 4.8 - 1.8^2$ $\text{s.d} = \sqrt{1.56} = 1.25$	M1 A1 M1 M1 A1		
(c)	$P(\text{points} > 3.05) = 0$	B1F	5	If <i>their</i> $E(X) + \text{s.d.} < 3$ accept 0.5
	<b>Total</b>		<b>1</b>	
			<b>8</b>	

Q	Solution	Marks	Total	Comments
<b>2 (a)(i)</b>	$n = 4$	B1	<b>1</b>	
<b>(ii)</b>	$(9.6 + 7.4 + 7.2 + 8.4) \div 4$ $y = 8.15$	M1 A1	<b>2</b>	Possibly implied 8.15 must be seen
<b>(iii)</b>	$8.0 + 8.8 + 7.0 + x = 4 \times 7.4$  $x = 5.8$	M1  A1	<b>2</b>	Move beyond moving average equation. 5.8 must be seen nms B2
<b>(b)(i)</b>	Correct position for $x$ Correct position for $y$	B1 B1	<b>2</b>	Within one square Within one square
<b>(ii)</b>	Short term (variation)..... about a downward (trend)	B1 B1	<b>2</b>	Accept negative/decreasing trend
<b>(iii)</b>	Residuals for Q2  $(+)0.3, (+)0.5$ and $(+)0.5$  Mean = 0.43	M1  A1  A1	<b>2</b>	Ignore sign for method mark. Attempt at min of 2 residuals  Must be +ve Their answer to 1d.p. Must use 3 residuals  Accept 0.4 to 0.45 Must use 3 residuals
<b>(iv)</b>	Read off from graph = 7.1 Add 0.4(3) 7.5(3) <b>million</b>	B1 M1 A1	<b>3</b>	Accept 7.05 to 7.15 Their 0.43 cao accept 7.5 to 7.6 <b>million</b>
<b>(v)</b>	Extrapolation far ahead is risky/not likely to be accurate.	E1	<b>3</b>	Anything indicating that things may be different two years later. Eg mention Olympics. Reason not necessary.
	<b>Total</b>		<b>16</b>	

Q	Solution	Marks	Total	Comments
<b>3(a)</b>	$2791 - (1+9+72+366+1173+11+22)$ $= 1137$	M1 A1	<b>2</b>	Or B2 for answer only seen.
<b>(b)</b>	Adding at least 10 numbers and $\div 10$ $65 \div 10 = 6.5$	M1 A1		Must specify divided by 10 SC 5.9 B1
<b>(c)(i)</b>	0.043(0)	B1	<b>2</b>	awrt
<b>(ii)</b>	$P(X > 10) = 1 - P(X \leq 10)$ $= 1 - 0.9332 = 0.0668$	M1 A1	<b>1</b>	Accept 0.067
<b>(iii)</b>	$P(X = 5) = P(X \leq 5) - P(X \leq 4)$ $= 0.3690 - 0.2237 = 0.145$	M1 A1	<b>2</b>	Or use of formula Answer alone scores B2
<b>(d)(i)</b>	9.6	B1	<b>2</b>	
<b>(ii)</b>	So Po(2.4) for 3 months $P(=0) = 0.0907$ <b>Or</b> Uses Po(1.625) and Po(0.775) And multiplies to give 0.0907	M1 A1  (M1) (A1)	<b>1</b>	Their '9.6' $\div 4$ as $\lambda$ for Poisson Accept 0.091
<b>(e)</b>	Poisson requires independence. eg Earthquake greater than 7.0 has associated (so non-independent) 6.0 or greater earthquakes. <b>Or</b> Poisson requires constant average rate. eg An earthquake of magnitude 7.0 will change the average rate.  Poisson inappropriate.	     E1  E1	<b>2</b>	Must <b>use</b> these Using formula, not tables  Must be in context
<b>Total</b>			<b>14</b>	

Q	Solution	Marks	Total	Comments
4(a)(i)	Cluster sampling	B1	1	
(ii)	Hypothesis test needs random sample	E1	1	Reference to sample
(b)	$H_0: \mu = 50$ $H_1: \mu \neq 50$ $z = \frac{(52.1 - 50)}{\sqrt{81}}$ $= +2.42$ <b>c.v. <math>z = \pm 2.5758</math> or <math>t_{80} = \pm 2.639</math></b>  So test statistic not in critical region. Accept $H_0$ PI  no significant evidence that mean mark for exam is not 50.	B1 B1 <b>M1</b>  <b>m1</b>  <b>A1</b> <b>B1</b>  A1F  E1F	1 1  8  2  2	$\mu$ or equivalent $\mu$ or equivalent <b>For general form for <math>z</math></b> Condone sign error <b>For use of <math>\sqrt{81}</math></b>  Must be +ve Or p value = 0.00776 Or p < 0.01  ft if M1 <b>and</b> A1 or B1 earned  Dep on A1F. For consistent conclusion in context. Must mention mark or exam.
(c)(i)	<b>Large sample</b> so mean can be treated as normally distributed. Conclusion still valid.	M1 A1	2	Any relevant comments with a correct conclusion. eg CLT
(ii)	The 81 may not be representative (or not random).  Conclusion not reliable.	M1 A1	2	Any relevant comments with a correct conclusion. eg biased <b>Not</b> reference to change in $n$ .
	<b>Alternatives for bold parts of (b)</b> $50 \pm 2.5758 \times \frac{7.8}{\sqrt{81}}$ $= (47.768, 52.232)$ <b>or</b> $50 \pm 2.639 \times \frac{7.8}{\sqrt{81}}$ $= (47.713, 52.287)$	(B1) (M1) (m1) (A1)  (B1) (M1) (m1) (A1)		For $z = \pm 2.5758$ For general form of equation For use of $\sqrt{81}$  For $t_{80} = \pm 2.639$ For general form of equation For use of $\sqrt{81}$
	<b>Total</b>		<b>14</b>	

Q	Solution	Marks	Total	Comments
<b>5(a)(i)</b>	28	B1	<b>1</b>	27 to 29 Must be an integer
<b>(ii)</b>	180 – “154” = 26 = 14.4%	M1 A1 A1		<b>3</b>
<b>(b)(i)</b>	Lowest end of whisker = 100 Q1 = 130 to 133 Q2 = 145 to 147 Q3 = 175 to 177 Top of whisker = 275	B1 B1 B1 B1 B1	<b>5</b>	
<b>(ii)</b>	Eg. Women have: Smaller range           OE Smaller IQR            OE Greater degree of skew   OE Lower median            OE	B1 B1 B1 B1		
<b>(c)</b>	People visiting the doctor may not have representative weights.	E1	<b>3</b>	Or similar in context
			<b>1</b>	
	<b>Total</b>		<b>13</b>	



<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>6(a)</b>	Generate random numbers in pairs	E1	<b>4</b>	ie 2-digit
	Rejecting any greater than 79 and any repeats.	E1		For either of these.
	Continue until 10 numbers obtained	E1		
	Include the corresponding councillors in the sample	E1		If candidate uses 01 to 80, must relate these numbers to the stated 00 to 79 to earn 4 <sup>th</sup> mark
<b>(b)</b>	09,63,20,71,78,11,19,13,54,26 (or 09,63,07,18,78,11,37,54,26,62)	B2	<b>2</b>	B2 for completely correct list of 10. B1 for 9 or 11 or one included erroneously.
<b>(c)(i)</b>	A systematic sample picks regularly throughout the list,...	E1	<b>2</b>	What a systematic sample does.
	but the list is alphabetical, not sorted by party.	E1		What the problem is
<b>(ii)</b>	$48/80 \times 10$ or $32/80 \times 10$ 6 Preservative and 4 Action.	M1 A1	<b>2</b>	Or B2 for 6 Preservative and 4 Action.
	<b>Total</b>		<b>10</b>	
	<b>TOTAL</b>		<b>75</b>	