

ADVANCED SUBSIDIARY (AS) General Certificate of Education 2013


# Biology

Assessment Unit AS 1

assessing Molecules and Cells

[AB111]

# **TUESDAY 11 JUNE, MORNING**



# TIME

1 hour 30 minutes.

# **INSTRUCTIONS TO CANDIDATES**

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Write your answers in the spaces provided in this question paper.

There is an extra lined page at the end of the paper if required.

Answer all eight questions.

You are provided with **Photograph 1.5** for use with Question 5 in this paper. Do not write your answers on this photograph.

# **INFORMATION FOR CANDIDATES**

The total mark for this paper is 75.

Section A carries 60 marks. Section B carries 15 marks.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You are reminded of the need for good English and clear presentation in your answers.

Use accurate scientific terminology in all answers.


You should spend approximately **20 minutes** on Section B.

You are expected to answer Section B in continuous prose.

Quality of written communication will be assessed in **Section B**, and awarded a maximum of 2 marks.

For Examiner's use only		
Question Number	Marks	
1		
2		
3		
4		
5		
6		
7		
8		
Total Marks		

8487




**BLANK PAGE** 

### **Section A**

- 1 Identify the word or phrase which is described by each of the following statements.
- StudentBounts.com The paired structures from which spindle fibres originate in animal cells
  - The precise part of a chromosome to which the spindle fibres attach
  - The stage of mitosis during which chromosomes lie across the equator of the spindle
  - The stage of mitosis when nuclear membranes form around the two groups of chromosomes
  - The precise stage of meiosis in which homologous chromosomes pair up to form bivalents

[5]

	agram shows how phospholipids and proteins are arranged in a face membrane.	Studente rony nark
	plain why the phospholipids form a bilayer in the cell-surface embrane.	
		_ [1]
the (i)	ese proteins act as carriers, enabling the membrane to be selection	ve.
	Explain how the structure of proteins allows them to select cert molecules and not others.	tain
(ii)		
(ii)	molecules and not others.	
(ii)	molecules and not others.	



All organic macromolecules contain the elements carbon, hydrogen and 3 oxygen. Some also contain a variety of other elements.

StudentBounty.com The following statements give information about the chemical composition and nature of five different macromolecules.

Molecules A and B contain only the elements carbon, hydrogen and oxygen.

- When hydrolysed, molecule A produces monomers that reduce Benedict's reagent.
- When hydrolysed, molecule **B** produces some sub-units that lower the pH of the solution.

Molecules C, D and E all contain other elements in addition to carbon, hydrogen and oxygen.

When hydrolysed, molecule **C** produces sub-units, all of which contain nitrogen and some also contain sulfur.


Molecules **D** and **E** both contain phosphate.

- When hydrolysed, molecule **D** creates two other types of sub-unit in addition to the phosphate. One type of sub-unit lowers the pH of the solution.
- When hydrolysed, molecule E creates four types of sub-unit. Each sub-unit has a nitrogen-containing molecule and a pentose, in addition to the phosphate.

Identify the macromolecules from the information that is given above.

Molecule A	
Molecule B	
Molecule C_	
Molecule D	
Molecule E	

[5]



# **BLANK PAGE**

(Questions continue overleaf)

StudentBounty.com Pumpkins, melons and cucumbers all belong to a family of plants (the 4 Cucurbitaceae) commonly known as gourds. As they are related species they would be expected to have many sections of their DNA in common. However, as they are different species, they would also be expected to have some unique sections of DNA.


Modern DNA technology allows scientists to extract DNA and to amplify specific relevant sections, such as Microsatellite Repeat Sequences (MRSs), that might show similarities or differences between the species. Following amplification, the relevant sections of DNA of each species can be separated and analysed.

(a) Outline how a scientist might amplify a specific section of DNA (e.g. containing a particular MRS).

\_ [3]

Twelve different sections containing MRSs were selected for	r Only park
amplification. Outline how the scientists could separate the selected amplified sections and then mark them to create a DNA fingerprint (profile) for analysis.	Rint Polynak
	19
[3]	
Describe how the scientists could now use the fingerprints (profiles) obtained from pumpkins, melons and cucumbers to show that these are related species and also to show that they are actually different species.	
Related species:	
Different species:	
[2]	


a)	Identify the structures labelled <b>A</b> and <b>B</b> .	SID &
	Α	2.0
	В	[2]
b)	Using the scale bar (on <b>photograph 1.5</b> ), calculate the magnificatio of this electronmicrograph. (Show your working.)	'n
	Answer	[3]
c)	Identify the precise stage of the cell cycle during which mitochondria would be most likely to divide. Explain why mitochondria divide at th stage.	
		[2]
d)	State the precise function of mitochondria in the cell.	[1]



[Turn over

StudentBounts.com (b) The Venus Flytrap plant (Dionaea muscipula) possesses modified leave two plates of a trap. When triggered, small invertebrates are caught and d mechanism of trap closure is believed to involve osmotic changes and the flo between the mesophyll layers within the plates.

A whole view of the plates is shown below, together with a cross-sectional view showing the mesophyll layers.



Examiner Only Marks Remark

[2]

The plate consists of two mesophyll layers, the upper and lower mesophyll. The water potentials of the cells of these layers is different and information regarding this is shown in the table below.

Potential/kPa	Cells of upper mesophyll	Cells of lower mesophyll
$\psi_{cell}$	0	
$\psi_{ m s}$	-250	-250
$\psi_{p}$		100

(i) Calculate the missing values and present these in the empty spaces in the table above.

> 12 www.StudentBounty.com iomework Help & P

StudentBounty.com When the trap is open, water movement between the mesophyll layers is not possible. However, when triggered to close, water will move from one layer to the other.

(ii) Determine the direction in which water will flow when the trap is triggered to close. Explain your answer.

\_\_\_\_\_ [2]

(iii) Explain what causes the change in the shape of the plates during closure of the trap.

\_\_\_\_\_ [2]

pr	nergy of a reaction. Two models of enzyme action have been roposed: the lock and key hypothesis and the induced fit model.	ion entre voly
(1)	Explain the term 'activation energy'.	
		[1]
(ii	<ul> <li>Describe one similarity and one difference between the lock an key hypothesis and the induced fit model of enzyme action.</li> <li>Similarity</li></ul>	
		_
	Difference	_
		[2]
le <sup>-</sup> th th	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found the cytoplasm, whereas its substrate catechol is usually confined to be vacuole. The reaction is a major cause of browning in fruits and egetables. The enzyme's action is summarised below.	nd
le th th ve	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found le cytoplasm, whereas its substrate catechol is usually confined to le vacuole. The reaction is a major cause of browning in fruits and	nd
le th th ve	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found is the cytoplasm, whereas its substrate catechol is usually confined to be vacuole. The reaction is a major cause of browning in fruits and egetables. The enzyme's action is summarised below.	n
le th th ve	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found is the cytoplasm, whereas its substrate catechol is usually confined to be vacuole. The reaction is a major cause of browning in fruits and egetables. The enzyme's action is summarised below. techol $\xrightarrow{catechol \ oxidase}$ Benzoquinone $\xrightarrow{non-enzymatic}{oxidation}$ Melanin (brown/black) Explain how dropping a banana or apple leads to the developm	n
le th th ve	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found is e cytoplasm, whereas its substrate catechol is usually confined to e vacuole. The reaction is a major cause of browning in fruits and egetables. The enzyme's action is summarised below. techol $\xrightarrow{catechol \ oxidase}$ Benzoquinone $\xrightarrow{non-enzymatic} \ oxidation}$ Melanin (brown/black) Explain how dropping a banana or apple leads to the developm of brown/black 'bruised' areas on the fruit.	n
le th th ve	he cells of many fruits and vegetables including apples, bananas a ttuce contain the enzyme catechol oxidase. The enzyme is found the cytoplasm, whereas its substrate catechol is usually confined to the vacuole. The reaction is a major cause of browning in fruits and egetables. The enzyme's action is summarised below. techol <u>catechol oxidase</u> Benzoquinone <u>non-enzymatic</u> <u>oxidation</u> Melanin (brown/black) Explain how dropping a banana or apple leads to the developm of brown/black 'bruised' areas on the fruit.	n

	(ii)	Suggest an explanation for the fact that lettuce leaves will brown faster if they are cut with a knife, rather than torn by hand.	Jon Bou	Only nark	OT
		[2]			
(c)	eac fund	e catechol oxidase molecule consists of four polypeptide chains, h of which is attached to a copper atom. The enzyme is not ctional without the copper atoms. State the term used to describe role of copper in the action of the enzyme.			
		[1]			
(d)		following are methods for decreasing the activity of catechol lase and thus preventing browning in fruits and vegetables:			
	1.	Cut fruit is coated with an antioxidant such as ascorbic acid, which prevents oxygen from reaching the cells of the fruit.			
	<b>2</b> .	Vegetables are blanched (immersed in boiling water for one minute) prior to preservation.			
	3.	Compounds such as cinnamic acid are added to fruit juice. Cinnamic acid is structurally similar to catechol.			
	Sele	ect the treatment 1, 2 or 3 which			
	•	represents an example of competitive inhibition			
	•	permanently changes the shape of the enzyme [2]			
(e)	oxic	vilage of fruit juice can also be prevented by removing the catechol lase enzyme. One way of doing this is to pour the juice through a umn containing a protease enzyme immobilised on glass beads.			
	the Onc dete	experiment was carried out to determine the optimum flow rate for juice through the column. A tap was used to adjust the flow rate. See the juice had been passed through the column, it was tested to ermine the concentration of catechol oxidase remaining. The ults of the investigation are shown in the table at the top of the next e.			


Flow rate/ mm <sup>3</sup> min <sup>−1</sup>	Concentration of catechol oxidase remaining in treated juice/arbitrary units
10	0
20	0
30	5
40	45
50	85
60	125
70	140

- (i) Plot the results using a suitable graphical technique. Use the graph paper opposite.
- (ii) It was concluded that the optimum flow rate for the juice through the column was 20 mm<sup>3</sup> min<sup>-1</sup>. Explain how this conclusion was reached.

\_\_\_ [2]

[4]


(iii) Describe and explain the trend shown by the results for flow rates of  $40-70 \text{ mm}^3 \text{ min}^{-1}$ .



### Section B

Quality of written communication is awarded a maximum of 2 marks in this section.

StudentBounty.com 8 Proteins have many roles in cells. For example, enzymes are involved in both extracellular and intracellular digestion, while conjugated proteins provide essential receptors and recognition sites on cell-surface membranes. Within the cell many organelles work together to make and transport these enzymes and conjugated proteins. The following diagram shows the relative positioning of these organelles.



Describe the involvement of each of the following:

- the nucleus and rough endoplasmic reticulum in the production of polypeptides (details of the process of polypeptide synthesis are **not** required)
- the Golgi body in the production of the functional enzymes and conjugated proteins
- vesicles and lysosomes in the transportation of products for secretion or for use within the cell. [13]

[2]

Quality of written communication


Ste.
 r Only nark
 - <b>Tony</b> <b>Park</b> - <b>Com</b>
 -
-
 -
-
 -
 -
 -
-
 -
 -
 -
-
 -
 -
-
_
-
 -


StudentBou	
 (Ches	Only nark
00	
	2
 	.01

Extra lined page	Studento	r Only
	- 18	oung Cung
		7.01



# THIS IS THE END OF THE QUESTION PAPER





Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified. GCE Biology Advanced Subsidiary (AS)

**Assessment Unit AS 1 Molecules and Cells** 

**Summer 2013** 



