Centre Number	Candidate Number	Name	
UNIVERS	General Cer	E INTERNATIONAL EXAMINATIONS rtificate of Education anced Level	
BIOLOGY	7.00	9700/05	
Paper 5 Prac	ctical Test A2		
		May/June 2005	
	wer on the Question Pap rials: As listed in Instruc		
READ THESE INSTRU			
Write in dark blue or bla You may use a soft pen			
If you have been given a details. If any details are missing, please fill in you in the space given at the	e incorrect or ur correct details	For Examiner's	s Use
Stick your personal labe		2	
provided.		Total	
		Iotai	
	This document consists of	f <b>7</b> printed pages and <b>1</b> blank page.	

1 You are required to investigate the light-dependent stage in photosynthesis in a leaf extract, using the dye DCPIP. When DCPIP is **reduced**, it changes from a **blue** colour, to **colourless**.

You are required to produce a leaf extract from material provided.

Procedure to make a leaf extract

- Place the piece of fresh leaf, **C1**, on a white tile.
- With a scalpel, carefully remove any large veins and discard them.
- Chop the rest of the leaf into small pieces.
- Place the chopped leaf into a plastic specimen tube.
- Using a syringe or pipette, add 2 cm<sup>3</sup> of cold buffer solution, labelled **C2**, to the specimen tube.
- Add a spatula full of sand, labelled **C3**. Carefully grind the chopped leaf in the specimen tube, using a glass rod for about one minute to obtain a green leaf extract.
- Clean the surface of the white tile with a paper towel.
- Place a Petri dish on the edge of the tile as shown in Fig. 1.1, so that the dish is tilted.
- Pour the leaf extract from the specimen tube into the Petri dish and allow it to drain to the edge of the dish as shown in Fig. 1.1.




Fig. 1.1

• Completely cover the Petri dish with aluminium foil to shield the contents from light. The foil should be made easy to remove.

## Leave the foil in position except when removing samples.

- Take a capillary tube and stand one end of the tube in the leaf extract. Some of the extract will immediately rise a short distance up the tube. Remove the tube and lay it on the tile as a colour standard. **Label this Tube 1**.
- You are provided with a 1% solution of DCPIP, labelled **C4**. Using a teat pipette, add five drops of **C4** solution to the leaf extract. Tilt the dish to mix the two liquids (or mix with the teat pipette).

## Ensure that the foil cover is replaced.

• Take a second capillary tube and collect some leaf extract/C4 mixture. Immediately replace the foil cover and wrap the tube quickly in aluminium foil to prevent any exposure to light. Lay the tube on the tile.

3

## Label this Tube 2.

• Take a third capillary tube and collect some leaf extract/C4 mixture. Place the tube near a bench lamp.

## Label this Tube 3.

- Carefully check the colour of the liquid in all of the tubes, every minute, ensuring that the foil on Tube 2, is replaced each time.
- Note the time taken for the dye in Tubes 2 and 3 to be reduced.
- If there is no change in colour after 10 minutes, stop the investigation.
- (a) Draw a table to show your results.

[4]


(b) Explain your results in relation to the light-dependent stage of photosynthesis.

(c) Suggest why C2 and C4 were kept in a beaker of ice so that the solutions were cold when they were being used.

(d)	Explain the purpose of using the buffer solution, C2.		
	[1]		
(e)	The buffer solution also contained some sucrose. Suggest why sucrose was added to the buffer solution.		

(f) Suggest how the investigation could be improved to make the results more reliable.

..... ..... ..... .....[3] [Total : 15]



5



(a) Determine the ratio of red blood cells to white blood cells.


ratio	 [2]	

- (b) You are provided with a slide of human blood, labelled **K1**. Choose an appropriate magnification so that you can clearly see both red and white blood cells in your field of view. The white blood cells have been stained blue for easy identification.
  - (i) Explain why it would be difficult to determine the exact ratio of red blood cells to white cells in slide **K1**.

(ii) Comment on the ratio of red blood cells to white blood cells when compared with the ratio you determined in (a).

(c) Fig. 2.2 is the same photomicrograph of blood, as is shown in Fig. 2.1.

6





In the spaces below, draw to the same scale, labelled diagrams of:

(i) a red blood cell.

(ii) a phagocyte.

(iii) a lymphocyte.

		[8]
(iv)	Draw a line, on Fig. 2.2, across the lymphocyte.	
	The actual diameter of a red blood cell is $8\mu$ m.	
	Use this information to calculate the width of the lymphocyte along the line draw	vn.

width of lymphocyte ......[2]

[Total : 15]

**BLANK PAGE** 

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.