



GCE AS/A LEVEL

2400U10-1 – NEW AS

BIOLOGY – Unit 1

Basic Biochemistry and Cell Organisation

P.M. THURSDAY, 26 May 2016

1 hour 30 minutes plus your additional time allowance

Surname _____

Other Names _____

Centre Number _____

Candidate Number 2 _____

| For Examiner's use only | | |
|--------------------------------|---------------------|---------------------|
| Question | Maximum Mark | Mark Awarded |
| 1. | 8 | |
| 2. | 16 | |
| 3. | 11 | |
| 4. | 19 | |
| 5. | 17 | |
| 6. | 9 | |
| Total | 80 | |

ADDITIONAL MATERIALS

In addition to this paper you will require a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink, black ball-point pen or your usual method.

Write your name, centre number and candidate number in the spaces on the front cover.


Answer ALL questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation pages at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The assessment of the quality of extended response (QER) will take place in question 6.



Answer ALL questions.

1. In 1951, Henrietta Lacks died of cervical cancer. A research scientist called George Gey, grew a sample of her tumour. He found that these cells multiplied rapidly and could be grown indefinitely in culture. They became the first immortal human cell line, which he named HeLa. HeLa cells are now grown in research laboratories in many different countries.
- (a) The photomicrographs opposite show images of HeLa cells during different stages of the cell cycle. The cells have been stained with a dye, which causes the DNA to be visible.

(i) Name the stages shown in **A** and **B**. [2]

Stage **A**: _____

Stage **B**: _____


1(a) (ii) Name the structures, labelled X, that can be seen in the photomicrograph. [1]

(iii) Explain why some stain would be seen in other parts of the cell. [2]

- 1(b) A biotechnology company that supplies HeLa cells to laboratories states that the number of HeLa cells double every 19 hours. They suggest that the starting culture of cells should have a density of 30 000 cells cm^{-3} and that the cells should be sub-cultured every 4 days.

Calculate the density of the cells that would be present after 4 days; give your answer in cells cm^{-3} in standard form. [3]

Cell density = _____ cm^{-3}



3. The photomicrograph opposite is of **SPIROGYRA**; an autotrophic organism that inhabits fresh water ponds and ditches.

(a) Identify the organelles labelled **A** and **B**. [2]

A: _____


B: _____

(b) The actual width of the cell between points **X – Y** was $32.3\ \mu\text{m}$. Calculate the magnification that was used to take the photomicrograph. [2]

magnification = _____


**photosynthetic
membrane**

**peptidoglycan
cell wall**



3(d) Opposite is a photomicrograph of the bacteria NOSTOC, which also inhabits fresh water ponds and ditches. They were once thought to belong to the same group of organisms as SPIROGYRA and thought to have the same cellular structure. Evidence from electron microscopy has now grouped these two separately.

Conclude what cell types are present in SPIROGYRA and NOSTOC. Identify TWO differences and ONE similarity (not labelled in the photomicrographs) between these two species that would be revealed by electron microscopy. [3]



4. Sucrose is a disaccharide of glucose and fructose. The enzyme sucrase, catalyses the hydrolysis of sucrose into its monosaccharides. A colorimeter can be used to record absorbance values, which can then be used to determine the rate of hydrolysis.

(a) (i) Complete the diagram opposite to show the hydrolysis of sucrose including the products formed. [2]

(ii) State the name of the bond broken during the reaction. [1]

(iii) Explain why glucose and fructose are referred to as structural isomers. [1]

4(a) (iv) A student was provided with two beakers, one containing sucrose and the other containing sucrose. Describe ONE biochemical test that the student could have carried out to distinguish between the two solutions. [2]

4(b) A student wanted to investigate the hypothesis that sucrase catalyses the hydrolysis of sucrose fastest at a neutral pH. She was provided with the following method:

- **Add 5 cm³ of buffer solution to a test tube**
- **Add 5 cm³ of sucrose solution to the test tube**
- **Add 1 cm³ of sucrase solution to the test tube and mix**
- **After 20 minutes add 1 cm³ of dinitrosalicylic acid (DNS)**
- **Pipette 5 cm³ of the mixture into a cuvette and place into a colorimeter and record the absorbance of light passing through the solution.**


Dinitrosalicylic acid (DNS) will react with monosaccharides to produce amino-nitrosalicylic acid (ANS). ANS causes a colour change to occur which can be detected by a colorimeter. The greater the concentration of ANS the greater the absorbance of light.


4(b) (i) State TWO additional variables that the student should control to ensure that the results recorded would be repeatable. [2]

4(b) (ii) The results are shown in the table below.

| pH OF SUCROSE AND SUCRASE SOLUTION | ABSORBANCE OF LIGHT (au) | | | |
|------------------------------------|--------------------------|---------|---------|------|
| | TRIAL 1 | TRIAL 2 | TRIAL 3 | MEAN |
| 3 | 0.84 | 0.63 | 0.76 | 0.74 |
| 4 | 0.89 | 0.95 | 0.76 | 0.87 |
| 5 | 0.85 | 0.91 | 0.72 | 0.83 |
| 7 | 0.27 | 0.34 | 0.29 | 0.30 |
| 9 | 0.13 | 0.11 | 0.12 | 0.12 |
| 10 | 0.05 | 0.04 | 0.06 | 0.05 |

Plot the mean data on the graph opposite. [4]






4(b) (iv) The student was also provided with the calibration curve (shown opposite) in order to calculate the concentration of glucose produced during the investigation.


The colour change, caused by ANS, is dependent on the concentration of monosaccharides, the greater the concentration the greater the absorbance value recorded.

Determine the maximum concentration of glucose produced during the investigation at pH 7. [2]


Epithelial cell




protein molecules




**double
layer of fat
(phospholipid)
molecules**




Cell 1




Cell 2




5(d) Acidification of fresh-water lakes, due to acid rain, has been linked to the death of fish such as carp. Scientists concluded that one of the causes of death in these animals is their inability to maintain blood plasma ion concentrations. Use the information to explain how they arrived at this conclusion. [4]




pyrophosphatase



pyrophosphate



phosphate



phenylalanine

6. **Pyrophosphatase is an enzyme found inside the nucleus of cells and is involved in DNA replication. The enzyme catalyses the conversion of a molecule of pyrophosphate to two phosphate ions. The diagrams opposite show the enzyme pyrophosphatase and its substrate pyrophosphate. Molecules of phenylalanine (an amino acid) and phosphate are also shown; both of these molecules are known to inhibit pyrophosphatase. (Drawings are not to the same scale).**

Describe and explain why pyrophosphatase can only hydrolyse pyrophosphate and the mechanism by which phenylalanine and phosphate inhibit pyrophosphatase. [9 QER]
