

ADVANCED SUBSIDIARY (AS) General Certificate of Education 2014


# Chemistry

Assessment Unit AS 2

assessing Module 2: Organic, Physical and Inorganic Chemistry

[AC122]

**TUESDAY 17 JUNE, AFTERNOON** 





TIME

1 hour 30 minutes.

## INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Answer **all seventeen** questions.

Answer **all ten** questions in **Section A**. Record your answers by marking the appropriate letter on the answer sheet provided. Use only the spaces numbered 1 to 10. Keep in sequence when answering. Answer **all seven** questions in **Section B**. Write your answers in the spaces provided in this question paper.

### INFORMATION FOR CANDIDATES

The total mark for this paper is 100.

Quality of written communication will be assessed in Question **13(b)(ii)**.

In Section A all questions carry equal marks, i.e. **two** marks for each question.

In Section B the figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question. A Periodic Table of the Elements, containing some data, is included in this question paper.

	miner's only
Question Number	Marks
Sect	ion A
1–10	
Sect	ion B
11	
12	
13	
14	
15	
16	
17	
Total Marks	

9575

#### Section A

For each of the following questions only **one** of the lettered responses (A–D) is correct.

# StudentBounts.com Select the correct response in each case and mark its code letter by connecting the dots as illustrated on the answer sheet.

- 1 Equal volumes of 1-chlorobutane and 1-iodobutane are warmed with aqueous silver nitrate in the presence of ethanol. Which one of the following is the reason why the 1-chlorobutane reacts more slowly?
  - А The C–CI bond is more polar than the C–I bond
  - The C–CI bond is stronger than the C–I bond В
  - С The C-I bond is more polar than the C-CI bond
  - The C–I bond is stronger than the C–CI bond D
- Which one of the following is correct as Group II is descended? 2

	Solubility of hydroxides	Solubility of sulfates
А	decreases	decreases
В	decreases	increases
С	increases	decreases
D	increases	increases

- Which one of the following is the colour of the flame produced when a barium compound is 3 placed in a blue Bunsen burner flame?
  - Crimson А
  - В Green
  - С Lilac
  - Orange D

StudentBounty.com Which one of the following shows the effect on the yield of ammonia in the Haber prov 4

#### Pressure increase

А yield decreases

- В vield decreases
- С yield increases

- **Temperature increase** yield decreases
  - vield increases
  - yield decreases
- D yield increases
- yield increases
- 5 Which one of the following mixtures will react to produce a compound with molecular formula C<sub>4</sub>H<sub>7</sub>N?
  - А 1-bromobutane and ammonia
  - В 1-bromobutane and potassium cyanide
  - 1-bromopropane and ammonia С
  - D 1-bromopropane and potassium cyanide
- The reaction shown below 6


 $\mathrm{C_2H_5Br} + \mathrm{KOH} \rightarrow \mathrm{C_2H_4} + \mathrm{KBr} + \mathrm{H_2O}$ 

is an example of

- dehydration. А
- В elimination.
- С free radical substitution.
- nucleophilic substitution. D

- StudentBounts.com Which one of the following is the mass of magnesium required to react with  $50.0 \,\mathrm{cm^3}$ 7 0.1 mol dm<sup>-3</sup> hydrochloric acid?
  - А 0.005g
  - В 0.060 g
  - С 0.120 g
  - D 0.240 g
- 8 Which one of the following lists the compounds in order of increasing boiling point?
  - A CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub> CH<sub>3</sub>CH<sub>2</sub>F CH<sub>3</sub>CH<sub>2</sub>OH B CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub> CH<sub>3</sub>CH<sub>2</sub>OH CH<sub>3</sub>CH<sub>2</sub>F C CH<sub>3</sub>CH<sub>2</sub>F CH<sub>3</sub>CH<sub>2</sub>OH CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub>
  - D CH<sub>3</sub>CH<sub>2</sub>OH CH<sub>3</sub>CH<sub>2</sub>F CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub>
- When an organic compound was reacted with chlorine, the organic product was found to 9 have a relative molecular mass which had increased by 69. Which one of the following is the reaction mechanism?
  - elimination А
  - electrophilic addition В
  - free radical substitution С
  - nucleophilic substitution D

**10** The Maxwell–Boltzmann distribution for a reaction mixture is shown below. N is the not of molecules with the most probable energy and E<sub>A</sub> is the activation energy.




Which one of the following shows the effect on  $\mathsf{E}_\mathsf{A}$  and on N of increasing the temperature?


	E <sub>A</sub>	Ν
А	constant	decreases
В	constant	increases
С	decreases	decreases
D	decreases	increases


			Section B	g
			Answer all <b>seven</b> questions in the spaces provided.	
11		an-1 Iatior	-ol is used to prepare 1-bromobutane according to the followin	g
			$C_4H_9OH + HBr \rightarrow C_4H_9Br + H_2O$	
	(a)	(i)	Give an equation to explain the term <b>percentage yield</b> .	
			percentage yield =	
				[1]
		(ii)	Assuming a 40% yield, what mass of butan-1-ol would be required to produce 5.48g of 1-bromobutane?	
			moles of 1-bromobutane	
			moles of butan-1-ol	
			mass of butan-1-ol	[0]
	(b)	(i)	Give an equation to explain the term <b>atom economy</b> .	_ [3]
			atom economy =	
				[1]
		(ii)	Calculate the atom economy for the formation of 1-bromobuta from butan-1-ol.	ine
				_ [1]

			Still	
		ive analysis can be used to distinguish between aqueous solutionng different metal ions.	DNS Examin Marks	NHB OL
(a)	whie	each of the following pairs of metal ions, give an aqueous reage ch can be used to distinguish between the aqueous solutions ar e the expected observations for each ion.	ent nd	
	(i)			
		Reagent	[1]	
		Observations		
	(ii)	Aluminium ions and zinc ions.		
		Reagent	[1]	
			[2]	
(b)			d to	
	(i)	What is observed when barium ions react with chromate ions?		
			[1]	
	(ii)	Write an ionic equation, including state symbols, for the reaction of barium ions with chromate ions.	n	
			[2]	
		7		n over
	con (a)	<ul> <li>(a) For white stat</li> <li>(i)</li> <li>(ii)</li> <li>(b) Add test</li> <li>(i)</li> </ul>	<ul> <li>(i) Iron(II) ions and iron(III) ions.</li> <li>Reagent</li></ul>	<ul> <li>(a) For each of the following pairs of metal ions, give an aqueous reagent which can be used to distinguish between the aqueous solutions and state the expected observations for each ion.</li> <li>(i) Iron(II) ions and iron(III) ions. <ul> <li>Reagent</li> <li>[1]</li> <li>Observations</li> <li>[2]</li> </ul> </li> <li>(ii) Aluminium ions and zinc ions. <ul> <li>Reagent</li> <li>[1]</li> <li>Observations</li> <li>[1]</li> <li>Observations</li> <li>[2]</li> </ul> </li> <li>(ii) Aluminium ions and zinc ions. <ul> <li>Reagent</li> <li>[1]</li> <li>Observations</li> <li>[2]</li> </ul> </li> <li>(ii) Aluminium ions and zinc ions. <ul> <li>Reagent</li> <li>[1]</li> <li>Observations</li> <li>[2]</li> </ul> </li> <li>(b) Addition of an aqueous solution of potassium chromate can be used to test for the presence of barium ions. <ul> <li>(i) What is observed when barium ions react with chromate ions?</li> <li>[1]</li> <li>(ii) Write an ionic equation, including state symbols, for the reaction of barium ions with chromate ions.</li> </ul></li></ul>

$CH_3CH_2CH = CFCH_2CH_3$ his fluorohydrocarbon using IUPAC rules. w and label the structures of the E and Z isomers of this rohydrocarbon.	Examin Marks Re.
w and label the structures of the E and Z isomers of this	
	_ [2]
	[1]
lain why one of the structures you have drawn is classified Z isomer.	d as
	_ [3]
ality of written communication	[2]
	Plain why one of the structures you have drawn is classified Z isomer.








		particular set of conditions incomplete combustion of propane according to the following equation:	he
		$C_3H_8(g) + 4O_2(g) \rightarrow CO_2(g) + 2CO(g) + 4H_2O(I)$	
		I volume of gases produced occupied a volume of 9000 dm <sup>3</sup> . Thas volume, under these conditions, is 30 dm <sup>3</sup> .	he
a)	Defi	ne the term <b>molar gas volume</b> .	
			[2]
b)	(i)	Calculate the number of moles of carbon monoxide produced in this combustion.	n
			[2]
	/!!)		
	(11)	Calculate the number of moles of oxygen used in this combust	ion.
	(11)	Calculate the number of moles of oxygen used in this combust	ion. [1]
	-	Calculate the number of moles of oxygen used in this combusti	
	-		[1]
	(iii)	Calculate the mass of propane, in kg, burned.	[1]
	(iii)	Calculate the mass of propane, in kg, burned.	[1]
(c)	(iii) (iv) Und	Calculate the mass of propane, in kg, burned.	. [1]

-

					Stur
16				ally as "bottled gas". It undergoes complete the following equation:	Examin Marks Re.
			-	$_{10} + 13O_2 \rightarrow 8CO_2 + 10H_2O$	12.00
	and	d has		Ipy of combustion of $-2876.5 \text{ kJ mol}^{-1}$ .	13
		Cal		t of energy released for every kilogram of carbon	
				[3]	
	(b)	star		sed to calculate enthalpy changes, such as the formation of butane, which cannot be measured	
		(i)	Define the term s	standard enthalpy of formation.	
				[3]	
		(ii)		n, with state symbols, which represents the y of formation of butane. [2]	
		(iii)		l enthalpy of combustion of butane and the calculate the standard enthalpy of formation of	
				Standard enthalpy of combustion (kJ mol <sup>-1</sup> )	
			Carbon (C)	-393.5	
			Hydrogen (H <sub>2</sub> )	-285.8	
				[3]	
9575				13	[Turn over

		Stud
I <b>7</b> Th	e formula for ethylene glycol is shown below:	Examin Marks Re
	CH <sub>2</sub> OH	Marks Re.
	│ <sup>∠</sup> CH₂OH	.6
(2)	Use IUPAC rules to give the systematic name of ethylene glycol.	
(a)		[4]
		[1]
(b)	What is the empirical formula of ethylene glycol?	
		[1]
(c)	Ethylene glycol reacts vigorously with an <i>excess</i> of ethanoyl chloric	le.
	(i) Suggest <b>two</b> observations in this reaction.	
		[2]
	(ii) Name the type of reaction occurring.	
	(ii) Name the type of reaction occurring.	[1]
		[1]
	(iii) Draw the structure of the organic product.	
		[2]
	(iv) Suggest a test for the inorganic product formed in this reaction	
		[2]
9575	14	

-	reflu	vlene glycol contains primary alcohol groups. When heated unc ux with excess acidified potassium dichromate the solution nges from orange to green.		Examin. Marks	Re. C	32.0
		Why are the alcohol groups in ethylene glycol classified as primary?				
			[1]			
	(ii)	Name the type of reaction occurring.				
			[1]			
	(iii)	Draw the structure of the organic product.				
			[1]			
	(iv)	Name the functional group present in the organic product.				
			[1]			
		What would be the most significant difference between the infra-red spectrum of the organic product and that of ethylene glycol?				
			[1]			
		e an equation for the reaction of ethylene glycol with an excess sphorus pentachloride.	s of			
			[2]			











Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.