

SECTION A

Answer ALL parts of this question in the spaces provided.

1. The following procedure can be used to prepare ethyl ethanoate (boiling temperature 77°C).

- Mix 20 cm³ of ethanol (an excess) and 12.6 g of ethanoic acid in a pear-shaped flask.
- Slowly add 8 cm³ of concentrated sulphuric acid, with cooling and mixing.
- Heat the mixture under reflux for 15 minutes.
- Allow the apparatus to cool and then re-arrange it for distillation.
- Collect everything that distils up to 80°C.
- Purify the distillate.

(a) Write the equation for the reaction between ethanol and ethanoic acid.

.....
.....

(1)

(b) What is the purpose of the concentrated sulphuric acid in this reaction?

.....
.....

(1)



(c) Draw a diagram to show how you should set up the apparatus to distil the mixture.

Leave
blank

(3)



(d) 10.6 g of pure ethyl ethanoate was collected.

Calculate the percentage yield of ethyl ethanoate obtained in this experiment.

(3)

(e) Ethyl ethanoate can also be prepared from ethanol and ethanoyl chloride.

(i) Write the equation for this reaction.

.....
(1)

(ii) Explain why the yield obtained by this method is significantly higher than that from ethanoic acid.

.....
.....
.....
(1)



Leave
blank

(f) The solid ester phenyl benzoate can be prepared from phenol and benzoyl chloride. It is purified by recrystallisation.

(i) Explain what makes a solvent suitable for use in this recrystallisation.

.....
.....
(1)

(ii) Explain why the mixture is filtered after dissolving the phenyl benzoate in the minimum amount of hot solvent.

.....
.....
(1)

(iii) Explain why the mixture is filtered after cooling.

.....
.....
(1)

(iv) Explain why the residue is washed with a small amount of cold solvent.

.....
.....
(1)

(v) State how you would check that the sample after recrystallisation was pure.

.....
.....
(2)

Q1

(Total 16 marks)

TOTAL FOR SECTION A: 16 MARKS



Leave
blank

SECTION B

Answer any TWO questions from this section in the spaces provided.

If you answer Question 2 put a cross in this box .

2. (a) Describe the structure of solid sodium chloride.

.....
.....
.....
.....

(2)

- (b) Explain, in terms of the bonding in solid sodium chloride and in the solution, why sodium chloride is soluble in water.

.....
.....
.....
.....
.....

(2)



Leave
blank

(c) Draw a labelled Hess's Law cycle for the dissolving of sodium chloride in water.

Use it and the data below to calculate the enthalpy of solution of sodium chloride.

Lattice enthalpy of NaCl	-771 kJ mol^{-1}
Hydration enthalpy of Na^+	-406 kJ mol^{-1}
Hydration enthalpy of Cl^-	-364 kJ mol^{-1}

(3)

QUESTION 2 CONTINUES ON THE NEXT PAGE



7

Turn over

Leave
blank

(d) Bromine is produced during the electrolysis of sodium bromide solution.

Identify the other **two** products of this electrolysis and write the half-equation for the reaction occurring at the anode.

.....
.....
.....
.....

(3)

(e) (i) Write the mechanism for the addition of bromine to ethene.

(3)



Leave
blank

(ii) When ethene reacts with a mixture of bromine and sodium chloride, 1,2-dibromoethane and 1-bromo-2-chloroethane are formed but no 1,2-dichloroethane.

Explain why some 1-bromo-2-chloroethane is formed but no 1,2-dichloroethane.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4)

Q2

(Total 17 marks)

--	--



If you answer Question 3 put a cross in this box .

3. (a) The table below shows the results of a kinetic investigation into the alkaline hydrolysis of 2-chloro-2-methylpropane, $(\text{CH}_3)_3\text{CCl}$.

Experiment	$[(\text{CH}_3)_3\text{CCl}]$ / mol dm^{-3}	$[\text{OH}^-]$ / mol dm^{-3}	Relative Initial Rate
A	0.2	0.1	1.0
B	0.3	0.1	1.5
C	0.1	0.2	0.5

- (i) Deduce the orders of reaction with respect to each of OH^- and 2-chloro-2-methylpropane, showing your working.

.....

.....

.....

.....

.....


(3)

- (ii) Use the orders that you have deduced in (i) to give the mechanism for the reaction of 2-chloro-2-methylpropane with aqueous hydroxide ions.

(3)



(b) Citronellal is the insect repellent found in a citronella candle.



(i) Identify the chiral carbon atom with an asterisk (*) on the diagram above.

(1)

(ii) Describe simple test tube experiments that would enable you to identify the **two** functional groups in citronellal.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4)

QUESTION 3 CONTINUES ON THE NEXT PAGE



Leave
blank

(iii) Draw the formulae of the major organic products formed when citronellal reacts with

sodium tetrahydridoborate(III), NaBH_4 , in aqueous ethanol

hydrogen bromide

(2)



Leave
blank

(iv) Give the equation for the complete combustion of citronellal, $C_{10}H_{18}O$.

Calculate the maximum volume, measured at room temperature and pressure, of carbon dioxide that could be formed by the complete combustion of 1.0 g of citronellal.

[Molar mass of citronellal = 154 g mol^{-1} and the molar volume of gas at room temperature and pressure = $24 \text{ dm}^3 \text{ mol}^{-1}$]

(4)

Q3

(Total 17 marks)

--	--

13



Turn over

Leave
blank

(b) In the solid state, phosphorus pentachloride contains the ions PCl_4^+ and PCl_6^- .

Suggest the shapes of these ions and justify the shape of ONE of the ions.

.....
.....
.....
.....
.....
.....
.....
.....

(3)

(c) Name a specific organic compound that has a functional group which reacts with phosphorus pentachloride and write the equation for the reaction.

.....
.....
.....

(2)

QUESTION 4 CONTINUES ON THE NEXT PAGE



Leave
blank

(d) Phosphorus pentachloride dissociates as shown in the reversible reaction



At 420 K and a pressure of 4.0 atm, phosphorus pentachloride was 67% dissociated.


Calculate K_p at this temperature and state its units.

(5)



(e) Phosphorus pentachloride reacts with water to produce phosphoric(V) acid, H_3PO_4 .

The titration curve when $0.100 \text{ mol dm}^{-3}$ of sodium hydroxide is added to 25.0 cm^3 of $0.100 \text{ mol dm}^{-3}$ of phosphoric(V) acid is shown below.



(i) Write the equation for the reaction that has occurred when 50.0 cm^3 of sodium hydroxide solution has been added.

.....

 (1)

(ii) Mark a point on the curve with a cross (x) where a buffer solution would be present.

(1)

Q4

(Total 17 marks)

TOTAL FOR SECTION B: 34 MARKS

TOTAL FOR PAPER: 50 MARKS

END



BLANK PAGE



BLANK PAGE



N 2 9 2 6 4 A 0 1 9 2 0

THE PERIODIC TABLE

Period	1	2	Group										3	4	5	6	7	0		
1	¹ H Hydrogen 1																			⁴ He Helium 2
2	⁷ Li Lithium 3	⁹ Be Beryllium 4																		²⁰ Ne Neon 10
3	²³ Na Sodium 11	²⁴ Mg Magnesium 12																		^{35.5} Cl Chlorine 17
4	³⁹ K Potassium 19	⁴⁰ Ca Calcium 20	⁴⁵ Sc Scandium 21	⁵¹ V Vanadium 23	⁵² Cr Chromium 24	⁵⁵ Mn Manganese 25	⁵⁶ Fe Iron 26	⁵⁹ Co Cobalt 27	⁵⁹ Ni Nickel 28	^{63.5} Cu Copper 29	^{65.4} Zn Zinc 30	⁷³ Ga Gallium 31	⁷⁵ Ge Germanium 32	⁷⁹ Se Selenium 34	⁸⁰ Br Bromine 35	⁸⁴ Kr Krypton 36				⁸⁴ Kr Krypton 36
5	⁸⁵ Rb Rubidium 37	⁸⁸ Sr Strontium 38	⁸⁹ Y Yttrium 39	⁹³ Nb Niobium 41	⁹⁶ Mo Molybdenum 42	⁹⁹ Tc Technetium 43	¹⁰¹ Ru Ruthenium 44	¹⁰³ Rh Rhodium 45	¹⁰⁶ Pd Palladium 46	¹⁰⁸ Ag Silver 47	¹¹² Cd Cadmium 48	¹¹⁵ In Indium 49	¹¹⁹ Sn Tin 50	¹²² Sb Antimony 51	¹²⁷ I Iodine 53	¹³¹ Xe Xenon 54				²²² Rn Radon 86
6	¹³³ Cs Caesium 55	¹³⁷ Ba Barium 56	¹³⁹ La Lanthanum 57	¹⁸¹ Ta Tantalum 73	¹⁸⁴ W Tungsten 74	¹⁸⁶ Re Rhenium 75	¹⁹⁰ Os Osmium 76	¹⁹² Ir Iridium 77	¹⁹⁵ Pt Platinum 78	¹⁹⁷ Au Gold 79	²⁰¹ Hg Mercury 80	²⁰⁴ Tl Thallium 81	²⁰⁷ Pb Lead 82	²⁰⁹ Bi Bismuth 83	²¹⁰ Po Polonium 84	²¹⁰ At Astatine 85				⁸⁶ Rn Radon 86
7	²²³ Fr Francium 87	²²⁶ Ra Radium 88	²²⁷ Ac Actinium 89	¹⁴¹ Pr Praseodymium 59	¹⁴⁴ Nd Neodymium 60	⁽¹⁴⁷⁾ Pm Promethium 61	¹⁵⁰ Sm Samarium 62	¹⁵² Eu Europium 63	¹⁵⁷ Gd Gadolinium 64	¹⁵⁹ Tb Terbium 65	¹⁶³ Dy Dysprosium 66	¹⁶⁵ Ho Holmium 67	¹⁶⁷ Er Erbium 68	¹⁶⁹ Tm Thulium 69	¹⁷³ Yb Ytterbium 70	¹⁷⁵ Lu Lutetium 71				²⁵⁷ Lr Lawrencium 103
	²³² Th Thorium 90	⁽²³¹⁾ Pa Protactinium 91	²³⁸ U Uranium 92	⁽²³⁷⁾ Np Neptunium 93	⁽²⁴²⁾ Pu Plutonium 94	⁽²⁴³⁾ Am Americium 95	⁽²⁴⁷⁾ Cm Curium 96	⁽²⁵¹⁾ Bk Berkelium 97	⁽²⁵¹⁾ Cf Californium 98	⁽²⁵³⁾ Es Einsteinium 99	⁽²⁵³⁾ Fm Fermium 100	⁽²⁵⁵⁾ Md Mendelevium 101	⁽²⁵⁶⁾ No Nobelium 102	⁽²⁵⁷⁾ Lr Lawrencium 103						

Key

Molar mass g mol⁻¹
Symbol
Name
Atomic number

¹
H
Hydrogen
1

⁴
He
Helium
2

