

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

Monday 19 June 2023 – Afternoon

A Level Chemistry A

H432/02 Synthesis and analytical techniques

**Time allowed: 2 hours 15 minutes
plus your additional time allowance**

YOU MUST HAVE:
the Data Sheet for Chemistry A

YOU CAN USE:
a scientific or graphical calculator
an HB pencil

Please write clearly in black ink.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

First name(s) _____

Last name _____

READ INSTRUCTIONS OVERLEAF



INSTRUCTIONS

Use black ink. You can use an HB pencil, but only for graphs and diagrams.

Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.

Answer ALL the questions.

Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

The total mark for this paper is 100.

The marks for each question are shown in brackets [].

Quality of extended response will be assessed in questions marked with an asterisk (*).

ADVICE

Read each question carefully before you start your answer.

BLANK PAGE

SECTION A

You should spend a **MAXIMUM** of 20 MINUTES plus your additional time allowance on this section.


Write your answer to each question in the box provided.

1 Which compound is used for proton exchange in NMR spectroscopy? [1]



Your answer

2 Which compound reacts with 2,4-dinitrophenylhydrazine but does NOT react with Tollens' reagent? [1]



Your answer

- 3 Propyne, $\text{CH}_3\text{C}\equiv\text{CH}$, is a member of the alkynes homologous series with the $\text{C}\equiv\text{C}$ functional group.


What is the general formula of the alkynes? [1]



Your answer

- 4 The structure of the painkiller paracetamol is shown below.

PARACETAMOL



Which functional groups are present in paracetamol? [1]


- A alcohol, amide
- B alcohol, arene, ketone, amine
- C phenol, amide
- D phenol, ketone, amine

Your answer

☐

- 5 Oxymetazoline, shown below, is used as a decongestant in the treatment of colds.

OXYMETAZOLINE



How many H atoms are in one molecule of oxymetazoline? [1]

- A 23
B 24
C 25
D 26


Your answer

6 Which statement supports the delocalised model of benzene and NOT the Kekulé model? [1]

- A Sigma bonds overlap to form a π -system.**
- B The carbon-carbon bond lengths are all the same.**
- C The enthalpy change of hydrogenation is more exothermic than expected.**
- D Benzene is more reactive than alkenes with bromine.**

Your answer ☐

7 What is the systematic name for the compound below? [1]



- A 3,5-dimethylocta-1,3,6-triene**
- B 3,5-dimethylocta-2,5,7-triene**
- C 4,6-dimethylocta-1,3,6-triene**
- D 4,6-dimethylocta-2,5,7-triene**

Your answer ☐





- 8 For complete combustion, 0.100 mol of an alkane requires 22.8 dm³ of O₂, measured at RTP.

Which alkane has undergone complete combustion? [1]

- A pentane
- B hexane
- C heptane
- D octane

Your answer

9 Which compound has the greatest number of peaks in its proton NMR spectrum? [1]

A	
B	
C	
D	

Your answer





10 Which ester is most likely to produce a mass spectrum with a fragment ion at $m/z = 43$? [1]

A	$ \begin{array}{c} \text{OH} \quad \text{O} \\ \quad \\ \text{H}_3\text{C}-\text{C}-\text{C}-\text{O}-\text{CH}_3 \\ \\ \text{CH}_3 \end{array} $
B	$ \text{HO}-\text{CH}_2-\overset{\text{O}}{\overset{ }{\text{C}}}-\text{O}-\text{CH}_2\text{CH}_2\text{CH}_3 $
C	$ \begin{array}{c} \text{OH} \quad \text{O} \\ \quad \\ \text{H}_3\text{C}-\text{CH}-\text{CH}_2-\text{C}-\text{O}-\text{CH}_3 \end{array} $
D	$ \begin{array}{c} \text{OH} \quad \text{O} \\ \quad \\ \text{CH}_3\text{CH}_2-\text{C}-\text{C}-\text{O}-\text{CH}_2\text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array} $


Your answer

11 The infrared spectrum of an organic compound is shown opposite.





Which compound could have produced this spectrum? [1]

A	
B	
C	
D	

Your answer




12 Which compound reacts with ethanoyl chloride? [1]

A	
B	
C	
D	


Your answer

13 Which compound(s) is/are aliphatic? [1]


1



2



3



A 1, 2 and 3

B Only 1 and 2


C Only 2 and 3

D Only 1

Your answer

☐

14 Which compound(s) is/are hydrolysed by HCl(aq) to produce butanoic acid? [1]



A 1, 2 and 3



B Only 1 and 2

C Only 2 and 3

D Only 1

Your answer

15 Which ion(s) contain(s) bond angles of approximately 120°? [1]



A 1, 2 and 3

B Only 1 and 2

C Only 2 and 3

D Only 1

Your answer

BLANK PAGE

SECTION B

16 This question is about hydrocarbons.

- (a)** The boiling points of some hydrocarbons containing 6 carbon atoms are shown below.


Hydrocarbon	Boiling point/°C
2,2-dimethylbutane	50
2-methylpentane	60
hexane	69

State and explain the trend in boiling points shown by these hydrocarbons.

[4]

- (b) 2-methylpentane reacts with bromine by radical substitution.

2-METHYLPENTANE



A mixture of organic products is formed, including 3-bromo-2-methylpentane, and compounds A and B.

- (i) Complete the table opposite to show the mechanism for the formation of 3-bromo-2-methylpentane and THREE possible equations for termination.

In your equations, use **STRUCTURAL** or **SKELETAL FORMULAE** and 'dots' (•) for the position of radicals. [6]

INITIATION

Equation: _____

Conditions: _____

PROPAGATION

→

→

TERMINATION

→

→

→

- (ii) Organic compound A is formed by the substitution of ALL 14 H atoms in 2-methylpentane by Br atoms.

Write the equation, using MOLECULAR FORMULAE, for the formation of compound A from 2-methylpentane.

_____ [2]

- (iii) Organic compound B is formed by the substitution of SOME of the 14 H atoms in 2-methylpentane by Br atoms.

0.8649 g of compound B is heated until it is vaporised.

Under the conditions used:
compound B has a volume of 72.0 cm^3
the molar gas volume is $40.0 \text{ dm}^3 \text{ mol}^{-1}$.

Determine a possible molecular formula of compound B.

molecular formula = _____ [3]

17 This question is about alcohols.

- (a) An UNSATURATED alcohol has 6 carbon atoms and contains ONE C=C bond.


Construct an equation for the complete combustion of this alcohol.

_____ [2]

- (b) Compound C, shown below, is refluxed with excess acidified potassium dichromate(VI) to form a single organic product and one other product.


Complete the equation for this reaction. [3]

COMPOUND C




- (c) Compound D, shown below, is refluxed with H_2SO_4 , as an acid catalyst, to form a mixture of three isomers with the molecular formula C_7H_{10} .

COMPOUND D



- (i) Draw the structures of the **THREE** isomers of C_7H_{10} formed from compound D. Use the space below. [3]



- (ii) A student converts compound D into a diiodoalkane.

Suggest suitable reagents for this reaction.

_____ [1]

- (d) There are 4 structural isomers of $C_4H_{10}O$ that are alcohols.

A student predicts that these structural isomers could be distinguished using carbon-13 NMR spectroscopy.

Explain whether the student is correct.

In your answer, show how the peaks in the carbon-13 NMR spectra are linked to the structure of each alcohol isomer.

[5]

18 1,3-dinitrobenzene is a solid at room temperature.

A chemist prepares 1,3-dinitrobenzene as outlined below.

STEP 1 12.5 cm³ of nitrobenzene (density = 1.20 g cm⁻³) is refluxed with concentrated nitric acid in the presence of concentrated sulfuric acid as a catalyst.

STEP 2 The mixture is cooled. Impure crystals of 1,3-dinitrobenzene appear.

STEP 3 The impure crystals are purified to obtain pure 1,3-dinitrobenzene.

The chemist obtains 15.0 g of pure 1,3-dinitrobenzene.

- (a) Outline the mechanism for this reaction, including the role of H_2SO_4 as a catalyst. Use the space below. [5]

(b) Determine the percentage yield of 1,3-dinitrobenzene.

Give your answer to 3 significant figures.


percentage yield = _____ % [3]

(c) Describe how to purify the impure crystals in STEP 3.

[3]

BLANK PAGE

19* A student intends to synthesise compound Z, as shown in the flowchart below.



[illegible]


Additional answer space if required.

BLANK PAGE


20 This question is about aromatic compounds containing the –COOH and –OH functional groups.

(a) Salicylic acid, shown below, is used in the manufacture of some important medicines.


SALICYLIC ACID




Complete the flowchart opposite for reactions of salicylic acid, by adding the organic products in each box. [4]




Br_2




$\text{Na}_2\text{CO}_3(\text{aq})$




propanoic anhydride
 $(\text{CH}_3\text{CH}_2\text{CO})_2\text{O}$



+



- (b) **PAS**, shown below, is an antibiotic used to treat several diseases including tuberculosis (TB).



- (i) A student predicts that PAS could polymerise to form a polymer containing **BOTH** ester and amide linkages.

Draw a section of this polymer.

The section should contain ONE amide and ONE ester linkage, which should be displayed. Use the space below. [3]

- (ii) For the treatment of TB, the maximum daily dosage of PAS that should be prescribed is 300 mg per kg of body mass.

A child weighs 20.0 kg.

Calculate the number of PAS molecules in the maximum daily dosage of PAS for this child.

number of PAS molecules = _____ [3]

21 This question is about α -amino acids.

- (a) The general formula of an α -amino acid is $\text{RCH}(\text{NH}_2)\text{COOH}$.**

Most α -amino acids show optical isomerism.

Explain the term OPTICAL ISOMERISM.

[1]

- (b) The α -amino acid valine has the R group of $-\text{CH}(\text{CH}_3)_2$.**

(i) What is the systematic name of valine?


[1]

(ii) Draw diagrams to show 3D structures of the optical isomers of valine. [2]

--	--

- (c) Three α -amino acids can react together to form compound E, shown below.

COMPOUND E



- (i) How many optical isomers are possible for compound E?

_____ [1]

- (ii) A student hydrolyses compound E with dilute hydrochloric acid, HCl(aq) .

Draw the structures of the organic products formed by this hydrolysis. Use the space below. [4]


22 This question is about reactions of acrolein, $\text{H}_2\text{C}=\text{CHCHO}$.

(a) Acrolein reacts with sodium cyanide in acidic conditions, $\text{NaCN}(\text{aq})/\text{H}^+(\text{aq})$.

(i) Outline the reaction mechanism for this reaction, showing the intermediate and the organic product.

The structure of acrolein has been provided.

Include curly arrows and relevant dipoles. [4]



(ii) Name this type of mechanism.

_____ [1]

(b) Complete the flowchart by filling in each box. [9]

unsaturated alcohol	reagent(s)	diol (major product)
	catalyst	

↑ reagent(s)

ACROLEIN

C=CC=O

reagent(s)

carboxylic acid

reagent(s)

acyl chloride

polymer (2 repeat units)

BLANK PAGE

23* An unknown organic compound is analysed.

The results are shown below and opposite.

ADDITION OF 2,4-DNP

No visible change


ELEMENTAL ANALYSIS BY MASS

C, 66.63%; H, 11.18%; O, 22.19%

MASS SPECTRUM


Molecular ion peak at $m/z = 144.0$

IR SPECTRUM



PROTON NMR SPECTRUM

The numbers by each peak are the relative peak areas.



Show ALL your reasoning.

48

Additional answer space if required.

END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.