

OXFORD CAMBRIDGE AND RSA EXAMINATIONS**Advanced GCE****CHEMISTRY****2814**

Chains, Rings and Spectroscopy

Wednesday **19 JUNE 2002** Afternoon 1 hour 30 minutes

Candidates answer on the question paper.

Additional materials:

Data Sheet for Chemistry

Scientific calculator

Candidate Name	Centre Number	Candidate Number										
	<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>						<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>					

TIME 1 hour 30 minutes**INSTRUCTIONS TO CANDIDATES**

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer **all** the questions.
- Write your answers in the spaces on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the *Data Sheet for Chemistry*.
- You are advised to show all the steps in any calculations.

FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	10	
2	11	
3	10	
4	6	
5	14	
6	15	
7	12	
8	12	
TOTAL	90	

This question paper consists of 12 printed pages.

1 A ketone **A** has the molecular formula C_3H_6O .

(a) Name **A** and draw its structure to show clearly its functional group.

name

structure:

[2]

(b) Ketone **A** can be **reduced** to an alcohol, **B**.

(i) Name **B** and draw its structure.

name

structure:

[2]

(ii) State a suitable reagent for this reduction.

..... [1]

(iii) Write a balanced equation for the reduction of **A** to **B**. You may use the symbol [H] in this redox reaction to represent the reducing agent.

..... [1]

(c) Describe a chemical method to detect the presence of a carbonyl group in a compound such as **A**. Explain how you would use the product from this chemical method to identify **A**.

.....

.....

.....

.....

.....

..... [4]

[Total : 10]

- 2 The reaction of benzene with bromine requires a halogen carrier but the reaction of phenol with bromine does not.

(a) (i) Write the equation for the reaction of benzene with bromine.

..... [2]

(ii) State a substance that will act as the halogen carrier for this reaction.

..... [1]

(b) The reaction of phenol with excess bromine gives the organic product **C**.

(i) Draw the structure of **C**.

[2]

(ii) Cold aqueous NaOH is added to compound **C**. Using structural formulae, predict the equation for the reaction that takes place.

[2]

(iii) Explain why the reaction of phenol with bromine does **not** require a halogen carrier.


.....
.....
.....
.....
..... [3]

(iv) State a general use for halogenated phenols.

..... [1]

[Total : 11]

- 3 A commercial synthesis of the ester **G** is shown below.



- (a) Stage I:

(i) Suggest a suitable reagent.

..... [1]

(ii) State the type of reaction occurring.

..... [2]

(iii) Write the equation for this reaction.

..... [1]

- (b) Stage II:

(i) Draw the displayed formula for the ester **G**.

[1]

(ii) Write the equation.

..... [1]

(iii) Suggest a general use for esters such as **G**.


..... [1]

(iv) **G** can also be made directly from **E** by reaction with $\text{CH}_3\text{COO}^-\text{Na}^+$. Suggest a possible mechanism for this reaction.

[3]

[Total : 10]

- 5 (a) A section of a polymer has the structure shown below.



- (i) Circle a repeat unit of this polymer on the diagram above. [1]
- (ii) Deduce the empirical formula of this polymer.
..... [1]
- (iii) Draw a structure for a monomer from which this polymer could be made. Your structure should show any multiple bonds. [1]

- (b) Proteins are natural polymers made from α -amino acids, such as glycine, $\text{H}_2\text{NCH}_2\text{COOH}$.

- (i) Name the functional group made during amino acid polymerisation and draw its displayed formula.

name of functional group


displayed formula of functional group:

- (ii) Name this type of polymerisation reaction.


..... [1]

- (iii) Draw a displayed and a skeletal formula for the dipeptide **H**, $C_4H_8N_2O_3$, made from glycine, H_2NCH_2COOH .

displayed formula of **H**



skeletal formula of **H**



[2]

- (iv) A student made 1.10 g of dipeptide **H** starting from 1.40 g of glycine.
Calculate the percentage yield obtained. Give your answer to 3 significant figures.

Percentage yield % [4]



- (v) When glycine is treated with hydrochloric acid a compound **J**, $C_2H_6ClNO_2$, is formed. Draw a structure for compound **J**.

[2]

[Total : 14]

[Turn over

6 Compounds **K** and **L** are structural isomers.

**K****L**

(a) (i) What is the molecular formula of these isomers?

..... [1]

(ii) Calculate the mass:charge ratio, m/e , you expect for the molecular ion peak in the mass spectrum of **K**, showing your working.

Answer [1]

(iii) A sample of **L** is sent for analysis to determine its percentage by mass of carbon and hydrogen. Calculate the expected results.

%C

%H


[2]

(b) Explain how infra-red spectra would allow you to distinguish between samples of **K** and **L**.

.....

 [3]

- (c) (i) Compound **K** gives the n.m.r. spectrum below. Identify which of the protons are responsible for each peak. Explain your reasoning.



.....

 [3]

- (ii) A sample of **K** is shaken with D_2O and the spectrum is re-run. Describe how the spectrum is changed.

.....
 [1]

- (iii) Suggest possible δ values for the peaks in the n.m.r. spectrum of compound **L**. For each peak, give the number of protons responsible.

.....

 [4]

[Total : 15]

OCR has made every effort to trace the copyright holders of items used in this Question paper, but if we have inadvertently overlooked any, we apologise.