



CHAINS + RINGS
Mark Scheme 2812
June 2003

The following annotations may be used when marking:

X	=	incorrect response (errors may also be underlined)
^	=	omission mark
bod	=	benefit of the doubt (where professional judgement has been used)
ecf	=	error carried forward (in consequential marking)
con	=	contradiction (in cases where candidates contradict themselves in the same response)
sf	=	error in the number of significant figures

Abbreviations, annotations and conventions used in the Mark Scheme:

/	=	alternative and acceptable answers for the same marking point
;	=	separates marking points
NOT	=	answers not worthy of credit
()	=	words which are not essential to gain credit
<u> </u> (underlining)	=	key words which <u>must</u> be used
ecf	=	allow error carried forward in consequential marking
AW	=	alternative wording
ora	=	or reverse argument

2812

Mark Scheme

June 2003



1.

(a)(i) + each by its own A_r to give 5 : 13.3 : 1.67 ✓ [1]

+ each by 1.67 to give 3 : 8 : 1 ✓ [1]

(ii) Evidence of working e.g. $36 + 8 + 16 = 60$ / that C_3H_8O adds up to 60 ✓ [1]

(b) unambiguous structure/formula of propan-1-ol & propan-2-ol to include:



✓ [2]

(c)(i) dichromate/ $\text{Cr}_2\text{O}_7^{2-}$ / MnO_4^- ✓ [1](ii) orange to green ✓✓ [2]
purple to green/brown/black/pink/colourless(iii) continuous boiling/evaporation and condensation /
heating & return of liquid to reaction flask/
simple sketch showing vertical condenser & heat ✓ [1]
(any reference to a closed system negates the mark)

(d)(i) OH/alcohol/hydroxy/hydroxyl – not hydroxide ✓ [1]

(ii) C=O/carbonyl – not CO ✓ [1]


(iii) carboxylic acid/- CO_2H /- COOH ✓ [1](e) propan-1-ol (no marks)
propan-1-ol oxidised to a carboxylic acid/ ✓ [1](f) $\text{C}_3\text{H}_8\text{O} + 2[\text{O}] \rightarrow \text{CH}_3\text{CH}_2\text{COOH} / \text{C}_3\text{H}_6\text{O}_2 + \text{H}_2\text{O}$ ✓✓ [2]
1 mark available if, $\text{CH}_3\text{CH}_2\text{COOH}$ & H_2O present in the equation

[Total : 15]

2.
 (a)(i) 1,1-dibromoethene ✓ [1]
 (ii) CHBr ✓ [1]
 (b)(i) (Br₂ is) decolourised ✓ [1]
 (ii) electrophilic addition ✓ [1]
 ✓ [1]

(c) allow names & unambiguous formulae throughout part (c)

(i) Isomer C reacts with H₂.




✓ [1]

conditions

suitable catalyst such as Ni/Pt/Pd


✓ [1]

(ii)



✓✓ [2]

(iii)



✓ [1]

conditions

phosphoric acid (catalyst)
temp ≥ 100 °C/ steam

✓ [1]
✓ [1]

[Total : 12]

2812

Mark Scheme

June 2003

- 3.
- (a) non-polar ✓ [1]
- hence particles not attracted to methane ✓ [1]
- (b)
- (free radical) substitution ✓ [1]
 - $\text{CH}_4 + \text{Br}_2 \rightarrow \text{CH}_3\text{Br} + \text{HBr}$ ✓ [1]
 - ultra violet/UV light ✓ [1]
 - $\text{Br}_2 \rightarrow 2 \text{Br}\cdot$ ✓ [1]
 - homolysis/ homolytic fission ✓ [1]
 - $\text{Br}\cdot + \text{CH}_4 \rightarrow \cdot\text{CH}_3 + \text{HBr}$ ✓ [1]
 - $\cdot\text{CH}_3 + \text{Br}_2 \rightarrow \text{CH}_3\text{Br} + \text{Br}\cdot$ ✓ [1]
 - any two free radicals $2 \text{Br}\cdot \rightarrow \text{Br}_2$ ✓ [1]
- free rads are difficult to control/react with anything/very reactive ✓ [1]
- identifies one of CH_2Br_2 / CHBr_3 / CBr_4 or can be polysubstituted ✓ [1]
- [10 max = 9]

1 QWC mark is available for using specific chemical terms.

chemical terms: initiation, propagation, termination, free radical substitution,
homolysis/ homolytic fission, photochemical


any two terms used correctly ✓ [1]

[Total : 11]

4.



(ii)



marking points:

dipoles

curly arrow from OH^- to $C^{\delta+}$

curly arrow from C-Cl bond to Cl

✓✓✓ [3]






[Total : 6]

2812

Mark Scheme

June 2003

5.



[Total : 15]