

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS****Advanced GCE****CHEMISTRY****2816/01**

Unifying Concepts in Chemistry

Wednesday

**21 JANUARY 2004**

Morning

1 hour 15 minutes

Candidates answer on the question paper.

Additional materials:

*Data Sheet for Chemistry*

Scientific calculator

Candidate Name

Centre Number

Candidate  
Number

--	--	--	--	--	--	--	--

**TIME** 1 hour 15 minutes**INSTRUCTIONS TO CANDIDATES**

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer **all** the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the *Data Sheet for Chemistry*.
- You are advised to show all the steps in any calculations.


FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	16	
2	17	
3	13	
4	14	
<b>TOTAL</b>	<b>60</b>	

---

**This question paper consists of 11 printed pages and 1 blank page.**

Answer all the questions.

- 1 The formation of ethyl ethanoate and water from ethanoic acid and ethanol is a reversible reaction which can be allowed to reach equilibrium. The equilibrium is shown below.



- (a) Write the expression for  $K_c$  for this equilibrium system.

[2]

- (b) A student mixed together 6.0 mol ethanoic acid and 12.5 mol ethanol. A small amount of hydrochloric acid was also added to catalyse the reaction. He left the mixture for two days to reach equilibrium in a water bath at constant temperature, after which time 1.0 mol ethanoic acid remained.

- (i) Complete the table below to show the equilibrium composition of the equilibrium mixture.

component	$\text{CH}_3\text{COOH}$	$\text{C}_2\text{H}_5\text{OH}$	$\text{CH}_3\text{COOC}_2\text{H}_5$	$\text{H}_2\text{O}$
initial amount / mol	6.0	12.5	0.0	0.0
equilibrium amount / mol				

[2]

- (ii) Calculate  $K_c$  to two significant figures. State the units, if any. The total volume of the equilibrium mixture is  $1.0 \text{ dm}^3$ .

$K_c = \dots\dots\dots$  units  $\dots\dots\dots$  [2]

- (c) The student was concerned that the mixture may **not** have reached equilibrium. What could he do to be sure that equilibrium had been reached?

.....  
.....  
.....[2]

- (d) The student added more ethanol to the mixture.

- (i) State, giving a reason, what would happen to the composition of the equilibrium mixture.

.....  
.....  
.....[2]

- (ii) What happens to the value of  $K_c$ ?

.....[1]

- (e) The student added more of the acid catalyst to the mixture. State, giving a reason, what would happen to the composition of the equilibrium mixture.

.....  
.....  
.....[2]

- (f) The student repeated the experiment at a higher temperature and found that the value of  $K_c$  decreased.

- (i) State, giving a reason, what would happen to the composition of the equilibrium mixture.

.....  
.....  
.....[2]

- (ii) What additional information does this information tell you about the reaction?

.....  
.....[1]

[Total: 16]

- 2 Hydrogen peroxide,  $\text{H}_2\text{O}_2$ , is a colourless liquid, widely used as an oxidising agent, antiseptic, and bleach for hair and cloth.

Hydrogen peroxide reacts with iodide ions,  $\text{I}^-$ , in the presence of acid,  $\text{H}^+(\text{aq})$ , forming iodine,  $\text{I}_2$ .

- (a) Suggest a balanced equation for the overall reaction between  $\text{H}_2\text{O}_2(\text{aq})$ ,  $\text{I}^-(\text{aq})$  and  $\text{H}^+(\text{aq})$  to form aqueous iodine.

.....[2]

- (b) Three experiments were carried out using different initial concentrations of  $\text{H}_2\text{O}_2(\text{aq})$ ,  $\text{I}^-(\text{aq})$  and  $\text{H}^+(\text{aq})$ . The initial rate of formation of  $\text{I}_2$  was measured for each experiment.

The experimental results are shown below.

experiment	$[\text{H}_2\text{O}_2(\text{aq})]$ /mol dm <sup>-3</sup>	$[\text{I}^-(\text{aq})]$ /mol dm <sup>-3</sup>	$[\text{H}^+(\text{aq})]$ /mol dm <sup>-3</sup>	rate /mol dm <sup>-3</sup> s <sup>-1</sup>
1	0.010	0.010	0.005	$1.15 \times 10^{-6}$
2	0.010	0.020	0.005	$4.60 \times 10^{-6}$
3	0.010	0.020	0.010	$4.60 \times 10^{-6}$

- (i) Showing all your reasoning, determine the reaction orders for  $\text{I}^-$  and for  $\text{H}^+$ .

.....  
 .....  
 .....  
 .....  
 .....  
 .....  
 .....[4]

- (ii) This reaction is first order with respect to  $\text{H}_2\text{O}_2$ .  
 Use this information and your answers to (i) to complete the rate equation for this reaction.


rate = .....[2]

- (iii) Calculate the rate constant  $k$  for this reaction. State the units for  $k$ .

rate constant,  $k$  ..... units ..... [3]

- (c) This reaction was shown to be first order with respect to  $\text{H}_2\text{O}_2$  by plotting a rate-concentration graph.

Using the axes below, sketch a graph to show how the rate of this reaction changes with increasing  $\text{H}_2\text{O}_2$  concentration.



[2]

- (d) Hydrogen peroxide readily decomposes to give water and oxygen.

Hydrogen peroxide is sold by volume strength. For example, 20-volume  $\text{H}_2\text{O}_2$  yields 20 volumes of oxygen gas for each volume of aqueous  $\text{H}_2\text{O}_2$  solution.

- (i) Construct an equation for the decomposition of hydrogen peroxide.

.....[1]

- (ii) Determine the concentration, in  $\text{mol dm}^{-3}$ , of 20-volume hydrogen peroxide.


Show all your working clearly.

answer .....  $\text{mol dm}^{-3}$  [3]

[Total: 17]

- 3 Carbonic acid,  $\text{H}_2\text{CO}_3$ , is a weak Bronsted-Lowry acid formed when carbon dioxide dissolves in water. Blood contains several buffer solutions and healthy blood is buffered to a pH of 7.40. The most important buffer solution in blood is a mixture of carbonic acid and hydrogencarbonate ions,  $\text{HCO}_3^-$ .

The equilibrium in the carbonic acid / hydrogencarbonate buffer system is shown below.



- (a) Carbonic acid is a weak Bronsted-Lowry acid.

What is meant by the following terms?

- (i) A Bronsted-Lowry acid.

.....  
..... [1]

- (ii) A weak acid.

.....  
..... [1]

- (iii) pH.


..... [1]

- (iv) A buffer solution.

.....  
..... [1]



- 4 Maleic anhydride (*cis*-butenedioic anhydride) is an important industrial chemical. The structure of maleic anhydride is shown below.



maleic anhydride  
 $C_4H_2O_3$


- (a) In industry, maleic anhydride is produced on a large scale by oxidation of butane in air over a hot catalyst.

- (i) Suggest the industrial source of butane.

.....[1]

- (ii) An incomplete equation for this reaction is given below.

Complete the equation.



[2]


- (iii) Calculate the mass, in kg, of maleic anhydride that could be made by completely converting  $30 \text{ m}^3$  of butane in this reaction. ( $1 \text{ m}^3 = 1000 \text{ dm}^3$ )

Assume that the molar volume of butane under the conditions used is  $24 \text{ dm}^3$ .

answer ..... kg [3]



- (b) Maleic anhydride can be converted into tartaric acid by reaction with water and a suitable oxidising agent.




Deduce the **empirical** formula of tartaric acid.

[2]

- (c) 'Cream of tartar' is often used in cookery. This compound can be prepared by reacting aqueous solutions of tartaric acid and potassium hydroxide in 1:1 molar proportions.

- (i) Complete the equation below for the preparation of 'cream of tartar'.



[2]

- (ii) Suggest another chemical that would react with aqueous tartaric acid. The chemical you choose should **not** be a hydroxide or an oxide.

State what you would expect to see and write an equation for your chosen reaction.

chemical .....

observation(s) .....

.....

equation

[4]

[Total: 14]

**END OF QUESTION PAPER**

**BLANK PAGE**

