

**Mark Scheme 2816/01**  
**June 2005**

UNIFYING CONCEPTS  
IN CHEMISTRY



| Abbreviations, annotations and conventions used in the Mark Scheme | / = alternative and acceptable answers for the same marking point<br>; = separates marking points<br>NOT = answers which are not worthy of credit<br>( ) = words which are not essential to gain credit<br><u>      </u> = (underlining) key words which <b>must</b> be used to gain credit<br>ecf = error carried forward<br>AW = alternative wording<br>ora = or reverse argument  |                                 |
|--|--|---------------------------------|
| Question   | Expected Answers   | Marks                           |
| 1 (a)  | (i) constant half-life ✓<br>(ii) rate = $k [N_2O_5]$ ✓<br>Common error will be to use '2' from equation.<br>(iii) curve downwards getting less steep ✓<br>curve goes through 1200,0.30; 2400,0.15; 3600,0.075 ✓<br>(iv) tangent shown on graph at $t = 1200\text{ s}$ ✓<br>(v) $3.7(2) \times 10^{-4}$ ✓ $\text{mol dm}^{-3} \text{s}^{-1}$ ✓<br>ecf possible from (ii) using $[N_2O_5]^x$<br>(2nd order answer: $2.2(3) \times 10^{-4}$ ) | [1]<br>[1]<br>[2]<br>[1]<br>[2] |
| (b)  | (i) slow step ✓<br>(ii) $(CH_3)_2C=CH_2 + H_2O \longrightarrow (CH_3)_3COH$ ✓<br>(iii) $H^+$ is a catalyst ✓<br>$H^+$ used in first step and formed in second step/<br>regenerated/ not used up ✓<br>(iv) rate = $k [(CH_3)_2C=CH_2] [H^+]$ ✓<br>common error will be use of $H_2O$ instead of $H^+$   | [1]<br>[1]<br>[2]<br>[1]        |
|  |  | <b>Total: 12</b>                |

| Abbreviations, annotations and conventions used in the Mark Scheme | / = alternative and acceptable answers for the same marking point<br>; = separates marking points<br>NOT = answers which are not worthy of credit<br>( ) = words which are not essential to gain credit<br><u>     </u> = (underlining) key words which <u>must</u> be used to gain credit<br>ecf = error carried forward<br>AW = alternative wording<br>ora = or reverse argument  |                              |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
|--|---|------------------------------|----------------|--------------------|-----|-----|-----|-----|-------|-------|------------------------------|--|------------------------------|--|--------------------------|---|------------|------------|--------------|--------------------------|
| Question   | Expected Answers  | Marks                        |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 2 (a)  | <p><b>High Pressure</b><br/>Equilibrium <math>\longrightarrow</math> right as fewer moles on right hand side and the shift reduces number of molecules/compensates for increasing pressure ✓<br/>Rate increases/ more collisions ✓</p> <p><b>High temperature</b><br/>Equilibrium <math>\longrightarrow</math> left as equilibrium goes to the left to compensate for increased temperature/absorbs the energy/in endothermic direction (ora) ✓<br/>Rate increases/ more successful collisions ✓</p> <p><b>Other effect</b><br/>High pressures expensive/ high temperatures expensive /high pressures cause safety problems ✓</p> <p>One correct statement followed by correct explanation ✓</p>  | [2]<br>[2]<br>[1]<br>[1]     |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 2 (b) (i)  | <table border="1"> <tr> <td>CO</td> <td>H<sub>2</sub></td> <td>CH<sub>3</sub>OH</td> </tr> <tr> <td>1.0</td> <td>2.0</td> <td>0.0</td> </tr> <tr> <td>0.9</td> <td>1.8 ✓</td> <td>0.1 ✓</td> </tr> <tr> <td>0.9/2.8 or 0.321 or 0.32/0.3</td> <td></td> <td>1.8/2.8 or 0.643 or 0.64/0.6</td> </tr> <tr> <td></td> <td>0.1/2.8 or 0.036 or 0.04</td> <td>✓</td> </tr> <tr> <td>3.21 (MPa)</td> <td>6.43 (MPa)</td> <td>0.36 (MPa) ✓</td> </tr> </table> <p>In 3rd and 4th rows, ecf from previous row</p> <p>(ii) <math>K_p = \frac{p(\text{CH}_3\text{OH})}{p(\text{CO}) \times p(\text{H}_2)^2}</math> ✓✓</p> <p>1 mark for <math>K_c</math>/ use of any [ ] /inverted/power missing.</p> <p>(iii) <math>K_p</math> stays the same ✓<br/>Equilibrium position moves to the right/yield increases ✓ in response to increase in reactants ✓</p> <p>(iv) <math>K_p = \frac{0.261}{3.70 \times 5.10^2} = 2.71 \times 10^{-3}</math> ✓ MPa<sup>-2</sup> ✓</p> <p>calc value <math>2.7120546 \times 10^{-3}</math>; answer and/or units ecf from (ii)</p> | CO                           | H <sub>2</sub> | CH <sub>3</sub> OH | 1.0 | 2.0 | 0.0 | 0.9 | 1.8 ✓ | 0.1 ✓ | 0.9/2.8 or 0.321 or 0.32/0.3 |  | 1.8/2.8 or 0.643 or 0.64/0.6 |  | 0.1/2.8 or 0.036 or 0.04 | ✓ | 3.21 (MPa) | 6.43 (MPa) | 0.36 (MPa) ✓ | [4]<br>[2]<br>[3]<br>[2] |
| CO   | H <sub>2</sub>  | CH <sub>3</sub> OH           |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 1.0  | 2.0   | 0.0                          |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 0.9  | 1.8 ✓   | 0.1 ✓                        |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 0.9/2.8 or 0.321 or 0.32/0.3                                       |   | 1.8/2.8 or 0.643 or 0.64/0.6 |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
|  | 0.1/2.8 or 0.036 or 0.04  | ✓                            |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| 3.21 (MPa)   | 6.43 (MPa)  | 0.36 (MPa) ✓                 |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
| (c)  | $\text{CH}_3\text{OH} + 1.5\text{O}_2 \longrightarrow \text{CO}_2 + 2\text{H}_2\text{O}$ ✓  | [1]                          |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |
|  |   | Total: 18                    |                |                    |     |     |     |     |       |       |                              |  |                              |  |                          |   |            |            |              |                          |

| Abbreviations, annotations and conventions used in the Mark Scheme |  | / = alternative and acceptable answers for the same marking point<br>: = separates marking points<br>NOT = answers which are not worthy of credit<br>( ) = words which are not essential to gain credit<br>_____ = (underlining) key words which <b>must</b> be used to gain credit<br>ecf = error carried forward<br>AW = alternative wording<br>ora = or reverse argument |
|--|--|---|
| Question   | Expected Answers   | Marks   |
| 3 (a) (i)  | completely dissociates/ionised ✓<br>proton donor ✓   | [2]   |
| (ii)   | NO <sub>3</sub> <sup>-</sup> ✓   | [1]   |
| (b) (i)  | pH = -log[H <sup>+</sup> ] / -log(0.015) ✓ = 1.82 / 1.8 ✓ (Not 2)  | [2]   |
| (ii)   | [H <sup>+</sup> ] = 0.0075 mol dm <sup>-3</sup><br>pH = -log(0.0075) = 2.12 / 2.1 ✓  | [1]   |
| (c) (i)  | K <sub>w</sub> = [H <sup>+</sup> (aq)] [OH <sup>-</sup> (aq)] ✓ <i>state symbols not needed</i>  | [1]   |
| (ii)   | [H <sup>+</sup> (aq)] = 10 <sup>-pH</sup> = 10 <sup>-13.54</sup> = 2.88/2.9 × 10 <sup>-14</sup> mol dm <sup>-3</sup> ✓<br>$[\text{NaOH}] / [\text{OH}^-(\text{aq})] = \frac{K_w}{[\text{H}^+(\text{aq})]} = \frac{1.0 \times 10^{-14}}{2.88 \times 10^{-14}}$ = 0.347 / 0.35 mol dm <sup>-3</sup> ✓  | [2]   |
| (d) (i)  | a solution that <del>minimises</del> /resists/opposes pH changes ✓   | [1]   |
| (ii)   | The buffer must contain both CH <sub>3</sub> COOH and CH <sub>3</sub> COONa / CH <sub>3</sub> COO <sup>-</sup> / weak acid and conjugate base ✓<br><br>Solution A is a mixture of CH <sub>3</sub> COOH and CH <sub>3</sub> COONa / / has an excess of acid / is acidic ✓<br><br>Solution B, contains only CH <sub>3</sub> COONa/ only CH <sub>3</sub> COO <sup>-</sup> / only the salt/ is neutral ✓<br><br>CH <sub>3</sub> COOH(aq) + NaOH(aq) → CH <sub>3</sub> COONa(aq) + H <sub>2</sub> O(l) / acid/alkali has been neutralised/ CH <sub>3</sub> COOH(aq) and NaOH react together ✓ | [4]   |
| (e)  | [H <sup>+</sup> ] increases ✓<br>H <sub>2</sub> O ionises more /<br>for H <sub>2</sub> O ⇌ H <sup>+</sup> + OH <sup>-</sup> , equilibrium moves to the right ✓<br><br>exo/endo is 'noise'  | [2]   |
|  |  | <b>Total: 15</b>  |

| Abbreviations, annotations and conventions used in the Mark Scheme | / = alternative and acceptable answers for the same marking point<br>; = separates marking points<br>NOT = answers which are not worthy of credit<br>( ) = words which are not essential to gain credit<br>— = (underlining) key words which <b>must</b> be used to gain credit<br>ecf = error carried forward<br>AW = alternative wording<br>ora = or reverse argument  |   |
|--|--|---|
| Question   | Expected Answers   | Marks   |
| 4 (a)  | moles of Cu = $0.68 \times 5/1000 = 0.0034$ ✓<br>mass of Cu = $0.0034 \times 63.5 = 0.216$ g ✓<br>% Cu = $0.216/0.28 = 77\%$ ✓<br><br>ratios:<br>Cu = $26.29/63.5 = 0.41$<br>N = $11.6/14 = 0.83$<br>O = $59.63/16 = 3.73$<br>H = $2.48/1 = 2.48$<br>OR<br>1<br>2<br>9<br>6 ✓<br><br>empirical formula = $\text{CuN}_2\text{O}_9\text{H}_6$ ✓<br><br>Formula with $3\text{H}_2\text{O}$ shown separately scores 1:<br>i.e. $\text{CuN}_2\text{O}_6 \cdot 3\text{H}_2\text{O}$ ✓<br>Correct formula shown with $(\text{NO}_3)_2$ scores 2nd mark:<br>$\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ ✓<br>(Correct answer automatically scores both marks) | [3]<br><br><br><br><br><br><br><br><br><br>[2]<br><br><br><br><br><br><br><br><br><br>[2] |
| (b)  | $\text{Cu} \rightarrow \text{Cu}^{2+}$ : Cu from 0 to +2 ✓<br>$\text{NO}_3^- \rightarrow \text{NO}$ : N from +5 to +2 ✓<br><br>$3\text{Cu} + 8\text{H}^+ + 2\text{NO}_3^- \rightarrow 3\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$ ✓<br><br>'simple balance' as the only creditworthy response scores 1 mark:<br>i.e. $\text{Cu} + 4\text{H}^+ + \text{NO}_3^- \rightarrow \text{Cu}^{2+} + \text{NO} + 2\text{H}_2\text{O}$  | [3]   |
| (c)  | moles of A = $90/24000 = 3.75 \times 10^{-3}$ ✓<br>$M_r$ of A = $0.24 / 3.75 \times 10^{-3} = 64$ ✓<br><br>Gas is $\text{SO}_2$ ✓<br><br>$\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$ /<br>$\text{Cu} + 4\text{H}^+ + \text{SO}_4^{2-} \rightarrow \text{Cu}^{2+} + \text{SO}_2 + 2\text{H}_2\text{O}$ /<br>$\text{Cu} + 3\text{H}^+ + \text{HSO}_4^- \rightarrow \text{Cu}^{2+} + \text{SO}_2 + 2\text{H}_2\text{O}$ ✓  | [4]   |
|  |  | Total: 14   |