



**General Certificate of Education**

**Mathematics 6360**

**MPC3      Pure Core 3**

**Mark Scheme**

*2010 examination - January series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2010 AQA and its licensors. All rights reserved.

#### COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

**Key to mark scheme and abbreviations used in marking**

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

**No Method Shown**

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.


Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**


## MPC3

Q	Solution	Marks	Total	Comments
1(a)	$y' = e^{-4x}(2x+2) - 4e^{-4x}(x^2+2x-2)$	M1	3	$y' = Ae^{-4x}(ax+b) \pm Be^{-4x}(x^2+2x-2)$ where $A$ and $B$ are non-zero constants All correct  or $-4x^2e^{-4x} - 6xe^{-4x} + 10e^{-4x}$
	$= e^{-4x}(2x+2-4x^2-8x+8)$	A1		
	$= 2e^{-4x}(5-3x-2x^2)$	A1		
	or $y = x^2 e^{-4x} + 2xe^{-4x} - 2e^{-4x}$			
	$y' = -4x^2e^{-4x} + 2xe^{-4x} + 2x - 4e^{-4x}$ $+ 2e^{-4x} + 8e^{-4x}$ $= -4x^2e^{-4x} - 6xe^{-4x} + 10e^{-4x}$ $= 2e^{-4x}(5-3x-2x^2)$	(M1) (A1)  (A1)		
(b)	$-(2x+5)(x-1) (=0)$	M1	5	OE Attempt at factorisation $(\pm 2x \pm 5)(\pm x \pm 1)$ or formula with at most one error  Both correct and no errors  SC $x = 1$ only scores M1A0  For $y = ae^b$ attempted  Either correct, follow through only from incorrect sign for $x$  CSO 2 solutions only  Note: withhold final mark for extra solutions Note: approximate values only for $y$ can score m1 only
	$x = \frac{-5}{2}, 1$	A1		
	$x=1, y=e^{-4}$	m1		
		A1F		
	$x = -\frac{5}{2}, y = e^{10} \left( -\frac{3}{4} \right)$	A1		
	<b>Total</b>		<b>8</b>	

## MPC3 (cont)

Q	Solution	Marks	Total	Comments
2(a)(i)		B1		correct shape passing through origin and stopping at $A$ and $B$
	$A\left(1, \frac{\pi}{2}\right)$ $B\left(-1, -\frac{\pi}{2}\right)$	B1 B1	3	SC $A(1, 90)$ and $B(-1, -90)$ scores B1
(ii)	line intersecting their curve (positive gradient, positive $y$ intercept) Correct statement	M1 A1	2	one solution only, stated or indicated on sketch - must be in the first quadrant (ie curve intersects line once) Must have scored B1 for graph in (a)(i)
(b)	$\left. \begin{array}{l} \text{LHS}(0.5) = 0.5 \quad \text{RHS}(0.5) = 1.1 \\ \text{LHS}(1) = 1.6 \quad \text{RHS}(1) = 1.3 \end{array} \right\}$ At 0.5 LHS < RHS, At 1 LHS > RHS $\therefore 0.5 < \alpha < 1$ <b>or</b> $f(x) = \sin^{-1}(x) - \frac{1}{4}x - 1$ $\left. \begin{array}{l} f(0.5) = -0.6 \\ f(1) = 0.3 \end{array} \right\}$ AWRT Change of sign $\Rightarrow 0.5 < \alpha < 1$ <b>or</b> $f(x) = \sin\left(\frac{1}{4}x + 1\right) - x$ $\left. \begin{array}{l} f(0.5) = 0.4 \\ f(1) = -0.1 \end{array} \right\}$ Attempt Change of sign $\Rightarrow 0.5 < \alpha < 1$ <b>or</b> $f(x) = 4\sin^{-1}x - x - 4$ $\left. \begin{array}{l} f(0.5) = -2.4 \\ f(1) = 1.3 \end{array} \right\}$ attempt Change of sign $\Rightarrow 0.5 < \alpha < 1$	M1 A1  (M1) (A1)  (M1) (A1)  (M1) (A1)	2	CSO  $f(x)$ must be defined  Allow $f(0.5) < 0$ $f(1) > 0$  $f(x)$ must be defined  $f(x)$ must be defined

## MPC3 (cont)

Q	Solution	Marks	Total	Comments
2(c)(i)	$x_2=0.902$ $x_3=0.941$	M1 A1	2	Sight of AWRT 0.902 or AWRT 0.941 These values only
(ii)		M1  A1	2	Staircase, (vertical line) from $x_1$ to curve, horizontal to line, vertical to curve  $x_2, x_3$ approx correct position on $x$ -axis
<b>Total</b>			<b>11</b>	

## MPC3 (cont)

Q	Solution	Marks	Total	Comments
3(a)	$\sin x = \frac{1}{3}$ , or sight of $\pm 0.34, \pm 0.11\pi$ or $\pm 19.47$ (or better)	M1	2	Penalise if incorrect answers in range; ignore answers outside range
	$x = 0.34, 2.8(0)$ AWRT	A1		
(b)	$\operatorname{cosec}^2 x - 1 = 11 - \operatorname{cosec} x$ $\operatorname{cosec}^2 x + \operatorname{cosec} x - 12 (=0)$ $(\operatorname{cosec} x + 4)(\operatorname{cosec} x - 3) (=0)$ $\operatorname{cosec} x = -4, 3$	M1 A1 m1	6	Correct use of $\cot^2 x = \operatorname{cosec}^2 x - 1$  Attempt at Factors Gives $\operatorname{cosec} x$ or $-12$ when expanded Formula one error condoned  Either Line
	$\left. \begin{array}{l} \sin x = -\frac{1}{4}, \frac{1}{3} \\ \sin x = -\frac{1}{4} \end{array} \right\}$ $\Rightarrow x = 3.39, 6.03$ AWRT $0.34, 2.8(0)$ AWRT	A1 B1F B1		
	<b>Alternative</b> $\frac{\cos^2 x}{\sin^2 x} = 11 - \frac{1}{\sin x}$ $\cos^2 x = 11 \sin^2 x - \sin x$ $1 - \sin^2 x = 11 \sin^2 x - \sin x$ $0 = 12 \sin^2 x - \sin x - 1$ $0 = (4 \sin x + 1)(3 \sin x - 1)$ $\sin x = -\frac{1}{4}, \frac{1}{3}$	(M1)  (A1) (m1) (A1) (B1F) (B1)		3 correct or their two answers from (a) and 3.39, 6.03 4 correct and no extras in range ignore answers outside range SC 19.47, 160.53, 194.48, 345.52    B1  Correct use of trig ratios and multiplying by $\sin^2 x$  Attempt at factors as above  As above
	<b>Total</b>		<b>8</b>	

## MPC3 (cont)

Q	Solution	Marks	Total	Comments										
4(a)		M1		Modulus graph V shape in 1 <sup>st</sup> quad going into 2 <sup>nd</sup> quad, touching $x$ -axis. Must cross $y$ -axis Condone not ruled 4 and 8 labelled										
(b)	$x = 2$ $x = 6$	B1 B1	2											
(c)	$x > 6$ $x < 2$	B1 B1	2											
<b>Total</b>			<b>6</b>											
5(a)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th><math>x</math></th> <th><math>y</math></th> </tr> </thead> <tbody> <tr> <td>1.5</td> <td>1.98100</td> </tr> <tr> <td>4.5</td> <td>3.22883</td> </tr> <tr> <td>7.5</td> <td>4.11496</td> </tr> <tr> <td>10.5</td> <td>4.74710</td> </tr> </tbody> </table> $\int = 3 \times \sum y$ $= 42.2$	$x$	$y$	1.5	1.98100	4.5	3.22883	7.5	4.11496	10.5	4.74710	B1 M1 A1		$x$ values correct PI 3+ $y$ values correct to 2sf or better or exact values 1.981, 3.228/9, 4.114/5, 4.747 for $y$ (or better)
$x$	$y$													
1.5	1.98100													
4.5	3.22883													
7.5	4.11496													
10.5	4.74710													
(b)(i)	$y = \ln(x^2 + 5)$ $e^y = x^2 + 5$ $x^2 = e^y - 5$	B1	1	OE AG Must see middle line, and no errors										
(ii)	$(\pi) \int (e^y - 5) (dy)$ $= (\pi) [e^y - 5y]_{(5)}^{(10)}$ $= (\pi) [(e^{10} - 50) - (e^5 - 25)]$ $V = \pi [e^{10} - e^5 - 25]$	M1 A1 m1 A1	4	Condone omission of brackets around $f(y)$ throughout F(10) – F(5) CSO including correct notation – must see $dy$ ISW if evaluated										
(c)	$(y =) \ln \left[ \left( \frac{x}{4} \right)^2 + 5 \right] + 3$	M1 B1 A1	3	$\frac{x}{4}$ seen, condone $\ln \frac{x^2}{4} + \dots$ ... + 3 CSO mark final answer (no ISW)										
<b>Total</b>			<b>12</b>											



## MPC3 (cont)

Q	Solution	Marks	Total	Comments	
6(a)	$f(x) > -3$	M1		' $> -3$ ', ' $x > -3$ ' or ' $f(x) \geq -3$ '	
(b)(i)	$y = e^{2x} - 3$ $y + 3 = e^{2x}$ $\ln(y + 3) = 2x$	A1	2	Allow $y > -3$	
	$(f^{-1}(x)) = \frac{1}{2} \ln(x + 3)$	M1		swap $x$ and $y$	
	<b>Alternative</b> $x \rightarrow \times 2 \rightarrow e \rightarrow -3$ $\div 2 \leftarrow \ln \leftarrow + 3 \leftarrow x$ (M1) (M1)	M1		attempt to isolate: $\ln(y \pm A) = Bx$ or reverse	
	$y = \frac{\ln(x + 3)}{2}$	A1	3	OE with no further incorrect working Condone $y = \dots$	
(ii)	$x + 3 = 1$	(A1)			
	$x = -2$	M1		for putting their $p(x) = 1$ from $k \ln(p(x))$ in their part (b)(i)	
(c)(i)	$(gf(x)) = \frac{1}{3(e^{2x} - 3) + 4}$ $(=) \frac{1}{3e^{2x} - 5}$	either OE	B1	1	substituting $f$ into $g$ ISW
(ii)	$\frac{1}{3e^{2x} - 5} = 1$ $1 = 3e^{2x} - 5$ $e^{2x} = 2$ $2x = \ln 2$ $x = \frac{1}{2} \ln 2$	OE	M1		Correct removal of their fraction
		m1			Correct use of logs leading to $kx = \ln \frac{a}{b}$
		OE	A1	3	CSO No ISW except for numerical evaluation
	<b>Total</b>		<b>11</b>		

## MPC3 (cont)

Q	Solution	Marks	Total	Comments	
7(a)	$\left(\frac{dy}{dx}\right) = \frac{\cos 4x \cdot 4 \cos 4x - \sin 4x \cdot -4 \sin 4x}{\cos^2 4x}$	M1	3	$\frac{\pm A \cos^2 4x \pm B \sin^2 4x}{\cos^2 4x}$	
	$= \frac{4 \cos^2 4x + 4 \sin^2 4x}{\cos^2 4x}$ or better	A1		Both terms correct	
	$= 4(1 + \tan^2 4x)$ CSO	A1		All correct	
	<b>or</b>				
	$\left(\frac{dy}{dx}\right) = \frac{\cos 4x \cdot 4 \cos 4x - \sin 4x \cdot -4 \sin 4x}{\cos^2 4x}$	(M1)		$\frac{\pm A \cos^2 4x \pm B \sin^2 4x}{\cos^2 4x}$	
	$= \frac{4 \cos 4x \cos 4x}{\cos 4x \cos 4x} + \frac{4 \sin 4x \sin 4x}{\cos 4x \cos 4x}$	(A1)			
	or better				
	$= 4(1 + \tan^2 4x)$ CSO	(A1)		All correct	
	(b)	$\frac{d^2 y}{dx^2} = 4 \times 2 \tan 4x \times \dots$	M1	5	$A \tan 4x \times f(4x)$
		$4 \sec^2 4x$	m1		$f(4x) = B \sec^2 4x$
$= 32 \tan 4x \sec^2 4x$		A1F	ft $8 \times$ their $p$ from part (a)		
$= 32 \tan 4x (1 + \tan^2 4x)$		m1	Previous two method marks must have been earned		
$= 32y(1 + y^2)$		A1	CSO		
<b>Alternative Solutions</b>					
$y' = 4 + 4 \tan^2 4x = 4 + 4 \frac{\sin^2 4x}{\cos^2 4x}$					
$y'' = 4 \times$		(M1)		$\frac{A \cos^3 4x \pm B \sin^3 4x}{\cos^4 4x}$ where $A$ and $B$ are	
$\left[ \frac{\cos^2 4x \cdot 2 \sin 4x \cdot 4 \cos 4x + \sin^2 4x \cdot 2 \cos 4x \cdot 4 \sin 4x}{\cos^4 4x} \right]$		(m1)		constants or trig functions. Where $A$ is $m \sin 4x$ and $B$ is $n \cos 4x$	
$= \frac{4 \times 8 \sin 4x \cos 4x [\cos^2 4x + \sin^2 4x]}{\cos^4 4x}$		(A1F)		ft $8 \times$ their $p$ from part (a)	
$= 32 \tan 4x \sec^2 4x$	(m1)		$k \tan 4x \sec^2 4x$		
$= 32y(1 + y^2)$	(A1)		CSO		
<b>or</b>					
$\frac{dy}{dx} = 4 \sec^2 4x$					
$\frac{d^2 y}{dx^2} = 4 \times 2 \sec 4x \cdot 4 \sec 4x \tan 4x$	(M1)		$A \sec 4x \times f(4x)$		
$= 32 \sec^2 4x \tan 4x$	(m1)		$f(4x) = B \sec 4x \tan 4x$		
$= 32(1 + \tan^2 4x) \tan 4x$	(A1F)		ft $8 \times$ their $p$ from part (a)		
$= 32y(1 + y^2)$	(m1)		Previous two method marks must have been earned		
	(A1)		CSO		

## MPC3 (cont)

Q	Solution	Marks	Total	Comments
7(b) or	$\frac{dy}{dx} = 4(1 + \tan^2 4x)$ $u = \tan 4x \quad \frac{dy}{dx} = 4 + 4u^2$ $\frac{d^2y}{dx^2} = (8)u \frac{du}{dx}$ $\frac{du}{dx} = 4 + 4 \tan^2 4x = 4 + 4u^2$ $\frac{d^2y}{dx^2} = 8u(4 + 4u^2)$ $= 32u(1 + u^2)$ $= 32y(1 + y^2)$	(M1) (m1) (A1) (m1) (A1)		
	<b>Total</b>		<b>8</b>	
8(a)	$\int x \sin(2x-1) dx$ $u = x \quad \frac{dv}{dx} = \sin(2x-1)$ $\frac{du}{dx} = 1 \quad v = -\frac{1}{2} \cos(2x-1)$ $(f=) -\frac{x}{2} \cos(2x-1)$ $-\int -\frac{1}{2} \cos(2x-1) (dx)$ $= -\frac{x}{2} \cos(2x-1) + \frac{1}{2} \int \cos(2x-1) (dx)$ $= -\frac{x}{2} \cos(2x-1) + \frac{1}{4} \sin(2x-1) + c$	M1 A1 m1 A1 A1	5	$\int \sin f(x), \frac{d}{dx}(x)$ attempted All correct – condone omission of brackets correct substitution of their terms into parts All correct – condone omission of brackets CSO condone missing + c and dx Condone missing brackets around 2x – 1 if recovered in final line ISW
(b)	$u = 2x-1$ $'du = 2 dx'$ $\int \frac{x^2}{2x-1} dx = \int \frac{(u+1)^2}{4u} \frac{du}{2}$ $= \left(\frac{1}{8}\right) \int \frac{u^2 + 2u + 1}{u} du$ $= \left(\frac{1}{8}\right) \int u + 2 + \frac{1}{u} du$ $= \left(\frac{1}{8}\right) \left[ \frac{u^2}{2} + 2u + \ln u \right]$ $= \frac{1}{8} \left[ \frac{(2x-1)^2}{2} + 2(2x-1) + \ln(2x-1) \right] + c$	M1 m1 A1 A1 B1 A1	6	OE All in terms of $u$ All correct PI from later working or $\left(\frac{1}{8}\right) \left[ \frac{(u+2)^2}{2} + \ln u \right]$ or $= \frac{1}{8} \left[ \frac{(2x+1)^2}{2} + \ln(2x-1) \right] + c$ CSO condone missing + c only ISW
	<b>Total</b>		<b>11</b>	
	<b>TOTAL</b>		<b>75</b>	