Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination June 2013

Mathematics

MFP3

Unit Further Pure 3

Monday 10 June 2013 9.00 am to 10.30 am

For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

Time allowed

• 1 hour 30 minutes

Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.



Answer all questions.

Answer each question in the space provided for that question.

1 It is given that y(x) satisfies the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{f}(x, y)$$

where

$$f(x, y) = (x - y)\sqrt{x + y}$$

and

$$y(2) = 1$$

Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where $k_1 = hf(x_r, y_r)$ and $k_2 = hf(x_r + h, y_r + k_1)$ and h = 0.2, to obtain an approximation to y(2.2), giving your answer to three decimal places. (5 marks)

QUESTION PART REFERENCE	Answer space for question 1
•••••	
••••••	
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	Answer space for question 1
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



The Cartesian equation of a circle is $(x+8)^2 + (y-6)^2 = 100$.

Using the origin O as the pole and the positive x-axis as the initial line, find the polar equation of this circle, giving your answer in the form $r = p \sin \theta + q \cos \theta$.

(4 marks)

QUESTION PART REFERENCE	Answer space for question 2
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 2
REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



3 (a) Find the values of the constants a, b and c for which $a + bx + cxe^{-3x}$ is a particular integral of the differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - 3y = 3x - 8e^{-3x}$$
 (5 marks)

- (b) Hence find the general solution of this differential equation. (3 marks)
- (c) Hence express y in terms of x, given that y = 1 when x = 0 and that $\frac{dy}{dx} \to -1$ as $x \to \infty$.

QUESTION PART REFERENCE	Answer space for question 3
•••••	



QUESTION PART REFERENCE	Answer space for question 3
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 3
••••••	
•••••	
••••••	
•••••	
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 3
••••••	
••••••	
•••••	
••••••	
•••••	
••••••	
••••••	
••••••	
• • • • • • • • • •	•••••••••••••••••••••••••••••••



4 Evaluate the improper integral

$$\int_0^\infty \left(\frac{2x}{x^2 + 4} - \frac{4}{2x + 3} \right) \mathrm{d}x$$

showing the limiting process used and giving your answer in the form $\ln k$, where k is a constant. (6 marks)

QUESTION	Anguran angga fan arrastian A
QUESTION PART REFERENCE	Answer space for question 4
•••••	
•••••	
•••••	
•••••	
,	
•••••	



QUESTION PART REFERENCE	Answer space for question 4
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



- **5 (a)** Differentiate ln(ln x) with respect to x. (1 mark)
 - (b) (i) Show that $\ln x$ is an integrating factor for the first-order differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} + \frac{1}{x \ln x} y = 9x^2, \quad x > 1$$
 (2 marks)

(ii) Hence find the solution of this differential equation, given that $y = 4e^3$ when x = e. (6 marks)

QUESTION PART REFERENCE	Answer space for question 5



QUESTION PART REFERENCE	Answer space for question 5
•••••	
•••••	



6		It is given that $y = (4 + \sin x)^{\frac{1}{2}}$.	
(a	1)	Express $y \frac{dy}{dx}$ in terms of $\cos x$.	(2 marks)
(b))	Find the value of $\frac{d^3y}{dx^3}$ when $x = 0$.	(5 marks)
(с	;)	Hence, by using Maclaurin's theorem, find the first four terms in the expans	sion, in
		ascending powers of x, of $(4 + \sin x)^{\frac{1}{2}}$.	(2 marks)
QUESTION PART REFERENCE	Ans	wer space for question 6	
••••••			
••••••			
••••••			
••••••			
•••••			
•••••			•••••
•••••			



QUESTION PART REFERENCE	Answer space for question 6



7 A differential equation is given by

$$\sin^2 x \frac{d^2 y}{dx^2} - 2\sin x \cos x \frac{dy}{dx} + 2y = 2\sin^4 x \cos x, \quad 0 < x < \pi$$

(a) Show that the substitution

$$y = u \sin x$$

where u is a function of x, transforms this differential equation into

$$\frac{\mathrm{d}^2 u}{\mathrm{d}x^2} + u = \sin 2x \tag{5 marks}$$

(b) Hence find the general solution of the differential equation

$$\sin^2 x \frac{\mathrm{d}^2 y}{\mathrm{d}x^2} - 2\sin x \cos x \frac{\mathrm{d}y}{\mathrm{d}x} + 2y = 2\sin^4 x \cos x$$

giving your answer in the form y = f(x).

(6 marks)

QUESTION PART REFERENCE	Answer space for question 7



QUESTION PART REFERENCE	Answer space for question 7




QUESTION PART REFERENCE	Answer space for question 7
••••••	
••••••	
••••••	
•••••	
•••••	
••••••	



QUESTION PART REFERENCE	Answer space for question 7
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



8 The diagram shows a sketch of a curve and a circle.



The polar equation of the curve is

$$r = 3 + 2\sin\theta$$
, $0 \le \theta \le 2\pi$

The circle, whose polar equation is r = 2, intersects the curve at the points P and Q, as shown in the diagram.

- (a) Find the polar coordinates of P and the polar coordinates of Q. (3 marks)
- (b) A straight line, drawn from the point P through the pole O, intersects the curve again at the point A.
 - (i) Find the polar coordinates of A. (2 marks)
 - (ii) Find, in surd form, the length of AQ. (3 marks)
 - (iii) Hence, or otherwise, explain why the line AQ is a tangent to the circle r=2.

 (2 marks)
- Find the area of the shaded region which lies inside the circle r=2 but outside the curve $r=3+2\sin\theta$. Give your answer in the form $\frac{1}{6}(m\sqrt{3}+n\pi)$, where m and n are integers.

QUESTION	
PART	Answer space for question 8
REFERENCE	



QUESTION PART REFERENCE	Answer space for question 8
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 8
•••••	
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 8
••••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
	END OF QUESTIONS



