



General Certificate of Education

Mathematics 6360

MFP2 Further Pure 2

Mark Scheme

2006 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key To Mark Scheme And Abbreviations Used In Marking

| | | | |
|--------------|--|-----|----------------------------|
| M | mark is for method | | |
| m or dM | mark is dependent on one or more M marks and is for method | | |
| A | mark is dependent on M or m marks and is for accuracy | | |
| B | mark is independent of M or m marks and is for method and accuracy | | |
| E | mark is for explanation | | |
| √ or ft or F | follow through from previous incorrect result | MC | mis-copy |
| CAO | correct answer only | MR | mis-read |
| CSO | correct solution only | RA | required accuracy |
| AWFW | anything which falls within | FW | further work |
| AWRT | anything which rounds to | ISW | ignore subsequent work |
| ACF | any correct form | FIW | from incorrect work |
| AG | answer given | BOD | given benefit of doubt |
| SC | special case | WR | work replaced by candidate |
| OE | or equivalent | FB | formulae book |
| A2,1 | 2 or 1 (or 0) accuracy marks | NOS | not on scheme |
| -x EE | deduct x marks for each error | G | graph |
| NMS | no method shown | c | candidate |
| PI | possibly implied | sf | significant figure(s) |
| SCA | substantially correct approach | dp | decimal place(s) |

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.


MFP2

| Q | Solution | Marks | Total | Comments |
|--------------|---|--------------------------|----------|--|
| 1(a) | $r^2 + r - 1 = A(r^2 + r) + B$ $A = 1, B = -1$ | M1 A1 A1F | 3 | Any correct method ft B if incorrect A and vice versa Or $\frac{r^2 + r - 1}{r^2 + r} = 1 - \frac{1}{r(r+1)}$ B1 $= 1 - \left(\frac{1}{r} - \frac{1}{r+1}\right)$ M1A1 |
| (b) | $r = 1 \quad 1 - \frac{1}{1} + \frac{1}{2}$ $r = 2 \quad 1 - \frac{1}{2} + \frac{1}{3}$ $r = 99 \quad 1 - \frac{1}{99} + \frac{1}{100}$ $\text{Sum} = 98 + \frac{1}{100}$ $= 98.01$ | M1 A1F m1 A1F | 4 | Do not allow M1 if merely $\sum \frac{1}{r} - \sum \frac{1}{r+1}$ is summed A1 for suitable (3 at least) number of rows Must have 98 or 99 OE Allow correct answer with no working 4 marks |
| Total | | | 7 | |
| 2(a) | $\dot{x} = 1 - t^2, \dot{y} = 2t$ $\dot{x}^2 + \dot{y}^2 = (1 - t^2)^2 + 4t^2$ $= (1 + t^2)^2$ | B1 M1 A1 | 3 | AG; must be intermediate line |
| (b) | $S = 2\pi \int_1^2 (1 + t^2) t^2 dt$ $= 2\pi \left[\frac{t^3}{3} + \frac{t^5}{5} \right]_1^2$ $= 2\pi \left[\frac{8}{3} + \frac{32}{5} - \frac{1}{3} - \frac{1}{5} \right]$ $= \frac{256\pi}{15}$ | M1A1 m1 A1F A1F | 5 | Must be correct substitutions for M1 Allow if one term integrated correctly Any form |
| Total | | | 8 | |

MFP2 (cont)

| Q | Solution | Marks | Total | Comments |
|----------------|--|--------------|--------------|--|
| 3(a)(i) | $\frac{e^k + e^{-k}}{2} - \frac{3(e^k - e^{-k})}{2} = -1$ | M1 | 3 | Allow if 2's are missing or if coshx and sinhx interchanged AG Condone x instead of k |
| | $-2e^k + 4e^{-k} = -2$ | A1 | | |
| | $e^{2k} - e^k - 2 = 0$ | A1 | | |
| (ii) | $(e^k + 1)(e^k - 2) = 0$ | M1 | 4 | Must state something to earn E1. Do not accept ignoring or crossing out. |
| | $e^k \neq -1$ | E1 | | |
| | $e^k = 2$ | A1 | | |
| | $k = \ln 2$ | A1F | | |
| (b)(i) | $\cosh x = 3 \sinh x$ or in terms of e^x | M1 | 4 | CAO |
| | $\tanh x = \frac{1}{3}$ or $2e^x = 4e^{-x}$ | A1 | | |
| | $x = \frac{1}{2} \ln \left(\frac{1 + \frac{1}{3}}{1 - \frac{1}{3}} \right)$ or $e^{2x} = 2$ | A1F | | |
| | $x = \frac{1}{2} \ln 2$ | A1 | | |
| (ii) | $\frac{dy}{dx} = \sinh x - 3 \cosh x$ or $-e^x - 2e^{-x}$ | M1 | 3 | Must give a reason |
| | $= 0$ when $\tanh x = 3$ or $e^{2x} = -2$ | A1 | | |
| | Correct reason | E1 | | |
| (iii) | $\frac{d^2y}{dx^2} = y = 0$ at $\left(\frac{1}{2} \ln 2, 0\right)$ ie one point | B1F | 1 | |
| Total | | | 15 | |

MFP2 (cont)

| Q | Solution | Marks | Total | Comments |
|---------|--|----------------|-----------|--|
| 4 |  | | | |
| (a)(i) | Circle Correct centre Enclosing the origin | B1 B1 B1 | 3 | |
| (ii) | Half line Correct starting point Correct angle | B1 B1 B1 | 3 | |
| (b) | Correct part of the line indicated | B1F | 1 | |
| | Total | | 7 | |
| 5(a)(i) | $\alpha + \beta + \gamma = 4i$ | B1 | 1 | |
| (ii) | $\alpha\beta\gamma = 4 - 2i$ | B1 | 1 | |
| (b)(i) | $\alpha + \alpha = 4i, \alpha = 2i$ | B1 | 1 | AG |
| (ii) | $\beta\gamma = \frac{4-2i}{2i} = -2i - 1$ | M1 A1 | 2 | Some method must be shown, eg $\frac{2}{i} - 1$ AG |
| (iii) | $q = \alpha\beta + \beta\gamma + \gamma\alpha$ $= \alpha(\beta + \gamma) + \beta\gamma$ $= 2i \cdot 2i - 2i - 1 = -2i - 5$ | M1 M1 A1 | 3 | Or $\alpha^2 + \beta\gamma$, ie suitable grouping AG |
| (c) | Use of $\beta + \gamma = 2i$ and $\beta\gamma = -2i - 1$ $z^2 - 2iz - (1 + 2i) = 0$ | M1 A1 | 2 | Elimination of say γ to arrive at $\beta^2 - 2i\beta - (1 + 2i) = 0$ M1A0 unless also some reference to γ being a root AG |
| (d) | $f(-1) = 1 + 2i - 1 - 2i = 0$ $\beta = -1, \gamma = 1 + 2i$ | M1 A1A1 | 3 | For any correct method A1 for each answer |
| | Total | | 13 | |

MFP2 (cont)

| Q | Solution | Marks | Total | Comments |
|-------------|--|--------------|--------------|---|
| 6(a) | $f(n+1) - 8f(n) = 15^{n+1} - 8^{n-1}$ $- 8(15^n - 8^{n-2})$ $= 15^{n+1} - 8 \cdot 15^n$ $= 15^n (15 - 8)$ $= 7 \cdot 15^n$ | M1A1 | 4 | For multiples of powers of 15 only For valid method ie not using 120^n etc |
| (b) | Assume $f(n)$ is $M(7)$ | M1 | | |
| | Then $f(n+1) - 8f(n) = 7 \times 15^n$ | M1 | | |
| | $f(n+1) = M(7) + M(7)$ $= M(7)$ | A1 | | |
| | $n = 2: f(n) = 15^2 - 8^0 = 224$ $= 7 \times 32$ | B1 | 4 | $n = 1$ B0 Must score previous 3 marks to be awarded E1 |
| | $P(n) \Rightarrow P(n+1)$ and $P(2)$ true | E1 | | |
| | Total | | 8 | |

MFP2 (cont)

| Q | Solution | Marks | Total | Comments |
|--------|---|--------------------|-----------|---|
| 7(a) | $z = e^{\frac{2k\pi i}{6}}, k = 0, \pm 1, \pm 2, 3$ | M1 A2,1,0 | 3 | OE M1A1 only if: (1) range for k is incorrect eg 0,1,2,3,4,5 (2) i is missing |
| (b)(i) | $\frac{w^2 - 1}{w} = w - \frac{1}{w} = 2i \sin \theta$ | M1A1 | 2 | AG |
| (ii) | $\frac{w}{w^2 - 1} = \frac{1}{2i \sin \theta}$ $= -\frac{i}{2 \sin \theta}$ | M1 A1 | 2 | AG |
| (iii) | $\frac{2i}{w^2 - 1} = \frac{-2iw^{-1}i}{2 \sin \theta}$ $= \frac{1}{\sin \theta} (\cos \theta - i \sin \theta)$ $= \cot \theta - i$ | M1 A1 A1 | 3 | AG Or for $\frac{1}{\sin \theta e^{i\theta}}$ |
| (iv) | $z = \frac{2i}{w^2 - 1}$ Or $z + 2i = \frac{2i}{w^2 - 1} + 2i$ $z + 2i = zw^2$ | M1 A1 | 2 | AG ie any correct method |
| (c)(i) | No coefficient of z^6 | E1 | 1 | |
| (ii) | $(w^2)^6 = 1 \quad w^2 = e^{\frac{k\pi i}{3}}$ $z = \cot \frac{k\pi}{6} - i, \quad k = \pm 1, \pm 2, 3$ | B1 M1 A2,1,0 | 4 | Alternatively: $z + 2i = e^{\frac{k\pi i}{3}} z$ B1 $z = \frac{2i}{e^{\frac{k\pi i}{3}} - 1}$ M1 roots A2,1,0 (NB roots are $\pm \sqrt{3} - i; \pm \frac{1}{\sqrt{3}} - i; -i$) |
| | Total | | 17 | |
| | TOTAL | | 75 | |