General Certificate of Education January 2007 Advanced Subsidiary Examination

MATHEMATICS Unit Mechanics 1A

MM1A/W



Friday 12 January 2007 9.00 am to 10.15 am

For this paper you must have:

• an 8-page answer book

• the **blue** AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 15 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MM1A/W.
- Answer all questions.
- Show all necessary working; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of calculators should be given to three significant figures, unless stated otherwise.
- Take $g = 9.8 \text{ m s}^{-2}$, unless stated otherwise.

Information

- The maximum mark for this paper is 60.
- The marks for questions are shown in brackets.
- Unit Mechanics 1A has a written paper and coursework.

Advice

• Unless stated otherwise, you may quote formulae, without proof, from the booklet.

MM1A/W

Answer all questions.

1 Two particles A and B have masses of 3 kg and 2 kg respectively. They are moving along a straight horizontal line towards each other. Each particle is moving with a speed of 4 m s^{-1} when they collide.



- (a) If the particles coalesce during the collision to form a single particle, find the speed of the combined particle after the collision. (3 marks)
- (b) If, after the collision, A moves in the same direction as before the collision with speed 0.4 m s^{-1} , find the speed of B after the collision. (3 marks)
- 2 A motorcycle accelerates uniformly along a straight horizontal road so that, when it has travelled 20 metres, its velocity has increased from 12 m s^{-1} to 16 m s^{-1} .
 - (a) Find the acceleration of the motorcycle. (3 marks)
 - (b) Find the time that it takes for the motorcycle to travel this distance. (3 marks)
- 3 A car, of mass 1500 kg, is towing a caravan, of mass 900 kg, along a straight horizontal road. The caravan is connected to the car by a horizontal tow bar. Resistance forces of magnitudes 400 N and 800 N act on the car and caravan respectively. The acceleration of the car and caravan is 0.8 m s^{-2} .



(a) Show that the magnitude of the force that the car exerts on the caravan is 1520 N.

(3 marks)

(b) Find the magnitude of the driving force produced by the car's engine. (3 marks)

4 A cricket ball is hit from the floor of a sports hall, which has a height of 6 metres. The initial velocity of the ball is 20 m s^{-1} at an angle of 60° above the horizontal.

Assume that the cricket ball is a particle which moves only under the influence of gravity.

- (a) Show that the ball hits the ceiling of the sports hall approximately 0.389 seconds after it was hit. (5 marks)
- (b) Find the horizontal distance travelled by the ball before it hits the ceiling. (2 marks)
- (c) Find the speed of the ball just before it hits the ceiling. (5 marks)
- 5 A girl in a boat is rowing across a river, in which the water is flowing at 0.1 m s^{-1} . The velocity of the boat relative to the water is 0.3 m s^{-1} and is perpendicular to the bank, as shown in the diagram.



The width of the river is 15 metres	(5 marks)
(i) Find the time that it takes the boat to cross the river.	(2 marks)
	Find the magnitude of the resultant velocity of the boat.Find the acute angle between the resultant velocity and the bank.The width of the river is 15 metres.(i) Find the time that it takes the boat to cross the river.

- (ii) Find the total distance travelled by the boat as it crosses the river. (2 marks)

Turn over for the next question

6 A trolley, of mass 100 kg, rolls at a constant speed along a straight line down a slope inclined at an angle of 4° to the horizontal.

Assume that a constant resistance force, of magnitude P newtons, acts on the trolley as it moves. Model the trolley as a particle.

- (a) Draw a diagram to show the forces acting on the trolley. (1 mark)
- (b) Show that P = 68.4 N, correct to three significant figures. (3 marks)
- (c) (i) Find the acceleration of the trolley if it rolls down a slope inclined at 5° to the horizontal and experiences the same constant force of magnitude P that you found in part (b). (4 marks)
 - (ii) Make one criticism of the assumption that the resistance force on the trolley is constant. (1 mark)
- 7 A particle is initially at the origin, where it has velocity $(5i 2j) m s^{-1}$. It moves with a constant acceleration $a m s^{-2}$ for 10 seconds to the point with position vector 75i metres.
 - (a) Show that a = 0.5i + 0.4j. (3 marks)
 - (b) Find the position vector of the particle 8 seconds after it has left the origin. (3 marks)
 - (c) Find the position vector of the particle when it is travelling parallel to the unit vector **i**. (6 marks)

END OF QUESTIONS