



**General Certificate of Education**

**Mathematics 6360**

**MFP2      Further Pure 2**

**Mark Scheme**

*2010 examination - January series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2010 AQA and its licensors. All rights reserved.

#### COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

---

**Key to mark scheme and abbreviations used in marking**

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

**No Method Shown**


Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

MFP2				
Q	Solution	Marks	Total	Comments
1(a)	$\text{LHS} = \frac{1}{4}(e^x + e^{-x})^2 - \frac{1}{4}(e^x - e^{-x})^2$ Correct expansion of either square Shown equal to 1	M1 A1 A1	3	AG
(b)(i)	$8\cosh^2 x - 3$	B1	1	
(ii)	Sketch of $y = \cosh x$	B1	1	Must cross $y$ -axis above $x$ -axis
(iii)	$\cosh x = (\pm)1.25$  $x = \ln(1.25 + \sqrt{1.25^2 - 1})$ $= \ln 2$ $\ln \frac{1}{2}$ by symmetry	B1F  M1 A1F A1F	4	OE; ft errors in (b)(i); allow $\pm$ missing  Accept $-\ln 2$ written straight down  Alternatively, if solved by using $e^{2x} - 2.5e^x + 1 = 0$ , allow M1 for $x = \ln\left(\frac{2.5 \pm \sqrt{2.5^2 - 4}}{2}\right)$
<b>Total</b>			<b>9</b>	
2				
(a)(i)	Circle  Correct centre  Touching $y$ -axis	B1  B1 B1	3	$x$ -coordinate $\approx -2 \times y$ -coordinate in correct quadrant; condone $(4, -2i)$
(ii)	Straight line parallel to $x$ -axis  through $(0, 1)$	B1 B1 B1	3	Assume $(0, 1)$ if distance up $y$ -axis is half distance to top of circle; no other shading outside circle
(b)	Shading: inside circle above line	B1F B1F	2	Whole question reflected in $x$ -axis loses 2 marks
<b>Total</b>			<b>8</b>	

## MFP2 (cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	$\beta = 2 - 2\sqrt{3}i$	B1	1	
(ii)	$\alpha\beta\gamma = -8$ $\alpha\beta = 16$ $\gamma = -\frac{1}{2}$	M1 B1 A1	3	Allow for +8 but not $\pm 16$
(iii)	Either $\frac{-p}{2} = \alpha + \beta + \gamma$ or $\frac{q}{2} = \alpha\beta + \beta\gamma + \gamma\alpha$  $p = -7, q = 28$	M1  A1F, A1F	3	SC if failure to divide by 2 throughout, allow M1A1 for either $p$ or $q$ correct ft  ft incorrect $\gamma$
	<b>Alternative to (a)(ii) and (a)(iii):</b> $(z^2 - 4z + 16)(az + b)$ $\alpha\beta = 16$ $a = 2, b = +1, \gamma = -\frac{1}{2}$ Equating coefficients $p = -7$ $q = 28$	(M1) (B1) (A1) (M1) (A1F) (A1F)		
(b)(i)	$r = 4, \theta = \frac{\pi}{3}$	B1, B1	2	
(ii)	$(2 + 2\sqrt{3}i)^n = \left(4e^{\frac{\pi i}{3}}\right)^n$ $= 4^n \left(\cos \frac{n\pi}{3} + i \sin \frac{n\pi}{3}\right)$	M1 A1	2	AG
(iii)	$(2 - 2\sqrt{3}i)^n = 4^n \left(\cos \frac{n\pi}{3} - i \sin \frac{n\pi}{3}\right)$ $\alpha^n + \beta^n + \gamma^n = 4^n \left(\cos \frac{n\pi}{3} + i \sin \frac{n\pi}{3}\right)$ $+ 4^n \left(\cos \frac{n\pi}{3} - i \sin \frac{n\pi}{3}\right) + \left(-\frac{1}{2}\right)^n$ $= 2^{2n+1} \cos \frac{n\pi}{3} + \left(-\frac{1}{2}\right)^n$	B1  M1 A1	3	AG
	<b>Total</b>		<b>14</b>	

## MFP2 (cont)

Q	Solution	Marks	Total	Comments
<b>4(a)</b>	$\frac{dx}{dt} = \sinh 2t$	B1		
	$\frac{dy}{dt} = 2 \cosh t$	B1		
	$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \sinh^2 2t + 4 \cosh^2 t$	M1		
	Use of $\sinh 2t = 2 \sinh t \cosh t$	m1		Or other correct formula for double angle
	$= 4 \cosh^2 t (\sinh^2 t + 1)$	A1		For taking out factor
	$= 4 \cosh^4 t$	A1F	6	ft errors of sign in $\frac{dx}{dt}$ or $\frac{dy}{dt}$
<b>(b)(i)</b>	$S = 2\pi \int_0^1 2 \sinh t \cdot 2 \cosh^2 t \, dt$	M1		Using the value obtained in (a)
	$= 8\pi \int_0^1 \sinh t \cdot \cosh^2 t \, dt$	A1	2	AG
<b>(ii)</b>	$S = 8\pi \left[ \frac{\cosh^3 t}{3} \right]_0^1$	M1		
	$= \frac{8\pi}{3} [\cosh^3 1 - 1]$	A1	2	OE eg $\frac{\pi}{3} \left( \left( e + \frac{1}{e} \right)^3 - 8 \right)$
<b>Total</b>			<b>10</b>	
<b>5(a)(i)</b>	$u_1 = S_1 = 1^2 \cdot 2 \cdot 3 = 6$	B1	1	AG
	<b>(ii)</b> $u_2 = S_2 - S_1 = 42$	B1	1	AG
<b>(iii)</b>	$u_n = S_n - S_{n-1}$	M1		
	$= n^2(n+1)(n+2) - (n-1)^2 n(n+1)$	A1		
	$= n(n+1)(4n-1)$	A1	3	AG
<b>(b)</b>	$\sum_{r=n+1}^{2n} u_r = S_{2n} - S_n$	M1		
	$= (2n)^2(2n+1)(2n+2) - n^2(n+1)(n+2)$	A1		
	$= 3n^2(n+1)(5n+2)$	A1	3	AG
<b>Total</b>			<b>8</b>	

## MFP2 (cont)

Q	Solution	Marks	Total	Comments
<b>6(a)</b>	$t = \tan \theta \quad dt = \sec^2 \theta \, d\theta$	B1		OE
	$I = \int \frac{dt}{(9 \cos^2 \theta + \sin^2 \theta) \sec^2 \theta}$	M1		OE
	$= \int \frac{dt}{t^2 + 9}$	A1	3	AG
<b>(b)</b>	$I = \left[ \frac{1}{3} \tan^{-1} \frac{t}{3} \right]_0^{\sqrt{3}}$	M1		M1 for $\tan^{-1}$
	$\frac{1}{3} \tan^{-1} \frac{\sqrt{3}}{3} \text{ or } \frac{1}{3} \tan^{-1} \frac{1}{\sqrt{3}}$	A1		
	$= \frac{\pi}{18}$	A1	3	AG
	<b>Total</b>		<b>6</b>	
<b>7(a)</b>	Assume true for $n = k$			
	$u_{k+1} = 2(3 \times 2^{k-1} - 1) + 1$	M1A1		
	$= 3 \times 2^k - 1$	A1		$2^{(k-1)+1}$ not necessarily seen
	True for $n = 1$ shown	B1		
	Method of induction clearly expressed	E1	5	Provided all 4 previous marks earned
<b>(b)</b>	$\sum_{r=1}^n u_r = \sum_{r=1}^n 3 \times 2^{r-1} - n$			
	$= 3(2^n - 1) - n$	M1A1		M1 for summation, ie recognition of a GP
	$= u_{n+1} - (n + 2)$	A1	3	AG
	<b>Total</b>		<b>8</b>	

## MFP2 (cont)

Q	Solution	Marks	Total	Comments
8(a)(i)	$\left(e^{\frac{2\pi i}{7}}\right)^7 = e^{2\pi i} = 1$	B1	1	Or $z^7 = e^{2k\pi i}$ $z = e^{\frac{2k\pi i}{7}}$ $k = 1$
(ii)	Roots are $\omega^2, \omega^3, \omega^4, \omega^5, \omega^6$	M1A1	2	OE; M1A0 for incomplete set SC B1 for a set of correct roots in terms of $e^{i\theta}$
(b)	Sum of roots considered $= 0$	M1 A1	2	$\left\{ \text{or } \sum_{r=0}^6 \omega^r = \frac{\omega^7 - 1}{\omega - 1} = 0 \right.$
(c)(i)	$\omega^2 + \omega^5 = e^{\frac{4\pi i}{7}} + e^{\frac{10\pi i}{7}}$ $= e^{\frac{4\pi i}{7}} + e^{-\frac{4\pi i}{7}}$ $= 2\cos\frac{4\pi}{7}$	M1 A1 A1	3	Or $\cos\frac{4\pi}{7} + i\sin\frac{4\pi}{7} + \cos\frac{4\pi}{7} - i\sin\frac{4\pi}{7}$ AG
(ii)	$\omega + \omega^6 = 2\cos\frac{2\pi}{7}$ ; $\omega^3 + \omega^4 = 2\cos\frac{6\pi}{7}$ Using part (b) Result	B1,B1 M1 A1	4	Allow these marks if seen earlier in the solution AG
	<b>Total</b>		<b>12</b>	
	<b>TOTAL</b>		<b>75</b>	