



**General Certificate of Education (A-level)  
June 2012**

**Mathematics**

**MS/SS1A**

**(Specification 6360)**

**Statistics 1A**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2012 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

## MS/SS1A/W

Q	Solution	Marks	Total	Comments
<b>1</b> <b>(a)</b>	Mean = <u>29.2</u> Mean = <u>29 to 30</u>	B2 (B1)	2	CAO AWFW ( $\Sigma fx = 730$ )
<b>Note</b>	If B0 then clear use of $\Sigma fx/25 \Rightarrow$ M1			
<b>(b)</b>	Mean = $\frac{(219 \times 5.50) + (73 \times 6.50) + (438 \times 9.5)}{25}$  $= \frac{5840}{25} = \underline{\underline{\pounds 233 \text{ to } \pounds 234}}$	M1  A1	  2	Allow divisor of 730 $\Rightarrow$ 8  AWFW; ignore units ( $\pounds 233.60$ )
<b>(c)</b>	29.2 > 28 <b>and/or</b> ( $\pounds 233$ to $\pounds 234$ ) < $\pounds 250$ so do <b>not purchase</b> business	B1  Adep1	  2	Either  Dependent on previous A1 and comparison with $\pounds 250$
	<b>Total</b>		<b>6</b>	

## MS/SS1A/W (cont)

Q	Solution	Marks	Total	Comments
2 (a)	$b$ (gradient) = <u>2.27</u> $b$ (gradient) = <u>2.2 to 2.3</u> $a$ (intercept) = <u>4.16 to 4.2</u> $a$ (intercept) = <u>3 to 7</u>	B2 (B1) B2 (B1)		AWRT (2.27075) AWFW <i>Treat rounding of correct answers as ISW</i> AWFW (4.16981) AWFW
	Attempt at $\sum x$ $\sum x^2$ $\sum y$ & $\sum xy$ ( $\sum y^2$ ) <b>or</b> Attempt at $S_{xx}$ & $S_{xy}$ ( $S_{yy}$ ) Attempt at <b>correct</b> formula for $b$ (gradient)	(M1) (m1)		480 24500 1140 & <b>57635</b> (135908) (all 4 attempted) 5300 & <b>12035</b> (27608) (both attempted)
	$b$ (gradient) = <u>2.27</u> $a$ (intercept) = <u>4.16 to 4.2</u>	(A1) (A1)	4	AWRT AWFW
<b>Notes</b>	<p>1 Values of <math>a</math> and <math>b</math> interchanged and equation <math>y = ax + b</math> stated in (a) <math>\Rightarrow</math> max of 4 marks</p> <p>2 Values of <math>a</math> and <math>b</math> interchanged and equation <math>y = a + bx</math> stated in (a) <math>\Rightarrow</math> 0 marks</p> <p>3 Values are <b>not</b> identified or simply <math>a = \#</math> and <math>b = \#</math>, then 2.2 to 2.3 <math>\Rightarrow</math> B1 and 3 to 7 <math>\Rightarrow</math> B1 but accept, for example, as identification, [<math>a = \#, b = \#</math> with <math>y = a + bx</math> but no substitution for <math>a</math> &amp; <math>b</math>] <b>or</b> [intercept(<math>a</math>) = #, gradient(<math>b</math>) = #]</p> <p>4 <math>b = \frac{2407}{1060}</math> CAO <math>\Rightarrow</math> B2, otherwise B1 if fraction equates to 2.2 to 2.3 (Notes 1, 2 &amp; 3 also apply)  <math>a = \frac{221}{53}</math> CAO <math>\Rightarrow</math> B2, otherwise B1 if fraction equates to 3 to 7 (Notes 1, 2 &amp; 3 also apply)</p> <p>5 Some/all of marks can be scored in (b), and in c(ii) &amp; (iii), even if some/all of marks are lost in (a) but marks lost in (a) <b>cannot</b> be recouped by subsequent working in (b)</p>			
(b)	Correct <b>straight</b> line drawn on scatter diagram Correct shortened and/or freehand line drawn on scatter diagram	B2 (B1)	2	Line must go from $x \leq 20$ to $x \geq 70$ and fall between the following 2 lines: Lower: (10, 25) (80, 180) Upper: (10, 30) (80, 190)
<b>Notes</b>	<p>1 If B0 but <b>seen</b> correct attempt at <math>\geq 2</math> points even if incorrectly evaluated <math>\Rightarrow</math> M1</p> <p>2 If B0 but <b>no seen</b> evidence to support <math>\geq 2</math> points (correct or incorrect) marked on scatter diagram <math>\Rightarrow</math> M0</p>			
(c)(i)	Correct <b>straight</b> line drawn on scatter diagram Correct shortened and/or freehand line drawn on scatter diagram	B2 (B1)	2	Line must go from $x \leq 20$ to $x \geq 70$ and fall between the following 2 lines: Lower: (10, 60) (80, 75) Upper: (10, 65) (80, 85)
<b>Notes</b>	<p>1 If B0 but <b>seen</b> correct attempt at <math>\geq 2</math> points even if incorrectly evaluated <math>\Rightarrow</math> M1</p> <p>2 If B0 but <b>no seen</b> evidence to support <math>\geq 2</math> points (correct or incorrect) marked on scatter diagram <math>\Rightarrow</math> M0</p>			
(ii)	<u>27 to 29</u>	B1	1	AWFW (calculation $\Rightarrow$ 27.75) Must clearly identify $x$ -value Thus (27 to 29, $y$ -value) $\Rightarrow$ B0
(iii)	At low temperatures more B (than A) dissolves At high temperatures more A (than B) dissolves	B1		Either; OE (eg a comparison using lines and/or data at a specific temperature but <b>not</b> at 0°C)
	Amount increases more rapidly for A (than B) Amount increases more slowly for B (than A)	B1	2	Either; OE Any comments about $b$ or $a$ $\Rightarrow$ B0 Comment about 'rate' must relate to temp
	<b>Total</b>		<b>11</b>	

## MS/SS1A/W (cont)

Q	Solution	Marks	Total	Comments
<b>3</b>				<b>Ratios (eg 194:640) are only penalised by 1 accuracy mark at first correct answer</b>
(a)(i)	$P(B = 3) =$ <u><b>194/640 or 97/320 or 0.303 or 30.3%</b></u>	B1	1	CAO or AWRT (0.303125)
(ii)	$P(T \geq 2) = \frac{172 + 256 + 135}{640}$ or $1 - \frac{77}{640}$ or $\frac{563}{640}$ $=$ <u><b>563/640</b></u> <u><b>or (0.879 to 0.88) or (87.9% to 88%)</b></u>	M1 A1	2	CAO AWFW (0.879688)
(iii)	$P(B = 3 \text{ \& } T \geq 2) =$ $\frac{72 + 99 + 16}{640}$ or $\frac{194 - 7}{640}$ or $\frac{187}{640}$ $=$ <u><b>187/640 or 0.292 or 29.2%</b></u>	M1 A1	2	CAO or AWRT (0.292188)
(iv)	$P(B \leq 3 \mid T = 2) =$ $\frac{(14 + 67 + 72)}{172}$ or $\frac{172 - 19}{172}$ or $\frac{153}{172}$ $=$ <u><b>153/172</b></u> <u><b>or (0.888 to 0.89) or (88.8% to 89%)</b></u>	M1 M1 A1	3	Correct numerator (accept both $\div 640$ ) Correct denominator CAO AWFW (0.889535)
(b)	$P(2T \cap 3T \cap \geq 4T \mid B = 3) = \frac{72}{194} \times \frac{99}{193} \times \frac{16}{192}$  $abc$ multiplied by 6 or 3 $=$ <u><b>0.095 to 0.0952</b></u>	M1 M1 M1 A1	4	Correct 3 values multiplied in numerator Correct 3 values multiplied in denominator $0.371 \times 0.513 \times 0.083$ (all AWRT) $\Rightarrow$ M1 M1 (OE products) $0 < (a, b \text{ \& } c) < 1$ AWFW (0.095187)
<b>Notes</b>	<b>1</b> Incorrect answer with no working $\Rightarrow$ 0 marks <b>2</b> The <b>3 correct fractions/decimals</b> identified but not multiplied (eg added) $\Rightarrow$ M1 M0 M0 A0 <b>3</b> The <b>3 correct fractions/decimals</b> identified together with 0.016 (AWRT) $\Rightarrow$ M1 M1 M0 A0 <b>4</b> A denominator of ${}^{194}C_3 = 1198144 \Rightarrow$ M2 (2 <sup>nd</sup> & 3 <sup>rd</sup> M1 marks)			
	<b>Total</b>		<b>12</b>	

## MS/SS1A/W (cont)

Q	Solution	Marks	Total	Comments
<b>4</b>				
(a)	$B(n, p)$	M1		Used correctly anywhere in question
	$P(C = 2) = \binom{10}{2}(0.275)^2(0.725)^8$	A1		Can be implied by a <b>correct</b> answer
	$= \underline{\mathbf{0.259\ to\ 0.26}}$	A1	3	AWFW (0.25977)
(b)	$n = \underline{\mathbf{40}}$ and $p = \underline{\mathbf{0.35}}$	B1		CAO both; used in (b)
(i)	$P(G \leq 15) = \underline{\mathbf{0.694\ to\ 0.695}}$	B1	2	AWFW (0.6946)
(ii)	$P(10 \leq G \leq 20) = \mathbf{0.9827\ or\ 0.9637}$ ( $p_1$ )	M1		Accept 3 dp rounding or truncation
	<b>MINUS</b> $\mathbf{0.0644\ or\ 0.1215}$ ( $p_2$ )	M1		Accept 3 dp rounding or truncation
	$= \underline{\mathbf{0.918\ to\ 0.92}}$ ( $p_3$ )	A1	3	AWFW (0.9183)
<b>Notes</b>	<b>1</b> $p_3 \leq 0$ or $p_3 \geq 1 \Rightarrow$ M0 M0 A0		<b>4</b>	$p_1 - (1 - p_2) \Rightarrow$ M1 M0 A0
	<b>2</b> $p_2 - p_1 \Rightarrow$ M0 M0 A0		<b>5</b>	$p_1 \times p_2 \Rightarrow$ M1 M0 A0
	<b>3</b> $(1 - p_2) - p_1 \Rightarrow$ M0 M0 A0		<b>6</b>	$(1 - p_2) - (1 - p_1) \Rightarrow$ M1 M1 (A1)
(c)	$n = \underline{\mathbf{50}}$ and $p = \underline{\mathbf{0.85}}$			
	$P(R > 40   p = 0.85) = P(A < 10   p = 0.15)$	M1		Attempt to change from $R$ to $A$
	$= \underline{\mathbf{0.7911\ or\ 0.8801}}$	m1		Accept 3 dp rounding or truncation
	$= \underline{\mathbf{0.79\ to\ 0.792}}$	A1		AWFW (0.7911)
	<b>OR</b>			
	$P(R \leq 40   p = 0.85) = \underline{\mathbf{0.2089\ or\ 0.1199}}$	(M1)		Accept 3 dp rounding or truncation
	$P(R > 40   p = 0.85) = \underline{\mathbf{1 - (0.2089\ or\ 0.1199)}}$	(m1)		Requires '1 -' Can be implied by 0.79 to 0.792 but <b>not</b> by 0.88 to 0.881
	$= \underline{\mathbf{0.79\ to\ 0.792}}$	(A1)	3	AWFW (0.7911)
	<b>Total</b>		<b>11</b>	

## MS/SS1A/W (cont)

Q	Solution	Marks	Total	Comments
<b>5</b>				<b>In (a)(i) &amp; (c), ignore the inclusion of a lower limit of 0; it has no effect on either answer</b>
(a)	<u>Weight, <math>W \sim N(2.75, 0.15^2)</math></u>			
(i)	$P(W < 2.8) = P\left(Z < \frac{2.8 - 2.75}{0.15}\right)$	M1		Standardising 2.8 with 2.75 and 0.15; allow (2.75 – 2.8)
	$= P(Z < \underline{0.33 \text{ or } 1/3})$	A1		AWRT/CAO; ignore inequality and sign May be implied by a <b>correct</b> answer
	$= \underline{0.629 \text{ to } 0.633}$	A1		AWFW (0.63056)
(ii)	$P(W > 2.5) = P(Z > -1.67) = P(Z < +1.67)$	M1		<b>Correct</b> area change May be implied by a <b>correct</b> answer <b>or</b> an answer <b>&gt; 0.5</b>
	$= \underline{0.951 \text{ to } 0.953}$	A1	5	AWFW (0.95221)
(b)	<u>Weight, <math>X \sim N(5.25, 0.20^2)</math></u>			
(i)	$P(5.1 < X < 5.3) = P(Z < 0.25) - P(Z < -0.75)$			
	$= \underline{0.59871}$	B1		Must have diff of 2 probs for each B1 Accept 0.599
	<b>MINUS</b> $[(1 - 0.77337) \text{ or } 0.22663]$	B1	2	Accept 0.773 or 0.227 Accept correct alternative methods AG; do <b>not</b> mark simply on answer
	$= \underline{0.372(08)}$			
(ii)	$P(0 \text{ in } 4) = [1 - 0.372]^4$	M1		Accept $[1 - c's (b)(i)]^4$
	$= 0.628^4 = \underline{0.155 \text{ to } 0.156}$	A1	2	AWFW (0.15554)
(c)	<u>Weight, <math>Y \sim N(10.75, 0.50^2)</math></u>			
	Variance of $\bar{Y}_6 = \underline{0.5^2/6 = 0.0416 \text{ to } 0.0417}$			
	<b>or</b>			
	Sd of $\bar{Y}_6 = \underline{0.5/\sqrt{6} = 0.204}$	B1		CAO or AFWW Stated or used CAO or AWRT
	$P(\bar{Y}_6 < 10.5) = P\left(Z < \frac{10.5 - 10.75}{\sqrt{0.0416}}\right) =$	M1		Standardising 10.5 with 10.75 and $\sqrt{0.0416}$ OE; allow (10.75 – 10.5)
	$P(Z < -1.22) = 1 - P(Z < 1.22) =$	m1		<b>Correct</b> area change May be implied by a correct answer <b>or</b> an answer <b>&lt; 0.5</b> ; but do <b>not</b> award for use of $z = \pm 0.22$
	$1 - (0.88877 \text{ to } 0.89065) = \underline{0.109 \text{ to } 0.112}$	A1	4	AWFW (0.11034) (1 – answer) $\Rightarrow$ B1 M1 max
	<b>Total</b>		<b>13</b>	



## MS/SS1A/W (cont)

Q	Solution	Marks	Total	Comments
<b>6</b> <b>(a)</b>	98% (0.98) $\Rightarrow z = \underline{2.32 \text{ to } 2.33}$	B1		AWFW (2.3263)
	CI for $\mu$ is $\bar{x} \pm z \times \frac{\sigma}{\sqrt{n}}$	M1		Used with $z$ (2.05 to 2.58), $\bar{x}$ (5.05, 5050 or 181.8), $\sigma$ (0.0075, 0.075, 0.75, 7.5 or 75) and $\div\sqrt{n}$ with $n > 1$
	Thus $5.05 \pm 2.3263 \times \frac{0.075}{\sqrt{36}}$	A1		$z$ (2.05 to 2.06 or 2.32 to 2.33 or 2.57 to 2.58), $\bar{x}$ (5.05) & $\sigma$ (0.075) or $\bar{x}$ (5050) & $\sigma$ (75) and $\div\sqrt{36}$ or 35
	Hence $\underline{5.05 \pm 0.03}$ or $\underline{5050 \pm 30}$			CAO/AWRT
	<b>OR</b>	Adep1	4	Dependent on previous A1 so can be scored with $z \neq 2.32 \text{ to } 2.33$ Ignore (absence of) quoted units AWRT to 3sf accuracy
	$\underline{(5.02, 5.08)}$ or $\underline{(5020, 5080)}$			
<b>Note</b>	Use of $t$ (2.43 to 2.72) $\Rightarrow$ B1 B0 M1 A0 A0 max			
<b>(b)</b>	<b>Clear correct comparison of 5 or 5000 with LCL or CI</b> so <b>agree</b> with (first) claim (about mean)	Adep1		Dependent on Adep1 in (a) Must use consistent units
	(8/36 or 0.22 or 22%) v (1/10 or 0.10 or 10%) <b>or</b> $8 \text{ v } 3.6$ (3 to 4)	B1		Mention of a value on LHS <b>and</b> a value on RHS
	so 8/36 OE $>/\neq$ 1/10 OE so <b>disagree</b> with (second) claim (about individuals)	Bdep1	3	Dependent on B1 Explicit comparison of values and correct conclusion
<b>Notes</b>	<b>1</b> It/(claimed) mean/(claimed) value < LCL/CI $\Rightarrow$ Adep0 <b>2</b> 98% have (mean) weights between CLs so ... $\Rightarrow$ Adep0 <b>3</b> Any reference to CI for second claim $\Rightarrow$ B0 Bdep0			Must indicate 5 or 5000 Claim refers to individual bottles
	<b>Total</b>		<b>7</b>	
	<b>TOTAL</b>		<b>60</b>	