

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
TOTAL	



General Certificate of Education
Advanced Level Examination
June 2014

Mathematics

MM05

Unit Mechanics 5

Thursday 12 June 2014 1.30 pm to 3.00 pm

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.
- The **final** answer to questions requiring the use of calculators should be given to three significant figures, unless stated otherwise.
- Take $g = 9.8 \text{ m s}^{-2}$, unless stated otherwise.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.



J U N 1 4 M M 0 5 0 1

Answer **all** questions.

Answer each question in the space provided for that question.

1 A particle moves, with simple harmonic motion, in a straight line between two points, *A* and *B*. During this motion, its maximum speed is 1.3 m s^{-1} . When the particle is 0.2 m from the midpoint of *AB*, its speed is 1.2 m s^{-1} .

Find the distance between the points *A* and *B*.

[6 marks]

QUESTION
PART
REFERENCE

Answer space for question 1



QUESTION
PART
REFERENCE

Answer space for question 1


A large rectangular area containing horizontal dotted lines for writing an answer.

Turn over ►



2

A simple pendulum consists of a small sphere of mass m kg attached to one end of a light inextensible string of length l metres. The other end of the string is attached to a fixed point. The sphere is released from rest with the string taut and at an angle of $\frac{\pi}{10}$ to the downward vertical. When the pendulum has been in motion for t seconds, the angle between the string and the downward vertical is θ , as shown in the diagram.



(a) Show that the motion can be modelled by the differential equation

$$\frac{d^2\theta}{dt^2} = -\frac{g}{l}\theta$$

and state clearly any assumptions that you make.

[4 marks]

(b) The length of the string is 39.2 cm.

(i) Find an expression for θ at time t .

[4 marks]

(ii) Find the time that it takes for θ to decrease from $\frac{\pi}{15}$ to $\frac{\pi}{30}$.

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 2

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



QUESTION
PART
REFERENCE

Answer space for question 2

A large rectangular area with horizontal dotted lines for writing.

Turn over ►



QUESTION
PART
REFERENCE

Answer space for question 2

A large rectangular area containing horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE

Answer space for question 2

A large rectangular area with horizontal dotted lines for writing an answer.

Turn over ►



3 A particle moves under the action of a force which is always directed towards the origin, O , on a curve defined in plane polar coordinates as $r = \frac{3}{1 + \sin \theta}$.

(a) Find r when $\theta = 0$.

[1 mark]

(b) Given that the transverse component of the velocity of the particle is 5 when $\theta = 0$, show that $\dot{\theta} = \frac{5}{3}(1 + \sin \theta)^2$.

[4 marks]

(c) (i) Find \dot{r} in terms of θ .

[2 marks]

(ii) Show that the radial component of acceleration is given by $\frac{k}{r^2}$, and state the value of the constant k .

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 3



QUESTION
PART
REFERENCE

Answer space for question 3

A large rectangular area with horizontal dotted lines for writing an answer.



Turn over ►

QUESTION
PART
REFERENCE

Answer space for question 3

A large rectangular area with horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE


Answer space for question 3

A large rectangular area with horizontal dotted lines for writing an answer.

Turn over ►



- 4 A spring has modulus of elasticity $mg\text{N}$ and natural length 0.2 metres. A particle of mass $m\text{ kg}$ is attached to one end of the spring. The other end of the spring is attached to a peg which moves up and down between two points, A and B . The midpoint of AB is O . The point A is 0.1 metres above O , and B is 0.1 metres below O . At time t seconds, the displacement of the peg from the point O is $0.1 \sin(4t)$ metres, where the downward direction is taken as positive. The displacement of the particle from O at time t is x metres, as shown in the diagram.



Assume that there is no air resistance.

- (a) Show that

$$\frac{d^2x}{dt^2} + 49x = 19.6 + 4.9 \sin(4t)$$

[5 marks]

- (b) At time $t = 0$, the particle is at rest with $x = 0.4$. Find an expression for x at time t .

[10 marks]

QUESTION
PART
REFERENCE

Answer space for question 4



QUESTION
PART
REFERENCE

Answer space for question 4

A large rectangular area with horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE

Answer space for question 4

A large rectangular area with horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE

Answer space for question 4

A large rectangular area with horizontal dotted lines for writing an answer.

Turn over ►



QUESTION
PART
REFERENCE

Answer space for question 5

A large rectangular area with horizontal dotted lines for writing an answer.

Turn over ►



QUESTION
PART
REFERENCE

Answer space for question 5

A large rectangular area containing horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE


Answer space for question 5

A large rectangular area with horizontal dotted lines for writing an answer.

Turn over ►



- 6** A rod, AB , of mass m and length a is pivoted at the end A and is free to rotate in a vertical plane. A light elastic string has natural length a and modulus of elasticity $4mg$. One end of the string is attached to B and the other end to the point C , which is a distance of a directly above A . The angle between the rod and the upward vertical is θ , as shown in the diagram.



- (a) The gravitational potential energy of the system, made up of the string and the rod, is taken to be zero at the level of A . Show that V , the total potential energy of the system, is given by

$$V = \frac{mga}{2} \left(12 - 7 \cos \theta - 16 \sin \left(\frac{\theta}{2} \right) \right) \quad \text{where } \frac{\pi}{3} \leq \theta \leq \frac{5\pi}{3}$$

[5 marks]

- (b) Explain why the condition $\frac{\pi}{3} \leq \theta \leq \frac{5\pi}{3}$ has to be applied to the expression for V .

[1 mark]

- (c) Find the values of θ for which the rod is in equilibrium.

[6 marks]

- (d) Determine whether each of the values of θ found in part (c) corresponds to a position of stable or unstable equilibrium.

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 6



QUESTION
PART
REFERENCE

Answer space for question 6

A large rectangular area containing horizontal dotted lines for writing an answer.



QUESTION
PART
REFERENCE

Answer space for question 6

A large rectangular area containing horizontal dotted lines for writing an answer.

END OF QUESTIONS



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

