
A-level Use of Mathematics Pilot Mathematics

USE1 – Algebra
Mark scheme

9360
June 2015

Version/Stage: Version 1.0: Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from aqa.org.uk

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
–x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Mark	Total	Comment
1(a)	$V = \frac{300}{P}$	B1	1	cao; just $k = 300$ is not sufficient
(b)	10, 7.5, 4, 3, 2	B1	1	cao
(c)	3 points plotted correctly. All points plotted correctly (ft) and joined by a smooth curve.	B1 B1	2	Ft from their (b) ± 1 square No straight lines, no double lines No more than 1 square from any point
(d)(i)	For drawing a tangent at $P = 45$ -0.15	M1 A1	2	For their curve, not at 50 Allow [-0.2,-0.1] if a tangent is drawn at 45. -0.148 or -0.15 with no working M1A1
(ii)	Cubic metres per pascal or m^3/Pa or m^3Pa^{-1}	B1	1	
(iii)	The rate of decrease [of volume] [is 0.15 m^3 per pascal]	E1	1	Not “every” Pascal. The rate of change/increase [of volume] is [their negative value]-0.15 is ok Positive rate is ok if consistent with their graph
Total			8	
2(a)	(£)0 (£)24	B1 B1	2	
(b)	For $P = 0$, the price is zero so she doesn't take any money. For $P =$ (their) 24, she doesn't sell any (or the price is too high).	E1 E1	2	Accept any sensible answer (it is revenue that is zero, not profit). Accept any sensible answer.
(c)	£12	B1	1	cao
(d)	$120 \times 12 - 5 \times 12^2$ £720	M1 A1ft	2	ft from (c) if in the range [0,24]
(e)	$(595 = 120P - 5P^2)$ $5P^2 - 120P + 595 = 0$ $P^2 - 24P + 119 = 0$ $P = \frac{-(-24) \pm \sqrt{24^2 - 4 \times 1 \times 119}}{2 \times 1}$ $= \frac{24 \pm 10}{2}$ $= 17 \text{ or } 7$	B1 M1 A1A1 cao	4	cao (B1 for any correct equation with zero on one side) Or M1 for $(P-17)(P-7)$ SC1 per correct answer if no equations of the required form are

				seen.
	Total		11	

Q	Solution	Mark	Total	Comment
3(a)	$\ln r = \ln a^w = w \ln a$	B1	1	Evidence of ln law used
(b)	3.00, 3.69, 4.09, 4.61	B1	1	All correct; condone 3 or 3.0
(c)	3 points correct	B1		ft
	All points correct	B1		ft
	Line of best fit	B1	3	Within 2 squares of the origin
(d)	$[\ln a] = \frac{\Delta y}{\Delta x}$	B1		Or for answer in range
	$= 2.48$			Allow [2.38, 2.58]
	$a = e^{2.48}$	M1		ft
	$= 12$			
(e)	$300 = 12^w$ or $\ln 300 = w \ln 12$	A1	3	cao [10.8, 13.2]
	$w = \frac{\ln 300}{\ln 12}$	B1		Correct answer with line not used SC1
	$= 2.3 \text{ mm}$	M1		ft from (d) ; PI
(f)		A1	3	Or $\log_{12} 300$
				ft from (d)
				(must be between 2 and 2.8)
(f)	Because it is outside the range of the original data/ extrapolated.	E1	1	Or something which implies this.
Total			12	
4(a)(i)	1 (m) 23 (m)	B1 B1	2	
(ii)	360/9	M1		
	=40 (seconds)	A1	2	
(iii)	$7 = 12 + 11 \sin[9(t + 7)]^\circ$			
	$11 \sin[9(t + 7)]^\circ = -5$			
	$\sin[9(t + 7)]^\circ = -5/11$	M1		
	$9(t + 7) = 180 + 27.0 = 207$			
	$t = 16 \text{ (seconds)}$	A1	2	Awrt 16
(b)	$h = 10 + 8.5 \sin(14.4t)^\circ$	B1		B1 for each constant used in the correct place.
		B1		-1 for any extras or h or $=$ or \sin missing
		B1	3	(deduct a max of 1 mark)
Total			9	