

# GCE 2004

## *June Series*



# Mark Scheme

## Mathematics A

### *Unit MAP6*

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from:

Publications Department, Aldon House, 39, Heald Grove, Rusholme, Manchester, M14 4NA  
Tel: 0161 953 1170

or

download from the AQA website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2004 AQA and its licensors

#### COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

*Dr Michael Cresswell Director General*

**Key to Mark Scheme**

**M**.....mark is for ..... method  
**m**.....mark is dependent on one or more M marks and is for..... method  
**A**.....mark is dependent on M or m marks and is for ..... accuracy  
**B**.....mark is independent of M or m marks and is for .....method and accuracy  
**E**.....mark is for ..... explanation  
**✓ or ft or F**..... follow through from previous incorrect result  
**CAO**..... correct answer only  
**AWFW** .....anything which falls within  
**AWRT** .....anything which rounds to  
**AG** ..... answer given  
**SC** ..... special case  
**OE**..... or equivalent  
**A2,1**..... 2 or 1 (or 0) accuracy marks  
**-x EE**..... deduct x marks for each error  
**NMS**..... no method shown  
**PI** ..... possibly implied  
**SCA** .....substantially correct approach  
**c**..... candidate  
**SF**..... significant figure(s)  
**DP** ..... decimal place(s)

**Abbreviations used in Marking**

**MC – x**..... deducted x marks for mis-copy  
**MR – x**..... deducted x marks for mis-read  
**ISW**..... ignored subsequent working  
**BOD**..... given benefit of doubt  
**WR**..... work replaced by candidate  
**FB** ..... formulae booklet

**Application of Mark Scheme**

**No method shown:**

Correct answer without working..... mark as in scheme  
 Incorrect answer without working ..... zero marks unless specified otherwise

**More than one method/choice of solution:**

2 or more complete attempts, neither/none crossed out mark both/all fully and award the mean mark rounded down  
 1 complete and 1 partial attempt, neither crossed out award credit for the complete solution only

**Crossed out work**

do not mark unless it has not been replaced

**Alternative solution** using a correct or partially correct method

award method and accuracy marks as appropriate

**MAP6**

Q	Solution	Marks	Total	Comments
1(a)	$\frac{1-4}{3} = \frac{-3+4}{-1} = \frac{2-4}{2} = -1$ $\frac{1-5}{2} = \frac{-3+1}{1} = \frac{2-6}{2} = -2$	B1	1	all three must be seen
(b)	$\begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} \times \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$ $= \begin{bmatrix} -4 \\ -2 \\ 5 \end{bmatrix}$ $\begin{bmatrix} x \\ y \\ z \end{bmatrix} \cdot \begin{bmatrix} -4 \\ -2 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -4 \\ -2 \\ 5 \end{bmatrix}$ <p>Equation of plane is  <math>4x + 2y - 5z + 12 = 0</math></p>	M1A1  A1F  M1A1F	6	<p>(b) <b>Alternative:-</b></p> $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix} + \lambda \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} + \mu \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix} \quad \text{M1}$ $x = 1 + 3\lambda + 2\mu$ <p>ft miscopy <math>y = -3 - \lambda + \mu</math> A1</p> $z = 2 + 2\lambda + 2\mu$ <p>eliminate <math>\lambda</math> M1A1F</p> <p>eliminate <math>\mu</math> A1F</p> <p>result A1F</p>
(c)	<p>Perpendicular distance from (0,0,0)</p> $= \frac{12}{\sqrt{4^2 + 2^2 + (-5)^2}}$ $= \frac{4}{5} \sqrt{5}$	M1A1F  A1	3	<p>(c) <b>Alternative</b></p> $\overrightarrow{OP} = -\frac{4}{15} \begin{bmatrix} 4 \\ 2 \\ -5 \end{bmatrix} \quad \text{M1A1F}$ <p>cao <math>= \frac{4\sqrt{5}}{5}</math> A1 cao</p>
<b>Total</b>			<b>10</b>	

**MAP6 (Cont)**

Q	Solution	Marks	Total	Comments
2(a)	y -axis	B1	1	
(b)	$\sin \theta = \frac{1}{3}, \quad \cos \theta = \frac{-2\sqrt{2}}{3}$  angle is $\pi - \sin^{-1} \frac{1}{3} = 2.8$	B1B1  B1	3	Correct answer with $\tan \theta = -\frac{1}{2\sqrt{2}}$ scores 3 marks  B0 here if B0 awarded in line above cao from correct $\cos \theta$ and $\sin \theta$  2.8 with no method B1  3.5 as an answer could be correct but needs scrutiny
<b>Total</b>			<b>4</b>	
3(a)	$\Delta = 2(0 - 2) - a(0 + 6) - a(-1 - 9)$  $= 4a - 4$	M1A1  A1F	3	M1 for correct method of expansion  ft on one error
(b)	a = 1	B1F	1	
(c)(i)	$x = t, \quad y = 3t$  $z = 5t$	M1A1  A1F	3	M1 for complete method  If answer given as $x = \frac{1}{3}y = \frac{1}{5}z$ o.e. deduct 1 mark  <b>Alternative</b>  $\begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \text{ B1 } \lambda \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \text{ M1A1F}$
(ii)	sheaf (oe) of planes	E1	1	
<b>Total</b>			<b>8</b>	

MAP6 (Cont)

Q	Solution	Marks	Total	Comments
4(a)	$\vec{AB} = \begin{bmatrix} 1 \\ 2 \\ -1-p \end{bmatrix} \quad \vec{AC} = \begin{bmatrix} 2 \\ -1 \\ 2-p \end{bmatrix}$ $\vec{AD} = \begin{bmatrix} -1 \\ -3 \\ 4-p \end{bmatrix}$	B2, 1, 0	2	
(b)	$\vec{AB} \times \vec{AC} = \begin{bmatrix} 2(2-p) + (-1-p) \\ -(2-p) + 2(-1-p) \\ -5 \end{bmatrix}$ $= \begin{bmatrix} 3-3p \\ -4-p \\ -5 \end{bmatrix}$ $(\vec{AB} \times \vec{AC}) \cdot \vec{AD} = \begin{bmatrix} 3-3p \\ -4-p \\ -5 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ -3 \\ 4-p \end{bmatrix}$ $= -11 + 11p$	M1A1F  A1F		Alternative $\begin{vmatrix} -1 & -3 & 4-p \\ 1 & 2 & -1-p \\ 2 & -1 & 2-p \end{vmatrix}$  expanded correctly gather terms $11p - 11$ M1 A2, 1, 0 m1 A1F
(c)	$ -11 + 11p  = 22$ $p = 3$ $p = -1$	M1A1F  M1A1F	5  4	Incorrect formula M0 here  but allow this M1 even if formula is incorrect, and A1F also
<b>Total</b>			<b>11</b>	

## MAP6 (Cont)

Q	Solution	Marks	Total	Comments
5(a)	$\mathbf{AX} = \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} p & q \\ r & s \end{bmatrix} = \begin{bmatrix} 3p+2r & 3q+2s \\ 4p+r & 4q+s \end{bmatrix}$	M1A1		M1 for method of multiplying matrices
	$\mathbf{XB} = \begin{bmatrix} p & q \\ r & s \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 5p-q \\ 5r-s \end{bmatrix}$	B1	3	
(b)(i)	$\mathbf{AX} = \mathbf{XB}$ $3p+2r=5p$ , $4p+r=5r$ $3q+2s=-q$ , $4q+s=-s$	M1A1F		2 equations are sufficient
	$p=r$ , $-2q=s$	A1		cao
	$\mathbf{X} = \begin{bmatrix} p & q \\ p & -2q \end{bmatrix}$	A1F	4	
(ii)	Det $\mathbf{X} = -3pq \neq 0$	B1F		Any valid unsimplified expression $\neq 0$
	$\mathbf{X}^{-1} = -\frac{1}{3pq} \begin{bmatrix} -2q & -q \\ -p & p \end{bmatrix}$	M1		For method of finding inverse
		m1		Appropriate use of determinant
		A1F	4	
(iii)	$\mathbf{X}^{-1} \mathbf{AX} = \mathbf{X}^{-1} \mathbf{XB} = \mathbf{IB} = \mathbf{B}$	M1A1	2	or directly (i.e. from original matrices) $\mathbf{X}^{-1} \mathbf{X} = \mathbf{I}$ must be seen
(iv)	Eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$	B1B1		OE deduct B1 once if eigenvectors and eigenvalues are not clearly corresponding
	Eigenvalues 5, -1	B1B1	4	
<b>Total</b>			<b>17</b>	

**MAP6 (Cont)**

Q	Solution	Marks	Total	Comments
6(a)	Any method	B1	1	Must be convincing
(b)(i)	$\overrightarrow{OM} = \frac{1}{2}(\mathbf{a} + 5\mathbf{b})$	M1A1		M1 method for either
	$\overrightarrow{ON} = \frac{1}{2}(3\mathbf{a} + 3\mathbf{b})$	A1	3	
(ii)	$\Delta OMN = \frac{1}{2} \overrightarrow{OM} \times \overrightarrow{ON} $			
	$= \frac{1}{8} (\mathbf{a} + 5\mathbf{b}) \times (3\mathbf{a} + 3\mathbf{b}) $	M1		M0 if modules sign missing
	Use of $\mathbf{a} \times \mathbf{a} = 0$	B1		
	Use of $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$	B1		
	$\Delta OMN = 1.5 \mathbf{a} \times \mathbf{b} $	A1F		Must score both B1 s for this A1
	$\Delta OQR = \frac{1}{2} 3\mathbf{a} \times 5\mathbf{b} $	B1		
	$\Delta OQR = 5 \Delta OMN$	A1	6	CAO
	<b>Total</b>		<b>10</b>	
	<b>Total</b>		<b>60</b>	