

General Certificate of Education  
June 2004  
Advanced Level Examination



**MATHEMATICS AND STATISTICS  
(SPECIFICATION B)  
Unit Pure 4**

**MBP4**

Monday 21 June 2004 Morning Session

**In addition to this paper you will require:**

- an 8-page answer book;
- the AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 15 minutes

**Instructions**

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MBP4.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.

**Information**

- The maximum mark for this paper is 60.
- Mark allocations are shown in brackets.

**Advice**

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

---

Answer **all** questions.

---

1 Differentiate each of the following with respect to  $x$ :

(a)  $(1 + x^2)^8$ ; *(2 marks)*

(b)  $\frac{5x}{x^3 + 2}$ . *(2 marks)*

2 The amount of money,  $\pounds P$ , in a special savings account at time  $t$  years after 1st January 2000 is given by

$$P = 100 \times 1.05^t$$

(a) State the amount of money in the account on 1st January 2000. *(1 mark)*

(b) Calculate, to the nearest penny, the amount of money in the account on 1st January 2004. *(1 mark)*

(c) Find the value of  $t$  when  $P = 150$ , giving your answer to 3 significant figures. *(3 marks)*

3 (a) Express  $\frac{13 - 2x}{(x + 4)(2x + 1)}$  in partial fractions. *(3 marks)*

(b) Hence, prove that

$$\int_0^4 \frac{13 - 2x}{(x + 4)(2x + 1)} dx = p \ln 3 - q \ln 2$$

where  $p$  and  $q$  are positive integers. *(4 marks)*

4 The polynomial  $p(x)$  is given by

$$p(x) = x^3 - 6x^2 + 12x - 11$$

- (a) Find the remainder when  $p(x)$  is divided by  $(x - 3)$ . *(2 marks)*
- (b) The equation  $p(x) = 0$  has a single real root  $\alpha$ .
- (i) Show that  $\alpha$  lies between 3 and 4. *(1 mark)*
- (ii) Use the bisection method to find an interval of width 0.25 in which  $\alpha$  lies. *(3 marks)*
- (c) (i) Find the binomial expansion of  $(x - 2)^3$ . *(2 marks)*
- (ii) Show that  $p(x) = (x - 2)^3 - k$ , stating the value of the constant  $k$ . *(1 mark)*
- (iii) Hence find the exact solution of the equation  $p(x) = 0$ , leaving your answer in surd form. *(2 marks)*

5 A circle with centre  $C(2, -5)$  has equation  $x^2 + y^2 + ax + 10y = 7$ , where  $a$  is a constant.

- (a) (i) Find the value of  $a$ . *(2 marks)*
- (ii) Show that the radius of the circle is 6 units. *(2 marks)*
- (b) The line  $l_1$  has equation  $24x + 7y = 5k + 3$ , where  $k$  is a constant.
- (i) Prove that the distance from  $C$  to  $l_1$  is  $\frac{|2 - k|}{5}$ . *(2 marks)*
- (ii) The line  $l_1$  intersects the circle. Show that

$$|2 - k| \leq 30$$

Hence find the range of values satisfied by  $k$ . *(4 marks)*

- 6 (a) Prove the identity

$$(3 \sin x + 5 \cos x)^2 \equiv 17 + 8 \cos 2x + 15 \sin 2x \quad (4 \text{ marks})$$

- (b) Hence find:

(i)  $\int (3 \sin x + 5 \cos x)^2 dx$ ; (3 marks)

- (ii) the volume of the solid formed when the region bounded by the curve with equation  $y = 3 \sin x + 5 \cos x$ , the coordinate axes and the line  $x = \frac{\pi}{4}$  is rotated through  $2\pi$  radians about the  $x$ -axis. (3 marks)

- (c) (i) Show that the equation

$$(3 \sin x + 5 \cos x)^2 = 4 \cos^2 x$$

can be written in the form

$$(3 \tan x + 5)^2 = 4 \quad (1 \text{ marks})$$

- (ii) Hence, or otherwise, solve the equation

$$(3 \sin x + 5 \cos x)^2 = 4 \cos^2 x$$

giving all solutions in radians in the interval  $0 < x < \pi$ . (5 marks)

- 7 The function  $f$  is defined for  $0 < x < \frac{\pi}{3}$  by  $f(x) = 5 \operatorname{cosec} 3x$ .

(a) Find  $f\left(\frac{\pi}{4}\right)$  in the form  $p\sqrt{2}$ . (2 marks)

(b) (i) Find the derivative  $f'(x)$ . (2 marks)

(ii) Hence find  $f'\left(\frac{\pi}{4}\right)$  in the form  $q\sqrt{2}$ . (2 marks)

- (c) Find the equation of the tangent to the curve with equation  $y = f(x)$  at the point where  $x = \frac{\pi}{4}$ . (1 mark)

**END OF QUESTIONS**