



## General Certificate of Education

# Mathematics and Statistics 6320

## *Specification B*

*MBP3 Pure 3*

# Mark Scheme

*2005 examination - June series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

## Key to Mark Scheme

<b>M</b>	mark is for	method
<b>m</b>	mark is dependent on one or more M marks and is for	method
<b>A</b>	mark is dependent on M or m marks and is for	accuracy
<b>B</b>	mark is independent of M or m marks and is for	accuracy
<b>E</b>	mark is for	explanation
<b>√ or ft or F</b>		follow through from previous incorrect result
<b>cao</b>		correct answer only
<b>cso</b>		correct solution only
<b>awfw</b>		anything which falls within
<b>awrt</b>		anything which rounds to
<b>acf</b>		any correct form
<b>ag</b>		answer given
<b>sc</b>		special case
<b>oe</b>		or equivalent
<b>sf</b>		significant figure(s)
<b>dp</b>		decimal place(s)
<b>A2,1</b>		2 or 1 (or 0) accuracy marks
<b>-x ee</b>		deduct x marks for each error
<b>pi</b>		possibly implied
<b>sca</b>		substantially correct approach

## Abbreviations used in Marking

<b>MC – x</b>	deducted x marks for mis-copy
<b>MR – x</b>	deducted x marks for mis-read
<b>isw</b>	ignored subsequent working
<b>bod</b>	given benefit of doubt
<b>wr</b>	work replaced by candidate
<b>fb</b>	formulae book

## Application of Mark Scheme

### **No method shown:**

Correct answer without working

mark as in scheme

Incorrect answer without working

zero marks unless specified otherwise

### **More than one method / choice of solution:**

2 or more complete attempts, neither/none crossed out

mark both/all fully and award the mean mark rounded down

1 complete and 1 partial attempt, neither crossed out

award credit for the complete solution only

### **Crossed out work**

do not mark unless it has not been replaced

**Alternative solution** using a correct or partially correct method

award method and accuracy marks as appropriate


**Mathematics and Statistics B Pure 3 MBP3 June 2005**

Q	Solution	Marks	Total	Comments
1(a)(i)	$\left(\frac{3}{4}, 0\right)$ and $\left(0, -\frac{3}{2}\right)$	B1 B1	2	
(ii)	Asymptote at $x=2$ and at $y=-4$	B1 B1	2	
(iii)		M1 A1	2	One branch of hyperbola roughly correct Good graph
(b)	$4x-3=2-x \Rightarrow x=1$ Using graph, $x < 1$ Also $x > 2$	M1 B1 B1	3	M0 for $(4x-3) < (2-x)$ or $(4x-3)(2-x) < (2-x)^2$ M1 $5(2-x)(x-1) < 0$ or quotient A1 Hence $x < 1, x > 2$ A1
<b>Total</b>			<b>9</b>	
2(a)(i)	$\alpha + \beta = 3$ ; $\alpha\beta = 5$	B1 B1	2	
(ii)	$\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$ $= 9 - 10 = -1$ Since $\alpha^2 + \beta^2 < 0$ , $\alpha, \beta$ not both real	M1 A1✓ E1✓	3	
(iii)	$\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ $= 27 - 45 = -18$	M1 A1	2	Or $\alpha^3 + \beta^3 = (\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta)$ etc ag
(b)	$\sum \text{roots} = \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} = \frac{\alpha^3 + \beta^3}{\alpha\beta}$ $= -\frac{18}{5}$ $x^2 - \left(-\frac{18}{5}\right)x + 5 (= 0)$ $5x^2 + 18x + 25 = 0$	M1 A1 M1 A1	4	Formation of quadratic using sum of roots and attempt at product of roots $= \alpha\beta = 5$ cso (integer coefficients and $= 0$ )
<b>Total</b>			<b>11</b>	


**MBP3 (cont)**

Q	Solution	Marks	Total	Comments
3(a)	$\det \mathbf{M} = 5k - 14$	B1	1	
(b)	$ \det \mathbf{M}  = 1$	M1		Condone $\det \mathbf{M} = 1$
	$\Rightarrow k = 3$	A1		
	or $k = 2.6$	A1	3	
(c)(i)	$\mathbf{M}^{-1} = \frac{1}{6} \begin{bmatrix} 5 & 7 \\ 2 & 4 \end{bmatrix}$	M1		Condone a "pair" of slips in matrix, or multiplication by/ omission of $\det \mathbf{M}$ for M1
		A1	2	
(ii)	$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 5 & 7 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \end{bmatrix}$	M1		Must premultiply by $\mathbf{M}^{-1}$ ft their inverse
	$x = 9,$	A1		
	$y = 5$	A1	3	Coords are (9,5)
<b>Total</b>			<b>9</b>	
4(a)	$\frac{4 \pm \sqrt{16 - 52}}{2}$	M1		Use of quadratic equation formula or completing square
	$\sqrt{-36} = 6i$ or $\sqrt{-9} = 3i$	B1		
	$\Rightarrow x = 2 \pm 3i$	A1	3	
(b)	$(p + 3i)^2 = p^2 + 6pi - 9$	B1		
	Comparing real/imag parts	M1		$6p = 12$ or $p^2 - 9 = q$
	$p = 2; \quad q = -5$	A1	3	
<b>Total</b>			<b>6</b>	
5	When $n = 1$ ; $\text{LHS} = \frac{1}{2}$ ; $\text{RHS} = 2 - \frac{3}{2} = \frac{1}{2}$ ;	B1		(True when $n = 1$ )
	Assume formula true for $n = k$	E1		Plus the conclusion; hence true ...
	Add $(k + 1)$ th term to both sides			
	namely $\frac{k + 1}{2^{k+1}}$	B1		
	$\text{RHS} = 2 - \frac{k + 2}{2^k} + \frac{k + 1}{2^{k+1}} = 2 - \frac{***}{2^{k+1}}$	M1		2 – attempt at common denominator
	$= 2 - \frac{\{2k + 4 - k - 1\}}{2^{k+1}} = 2 - \frac{k + 3}{2^{k+1}}$	A1	5	
	Result true for $n = k + 1$			
	Hence true for $n = 1, 2, 3$ etc by induction			Must have conclusion to earn E1 mark above
<b>Total</b>			<b>5</b>	

**MBP3 (cont)**

Q	Solution	Marks	Total	Comments
6(a)(i)	Maximum value of $r = 5$ when $\theta = \pi$	B1 B1	4	condone angles mod $2\pi$ 
	Minimum value of $r = 1$ when $\theta = 0$	B1 B1		
(ii)	Symmetry about $\theta = 0$ Correct graph – approx 5: 1 ratio	B1 B1	2	
(b)(i)	$8c^2 + 2c - 3 = 0$  $\Rightarrow (2c - 1)(4c + 3) = 0$	M1	2	Attempt to factorise or solve quad eqn
	$\cos \theta = \frac{1}{2}, \quad \cos \theta = -\frac{3}{4}$	A1		
(ii)	Use of $r = 3 - 2 \cos \theta$ to find $r$	M1	4	or using $r = 8 \cos^2 \theta$ one pair of matching $r$ and $\theta$ ft second pair of matching $r$ and $\theta$ ft
	$\left[2, \frac{\pi}{3}\right], \left[2, -\frac{\pi}{3}\right], \left[\frac{9}{2}, \cos^{-1}(-0.75)\right]$	A1✓ A1✓		
	$\left[\frac{9}{2}, -\cos^{-1}(-0.75)\right]$	A1		
<b>Total</b>			<b>12</b>	
7(a)(i)	£15 000	B1	1	
(ii)	£5 000	B1	1	
(b)(i)	$\frac{dV}{dt} = -2500t^{\frac{1}{2}}$  When $t = 4$ ; $\frac{dV}{dt} = -1250$	M1 A1	3	cso
		A1		
(ii)	Car is depreciating (at this instant in time) at a rate of £1 250 per year	E1✓ E1✓	2	ft increasing in value if $> 0$ in (b) (i)
(c)(i)	$\log V = \log a + \log b^{-t}$ $= \log a - t \log b$	M1 A1	2	One rule of logs used properly
(ii)	$\log 11\,500 = \log a - \log b$ and		4	or $11\,500 = a/b$ & $5\,000 = a/b^4$ or $b^3 = 2.3$  Condone 15 180
	$\log 5\,000 = \log a - 4 \log b$	B1✓		
	$3 \log b = \log (115/50)$ or $3 \log a = \log **$	M1		
	$b = 1.32$ $a = 15\,200$	A1 A1		
<b>Total</b>			<b>13</b>	

**MBP3 (cont)**

Q	Solution	Marks	Total	Comments
8(a)	$p \otimes e = p \Rightarrow p + e + pe = p \Rightarrow e = 0$	B1	1	or $p \otimes 0 = p + 0 + 0 = p$ etc
(b)	$p \otimes 3 = p + 3 + 3p$ $p \otimes 3 = 0$ $4p = -3$ ; Hence $p = -\frac{3}{4}$	M1 M1 A1	3	or $3 \otimes q = 3 + q + 3q$ or $3 \otimes q = 0$
(c)(i)	$(p \otimes q) \otimes r$ $= (p + q + pq) + r + (p + q + pq)r$ $= p + q + r + pq + qr + rp + pqr$	M1 A1	2	full marks at this stage if correct
(ii)	$p \otimes (q \otimes r)$ considered $= p + (q + r + qr) + p(q + r + qr)$ Shown to equal $(p \otimes q) \otimes r$ yes, it is associative	M1 A1	2	
<b>Total</b>			<b>8</b>	
9(a)		M1 B1 A1	3	Hyperbola one branch correct or two half branches correct  (2,0) and (-2,0) marked or stated  good symmetrical hyperbola
(b)	Translation through $\begin{bmatrix} -1 \\ 0 \end{bmatrix}$ Stretch in y- direction Scale factor 2	M1 A1 M1 A1	4	Move left by one etc scores M1, A0
<b>Total</b>			<b>7</b>	
<b>TOTAL</b>			<b>80</b>	