

General Certificate of Education
June 2005
Advanced Level Examination



**MATHEMATICS AND STATISTICS
(SPECIFICATION B)
Unit Pure 6**

MBP6

Tuesday 28 June 2005 Afternoon Session

In addition to this paper you will require:

- a 12-page answer book;
- the AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 45 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MBP6.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.

Information

- The maximum mark for this paper is 80.
- Mark allocations are shown in brackets.

Advice

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

Answer **all** questions.

- 1 Show that the radius of curvature, at the point where $x = a$, of the curve with equation $y = \cosh x$ is $\cosh^2 a$. (4 marks)

- 2 By writing $\sinh y = x$, prove that

$$\sinh^{-1} x = \ln \left\{ x + \sqrt{x^2 + 1} \right\} \quad (4 \text{ marks})$$

- 3 (a) Solve the equation $7 - \cosh x = \sinh x$, giving your answer in the form $\ln k$, where k is an integer. (3 marks)
- (b) On the same diagram, sketch the graphs of $y = \sinh x$ and $y = 7 - \cosh x$, indicating the coordinates of the points where the curves cross the axes. (4 marks)

- 4 A curve is defined parametrically by

$$x = t^2 + \frac{2}{3}t^3, \quad y = t + t^2$$

- (a) Verify that $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (1 + 2t + 2t^2)^2$. (3 marks)
- (b) (i) Determine the length of the arc of this curve between the points where $t = 0$ and $t = 1$. (3 marks)
- (ii) When this arc is rotated about the x -axis through 2π radians, it generates a surface of revolution with area S . Find S in terms of π . (4 marks)

5 (a) (i) Show that $\frac{1}{\cos \theta + i \sin \theta} = \cos \theta - i \sin \theta$. (1 mark)

(ii) Given that $z = \cos \frac{\pi}{4} + i \sin \frac{\pi}{4}$, evaluate

$$\left(z + \frac{1}{z}\right)^{12} \quad (4 \text{ marks})$$

(b) (i) Use de Moivre's Theorem to simplify the expression

$$[r(\cos \theta + i \sin \theta)]^{-n} \quad (1 \text{ mark})$$

(ii) Using this result, or otherwise, find the integer m for which

$$\left(\frac{1}{2} - \frac{1}{2}i\right)^{-12} = m \quad (5 \text{ marks})$$

6 All points (x, y) on a particular curve satisfy the differential equation

$$\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 4y = 4 + 27e^{-x}$$

The point $P(0, 6)$ lies on this curve, and the gradient of the curve at P is zero.

(a) Without solving the differential equation, evaluate $\frac{d^2y}{dx^2}$ at P and hence describe P as fully as possible. (3 marks)

(b) By solving the differential equation, find the equation of this curve. (11 marks)

7 (a) Use the identity $\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$ to prove that:

(i) $\sin 2\theta = 2 \sin \theta \cos \theta$; (1 mark)

(ii) $\sin 3\theta + \sin \theta = 4 \sin \theta - 4 \sin^3 \theta$. (4 marks)

(b) Show that $(2s - 1)$ is a factor of $8s^3 - 8s + 3$. (1 mark)

(c) Solve the equation

$$\sin 3\theta + \sin \theta = \frac{3}{2}$$

giving your answers in radians in the interval $0 < \theta < \pi$. (9 marks)

Turn over ►

- 8 (a) (i) Write down the matrix \mathbf{R} which represents an anticlockwise rotation through $\frac{\pi}{4}$ radians about O . (1 mark)
- (ii) Describe the transformation with matrix $\mathbf{S} = \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$, where k is a positive constant. (2 marks)
- (iii) Show that $\mathbf{R}^{-1}\mathbf{S}\mathbf{R} = \frac{1}{2} \begin{bmatrix} k+2 & k \\ -k & 2-k \end{bmatrix}$. (3 marks)
- (b) Show that the matrix $\mathbf{M} = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix}$ has just one eigenvalue λ . State the value of λ and find a corresponding eigenvector. (4 marks)
- (c) The transformation \mathbf{T} is defined by $\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{M} \begin{bmatrix} x \\ y \end{bmatrix}$.
- (i) Write down a cartesian equation of the line of invariant points of \mathbf{T} , and give a reason for your choice of answer. (2 marks)
- (ii) Show that, for some value of k , $\mathbf{M} = \mathbf{R}^{-1}\mathbf{S}\mathbf{R}$ and hence give a full geometric description of \mathbf{T} . (3 marks)

END OF QUESTIONS