



## General Certificate of Education

# Mathematics and Statistics 6320

## *Specification B*

### *MBS5 Statistics 5*

# Mark Scheme

## *2005 examination - June series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

## Key to Mark Scheme

<b>M</b>	mark is for	method
<b>m</b>	mark is dependent on one or more M marks and is for	method
<b>A</b>	mark is dependent on M or m marks and is for	accuracy
<b>B</b>	mark is independent of M or m marks and is for	accuracy
<b>E</b>	mark is for	explanation
<b>√ or ft or F</b>		follow through from previous incorrect result
<b>cao</b>		correct answer only
<b>cso</b>		correct solution only
<b>awfw</b>		anything which falls within
<b>awrt</b>		anything which rounds to
<b>acf</b>		any correct form
<b>ag</b>		answer given
<b>sc</b>		special case
<b>oe</b>		or equivalent
<b>sf</b>		significant figure(s)
<b>dp</b>		decimal place(s)
<b>A2,1</b>		2 or 1 (or 0) accuracy marks
<b>-x ee</b>		deduct x marks for each error
<b>pi</b>		possibly implied
<b>sca</b>		substantially correct approach

## Abbreviations used in Marking

<b>MC – x</b>		deducted x marks for mis-copy
<b>MR – x</b>		deducted x marks for mis-read
<b>isw</b>		ignored subsequent working
<b>bod</b>		given benefit of doubt
<b>wr</b>		work replaced by candidate
<b>fb</b>		formulae book

## Application of Mark Scheme

### **No method shown:**

Correct answer without working

mark as in scheme

Incorrect answer without working

zero marks unless specified otherwise

### **More than one method / choice of solution:**

2 or more complete attempts, neither/none crossed out

mark both/all fully and award the mean mark rounded down

1 complete and 1 partial attempt, neither crossed out

award credit for the complete solution only

### **Crossed out work**

do not mark unless it has not been replaced

**Alternative solution** using a correct or partially correct method


award method and accuracy marks as appropriate

**Mathematics and Statistics B Statistics 5 MBS5 June 2005**

<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>1(a)</b>	See graph on next page	M1 B1 A1	3	Scales and labels Reasonably accurate plot – allow one small slip
<b>(b)</b>	$y = -7.90 + 1.04x$  $x = 10 \quad y = 2.5 \quad x = 160 \quad y = 158.3$ + line	B2 B1 M1 A1	5	1.04 (1.035 to 1.045) allow M1A1 if method shown – 7.90(– 7.89 to – 7.91) Method for their line Accurate line - by eye
<b>(c)</b>	B $4 - (-7.90) - 1.04 \times 12 = -0.56$ G $88 - (-7.90) - 1.04 \times 86 = 6.55$	M1 m1 A1	3	Method for residuals - ignore sign Method for residuals - consistent signs must be demonstrated - eg. disallow if one residual is zero – 0.56(– 0.55 to – 0.6) and 6.55(6.45 to 6.6)
<b>(d)</b>	B has a small residual but 4 is a poor estimate of 12 G has relatively large residual but 88 is a good estimate of 86 Small residual indicates consistent with pattern of other estimates - not necessarily good or bad.	E1 E1 E1	3	Small residual not necessarily good Illustrated by B or G Small residual $\Rightarrow$ consistent with pattern or other relevant comment
<b>(e)</b>	Actual age is reduced by 7. Equation becomes $y = -0.90 + 1.04x$ - which is very close to ideal $y = x$ . Eamon's estimates better than appeared in part (b)	E1 E1	2	Estimates improved - disallow if no or clearly incorrect reason Corrected equation or ideal is $y = x$ or other sensible comment
<b>Total</b>			<b>16</b>	

**MBS5 (cont)**

**Graph for question 1**




## MBS5 (cont)

Q	Solution	Marks	Total	Comments
2(a)	$z = \frac{20-26}{8} = -0.75$	M1	5	Method for z - ignore sign
	Probability no need to refill $1 - 0.77337 = 0.227$	M1 A1		Completely correct method 0.227 (0.226 to 0.227)
	$z = \frac{40-26}{8} = 1.75$	M1		Correct method
	Probability exactly one refill i.e between 20 and 40 = $0.95994 - 0.2263 = 0.733$	A1		0.733 (0.733 to 0.734)
(b)(i)	$0.22623^5 = 0.000598$	M1 A1	2	0.000598 (0.00059 to 0.00061)
(ii)	$z = \frac{20-26}{\frac{8}{\sqrt{5}}} = -1.677$	M1	3	Use of $\frac{8}{\sqrt{5}}$
	probability mean less than 20 = $1 - 0.9532 = 0.0468$	m1 A1		Completely correct method 0.0468(0.046 to 0.048)
(c)	$\mu - 0.9945\sigma = 20$	B1	6	0.9945 (0.994 to 0.995)
	$\mu - 1.175\sigma = 40$	B1		1.175 (1.17 to 1.18)
		M1		Good attempt at equations - ignore sign
	$2.169\sigma = 20$	m1		Completely correct equations
	$\sigma = 9.222$	m1		Method of solution
	$\mu = 29.2$	A1	$\sigma = 9.222$ (9.21 to 9.23) and $\mu = 29.2$ (29.1 to 29.3)	
<b>Total</b>			<b>16</b>	

## MBS5 (cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	$197 \pm 1.96 \times \frac{103}{\sqrt{90}}$	M1	4	Use of $\frac{103}{\sqrt{90}}$
	197 ± 21.3 176 ~ 218	B1 m1 A1		1.96 Completely correct method - their $z$ 197 ± 21.3(21.25 to 21.35) or 176(175.5 to 176) and 218 (218 to 218.5)
(ii)	42.6	B1	1	42.6(42.5 to 42.6)
(iii)	$2z \times \frac{103}{\sqrt{90}} = 30$	M1	5	Reasonable attempt at equation containing $z$ - ignore omission of 2
	$z = 1.382$	m1 m1		Completely correct equation containing $z$ Method for finding $z$
(iv)	$1 - 2(1 - 0.9164) = 0.833$ 83.3%	M1 A1	4	Method for probability - their $z$ 83.3 (83 to 83.5)
	$2 \times 2.5758 \times \frac{103}{\sqrt{n}} = 30$ $n = 312.8$	B1 M1 m1 A1		2.5758 (2.57 to 2.58) Reasonable attempt at equation involving $n$ - ignore omission of 2, incorrect $z$ Method of solution of equation 313 cao
(b)(i)	313 needed large sample $\Rightarrow$ sample mean normally distributed	E1 E1	2	Large sample / CLT <b>Mean</b> normally distributed
(ii)	Mean less than 2 s.d. above zero $\Rightarrow$ non-trivial probability of negative values which are not possible	E1 E1	2	Mean less than 2 s.d. above zero / possibility of negative values / money discrete variable
<b>Total</b>			<b>18</b>	



**MBS5 (cont)**

<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>4(a)</b>	0.3	B1	1	0.3 cao
<b>(b)(i)</b>	$\frac{10}{30} = \frac{1}{3}$	B1	1	$\frac{1}{3}$ acf
<b>(ii)</b>	$\frac{1}{3} \times 0.06 = 0.02$	M1	1	Method - their (b)(i)
<b>(iii)</b>	$\frac{15}{30}(0.30 + 0.15) = 0.225$	M1 A1	2	Method - generous 0.225
<b>(iv)</b>	$\frac{1}{3} \times 0.06 + \frac{1}{2} \times 0.15 + \frac{5}{30} \times 0.18 = 0.125$	M1 m1 A1	3	Attempt at P(4* comedy) + P(4* drama) + P(4* other) Completely correct method 0.125 cao
<b>(v)</b>	$\frac{3}{4} \times (0.20 + 0.35) + \frac{1}{4} (0.40 + 0.10) = 0.5375$	M1 m1 A1	3	Reasonable attempt Completely correct method 0.5375 (0.537 to 0.538)
<b>(c)</b>	$3 \times \frac{10}{30} \times \frac{9}{29} \times \frac{15}{28} = 0.166$	B1 M1 A1	3	3 Allow omission of or incorrect '3' - allow with replacement 0.166 (0.166 to 0.1665)
<b>Total</b>			<b>14</b>	

## MBS5 (cont)

Q	Solution	Marks	Total	Comments
5(a)(i)	$H_0 : \mu = 40$	B1		One correct hypothesis – generous
	$H_1 : \mu \neq 40$ (allow $\mu > 40$ and 1.6449)	B1		Both hypotheses correct – ungenerous
	$z = \frac{46.5 - 40}{\frac{12}{\sqrt{8}}} = 1.53$	M1		Allow $H_1 \mu > 40$
		m1		Use of $\frac{12}{\sqrt{8}}$
	c.v $\pm 1.96$ ; 1.53 lies between $\pm 1.96$ so accept $H_0$ , mean is 40 mins	A1 B1		Completely correct method for $z$ ignore sign 1.53 (1.525 to 1.535) 1.96 - ignore sign (cv 1.895 for one tail test)
		A1✓		Correct conclusion – must be compared with $z$
				<b><i>N.B. apply this mark scheme to (a)(ii) and vice versa if more favourable to candidate</i></b>
(ii)	$H_0 : \mu = 50$	B1		Both hypotheses correct – ungenerous
	$H_1 : \mu \neq 50$ (allow $\mu < 50$ and $-1.6449$ )			
	$z = \frac{46.5 - 50}{\frac{12}{\sqrt{8}}} = -0.825$	A1		$-0.825$ ( $-0.8245$ to $-0.8255$ )
	c.v $\pm 1.96$ ; $-0.825$ lies between $\pm 1.96$ so accept $H_0$ , mean is 50 mins	A1✓	10	Correct conclusion must be compared with both tails or lower tail of t
(b)	Claim 1. <b>C</b> Not true - no null hypothesis rejected so no Type 1 error made	E2,1		Correct conclusion for correct reason - be generous for E1 but disallow no or clearly incorrect reason
	Claim 2. <b>B</b> Possibly true - true if population mean is equal to neither 40 nor 50	E2,1		Correct conclusion for correct reason - be generous for E1 but disallow no or clearly incorrect reason
	Claim 3. <b>A</b> Definitely true - since mean cannot equal both 40 and 50	E2,1	6	Correct conclusion for correct reason - be generous for E1 but disallow no or clearly incorrect reason
	<b>Total</b>		<b>16</b>	
	<b>TOTAL</b>		<b>80</b>	