

ADVANCED SUBSIDIARY (AS) General Certificate of Education January 2014

Mathematics

Assessment Unit M1 assessing Module M1: Mechanics 1

[AMM11]

FRIDAY 24 JANUARY, AFTERNOON

MARK SCHEME

GCE ADVANCED/ADVANCED SUBSIDIARY (AS) MATHEMATICS

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

- M indicates marks for correct method.
- W indicates marks for working.
- MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.


When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

1 Moments about C,

(clockwise is +ve)

4(3a) - 5(a) - 2(a) - 6(4a) = -19a MW2

2 (i) Forces acting on the crate



MW2

M1

MW1

M1

MW2

M1

MW1

(ii) Resolve vertically

 $R + 15\sin(30^\circ) = 5 \text{ g}$ M1

$$R = 41.5$$
 W1

Resolve horizontally

 $F_r = 15 \cos(30^\circ) = 12.99$ MW1

$$F_r = \mu R$$
 M1

$$\mu = \frac{F_r}{R} = \frac{12.99}{41.5} = 0.313$$
 MW1

(iii) Vertically, $R = 5 \text{ g} - T \sin(30^\circ)$

Horizontally,
$$F = ma$$

$$T \cos (30^\circ) - 0.313 (5 \text{ g} - T \sin (30^\circ)) = 5(2)$$


$$T = 24.8 \text{ N}$$
 W1

13

AVAILABLE MARKS

4


AVAILABLE MARKS 3 (i) u = 1*a* = 0.15 M1 s = ? $s_1 = ut + \frac{1}{2}at^2$ =(1)t + $\frac{1}{2}(0.15)t^2$ t = tDist for Kathryn is $= t + 0.075t^2$ W1 u = 2v = 2a = 0s = ? $s_2 = ut + \frac{1}{2}at^2$ t = tDist for Victoria is =2tM1 W1 AB = 90 $s_1 + s_2 = 90$ M1 $\therefore \quad (t + 0.075t^2) + (2t) = 90$ W1 $0.075t^2 + 3t - 90 = 0$ $t^2 + 40t - 1200 = 0$ (t-20)(t+60) = 0t = 20 or - 60MW1 $\therefore t = 20$ W1 **(ii)** B 90



MW2

(i)	Conservation of momentum	1	M1	AVAILABLE MARKS	
		+ Bm(u) = 2m(2u) + m(V)			
	V = -u Velocity of A = $-u$		W1		
(ii)	Valid explanation of sign cl	MW1			
(iii)		I = mv - mu	M1		
	Impulse to B	= 2m(2u) - 2m(u)			
		= 2mu Ns	W2	7	

5 (i)



6	(i)	v	=	$\int a dt$	M1

$$v = 2t^2 + 2t + c \tag{W1}$$

v = 0 at t = 0 therefore c = 0 MW1

at
$$t = 3$$
, $v = 2(3)^2 + 2(3) = 24 \text{ m s}^{-1}$ W1

(ii) Stops during 2nd 'part' of motion (uniform acceleration)

$$u = 24, a = -4$$

Stops when $v = 0$

$$v = u + at$$
 M1

$$0 = 24 - 4(T)$$

$$T = 6$$
 W1

At rest at
$$3 + 6 = 9$$
 s MW1

(iii) Distance between t = 0 and t = 3 (variable acceleration)

$$s = \int v dt$$
 M1

$$s = \frac{2}{3}t^3 + t^2 + c$$
 W1

$$s = 0$$
 at $t = 0$ therefore $c = 0$ MW1

at
$$t = 3$$
, $s = \frac{2}{3}(3)^3 + (3)^2 = 27$ MW1

Distance between t = 3 and t = 9 (constant acceleration)

$$u = 24, t = 6, a = -4$$

$$s = \frac{1}{2}(u+v)t$$

$$s = 72$$
MW1

Total distance =
$$27 + 72 = 99 \text{ m}$$
 MW1

15

AVAILABLE MARKS

M1

M1

7	(i)	T		AVAILABLE MARKS
		$F_r \leftarrow 250 \mathrm{g}$	MW2	
	(ii)	Pipe is a rod (only has length) Pipe is rigid (doesn't bend)	MW1	
	(iii)	Moments about A	M1	
		Clockwise = anticlockwise		
		$2500(15\cos 20^\circ) + T\cos 70^\circ(30\sin 20^\circ) = T\sin 70^\circ(30\cos 20^\circ)$ or $T\sin 50^\circ \times 30 = 15 \times 250g\cos 20^\circ$	W3	
		T = 1533.27		
		T = 1530(3 s.f.)	W1	
	(iv)	Resolve vertically	M1	
		$N + T\sin 70^\circ = 250 \text{ g}$	W1	
		<i>N</i> = 1059.103	W1	
		Resolve horizontally	M1	
		$F_r = T\cos 70^\circ$		
		$F_r = 524.44$	W1	
		$R = \sqrt{N^2 + F_r^2}$	M1	
		R = 1180 N	W1	15
			Total	75