

**MARK SCHEME for the October/November 2009 question paper
for the guidance of teachers**

9709/31

9709 MATHEMATICS

Paper 31, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through \surd ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

- 1 *EITHER*: State or imply non-modular inequality $(2 - 3x)^2 < (x - 3)^2$, or corresponding equation, and make a reasonable solution attempt at a 3-term quadratic M1
Obtain critical value $x = -\frac{1}{2}$ A1
Obtain $x > -\frac{1}{2}$ A1
Fully justify $x > -\frac{1}{2}$ as only answer A1
- OR1*: State the relevant critical linear equation, i.e. $2 - 3x = 3 - x$ B1
Obtain critical value $x = -\frac{1}{2}$ B1
Obtain $x > -\frac{1}{2}$ B1
Fully justify $x > -\frac{1}{2}$ as only answer B1
- OR2*: Obtain the critical value $x = -\frac{1}{2}$ by inspection, or by solving a linear inequality B2
Obtain $x > -\frac{1}{2}$ B1
Fully justify $x > -\frac{1}{2}$ as only answer B1
- OR3*: Make recognisable sketches of $y = 2 - 3x$ and $y = |x - 3|$ on a single diagram B1
Obtain critical value $x = -\frac{1}{2}$ B1
Obtain $x > -\frac{1}{2}$ B1
Fully justify $x > -\frac{1}{2}$ as only answer B1 [4]
[Condone \geq for $>$ in the third mark but not the fourth.]
- 2 *EITHER*: Use laws of indices correctly and solve a linear equation for 3^x , or for 3^{-x} M1
Obtain 3^x , or 3^{-x} in any correct form, e.g. $3^x = \frac{3^2}{(3^2 - 1)}$ A1
Use correct method for solving $3^{\pm x} = a$ for x , where $a > 0$ M1
Obtain answer $x = 0.107$ A1
- OR*: State an appropriate iterative formula, e.g. $x_{n+1} = \frac{\ln(3^{x_n} + 9)}{\ln 3} - 2$ B1
Use the formula correctly at least once M1
Obtain answer $x = 0.107$ A1
Show that the equation has no other root but 0.107 A1 [4]
[For the solution 0.107 with no relevant working, award B1 and a further B1 if 0.107 is shown to be the only root.]
- 3 (i) Use the iterative formula correctly at least once M1
State final answer 2.78 A1
Show sufficient iterations to at least 4 d.p. to justify its accuracy to 2 d.p., or show there is a sign change in an appropriate function in (2.775, 2.785) A1 [3]
- (ii) State a suitable equation, e.g. $x = \frac{3}{4}x + \frac{15}{x^3}$ B1
State that the exact value of α is $\sqrt[4]{60}$, or equivalent B1 [2]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

- 4 Use product or quotient rule M1
 Obtain derivative in any correct form A1
 Equate derivative to zero and obtain an equation of the form $a \sin 2x = b$, or a quadratic in $\tan x$, $\sin^2 x$, or $\cos^2 x$ M1*
 Carry out correct method for finding one angle M1(dep*)
 Obtain answer, e.g. 0.365 A1
 Obtain second answer 1.206 and no others in the range (allow 1.21) A1 [6]
 [Ignore answers outside the given range.]
 [Treat answers in degrees, 20.9° and 69.1° , as a misread.]
- 5 (i) EITHER: Use double angle formulae correctly to express LHS in terms of trig functions of 2θ M1
 Use trig formulae correctly to express LHS in terms of $\sin \theta$, converting at least two terms M1
 Obtain expression in any correct form in terms of $\sin \theta$ A1
 Obtain given answer correctly A1
 OR: Use double angle formulae correctly to express RHS in terms of trig functions of 2θ M1
 Use trig formulae correctly to express RHS in terms of $\cos 4\theta$ and $\cos 2\theta$ M1
 Obtain expression in any correct form in terms of $\cos 4\theta$ and $\cos 2\theta$ A1
 Obtain given answer correctly A1 [4]
- (ii) State indefinite integral $\frac{1}{4} \sin 4\theta - \frac{4}{2} \sin 2\theta + 3\theta$, or equivalent B2
 (award B1 if there is just one incorrect term)
 Use limits correctly, having attempted to use the identity M1
 Obtain answer $\frac{1}{32} (2\pi - \sqrt{3})$, or any simplified exact equivalent A1 [4]
- 6 (i) EITHER: State that the position vector of M is $2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, or equivalent B1
 Carry out a correct method for finding the position vector of N M1
 Obtain answer $3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$, or equivalent A1
 Obtain vector equation of MN in any correct form, e.g. $\mathbf{r} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k} + \lambda(\mathbf{i} - 3\mathbf{j} + 3\mathbf{k})$ A1
 OR: State that the position vector of M is $2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, or equivalent B1
 Carry out a correct method for finding a direction vector for MN M1
 Obtain answer, e.g. $\mathbf{i} - 3\mathbf{j} + 3\mathbf{k}$, or equivalent A1
 Obtain vector equation of MN in any correct form, e.g. $\mathbf{r} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k} + \lambda(\mathbf{i} - 3\mathbf{j} + 3\mathbf{k})$ A1 [4]
 [SR: The use of $AN = AC/3$ can earn M1A0, but $AN = AC/2$ gets M0A0.]
- (ii) State equation of BC in any correct form, e.g. $\mathbf{r} = 3\mathbf{i} + 2\mathbf{j} - 3\mathbf{k} + \mu(\mathbf{i} - 5\mathbf{j} + 5\mathbf{k})$ B1
 Solve for λ or for μ M1
 Obtain correct value of λ , or μ , e.g. $\lambda = 3$, or $\mu = 2$ A1
 Obtain position vector $5\mathbf{i} - 8\mathbf{j} + 7\mathbf{k}$ A1 [4]
- 7 (i) Substitute $x = -2 + i$ in the equation and attempt expansion of $(-2 + i)^3$ M1
 Use $i^2 = -1$ correctly at least once and solve for k M1
 Obtain $k = 20$ A1 [3]
- (ii) State that the other complex root is $-2 - i$ B1 [1]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

- (iii) Obtain modulus $\sqrt{5}$ B1
 Obtain argument 153.4° or 2.68 radians B1 [2]
- (iv) Show point representing u in relatively correct position in an Argand diagram B1
 Show vertical line through $z = 1$ B1
 Show the correct half-lines from u of gradient zero and 1 B1
 Shade the relevant region B1 [4]
 [SR: For parts (i) and (ii) allow the following alternative method:
 State that the other complex root is $-2 - i$ B1
 State quadratic factor $x^2 + 4x + 5$ B1
 Divide cubic by 3-term quadratic, equate remainder to zero and solve for k , or, using
 3-term quadratic, factorise cubic and obtain k M1
 Obtain $k = 20$ A1]
- 8 (i) State or imply partial fractions are of the form $\frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{3x+2}$ B1
 Use any relevant method to obtain a constant M1
 Obtain one of the values $A = 1, B = 2, C = -3$ A1
 Obtain a second value A1
 Obtain the third value A1 [5]
- (ii) Use correct method to obtain the first two terms of the expansion of $(x+1)^{-1}, (x+1)^{-2}, (3x+2)^{-1}$
 or $(1 + \frac{3}{2}x)^{-1}$ M1
 Obtain correct unsimplified expansion up to the term in x^2 of each partial
 fraction $A1\sqrt{ } + A1\sqrt{ } + A1\sqrt{ }$
 Obtain answer $\frac{3}{2} - \frac{11}{4}x + \frac{29}{8}x^2$, or equivalent A1 [5]
- [Symbolic binomial coefficients, e.g. $\binom{-1}{1}$, are not sufficient for the first M1. The f.t. is on A, B, C .]
 [The form $\frac{Dx+E}{(x+1)^2} + \frac{C}{3x+2}$, where $D = 1, E = 3, C = -3$, is acceptable. In part (i) give
 B1M1A1A1A1.
 In part (ii) give M1A1√A1√ for the expansions, and, if $DE \neq 0$, M1 for multiplying out fully and A1
 for the final answer.]
 [If B or C omitted from the form of fractions, give B0M1A0A0A0 in (i); M1A1√A1√ in (ii), max
 4/10]
 [If D or E omitted from the form of fractions, give B0M1A0A0A0 in (i); M1A1√A1√ in (ii), max
 4/10]
 [In the case of an attempt to expand $(5x+3)(x+1)^{-2}(3x+2)^{-1}$, give M1A1A1 for the expansions, M1
 for multiplying out fully, and A1 for the final answer.]
 [Allow use of Maclaurin, giving M1A1√A1√ for differentiating and obtaining $f(0) = \frac{3}{2}$ and
 $f'(0) = -\frac{11}{4}$, A1√ for $f''(0) = \frac{29}{4}$, and A1 for the final answer (the f.t. is on A, B, C if used).]
- 9 (i) State coordinates (1, 0) B1 [1]
- (ii) Use correct quotient or product rule M1
 Obtain derivative in any correct form A1
 Equate derivative to zero and solve for x M1
 Obtain $x = e^2$ correctly A1 [4]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	31

- (iii) Attempt integration by parts reaching $a\sqrt{x} \ln x \pm a \int \sqrt{x} \frac{1}{x} dx$ M1*
- Obtain $2\sqrt{x} \ln x - 2 \int \frac{1}{\sqrt{x}} dx$ A1
- Integrate and obtain $2\sqrt{x} \ln x - 4\sqrt{x}$ A1
- Use limits $x = 1$ and $x = 4$ correctly, having integrated twice M1(dep*)
- Justify the given answer A1 [5]
- 10 (i) State or imply $\frac{dA}{dt} = kV$ M1*
- Obtain equation in r and $\frac{dr}{dt}$, e.g. $8\pi r \frac{dr}{dt} = k \frac{4}{3} \pi r^3$ A1
- Use $\frac{dr}{dt} = 2, r = 5$ to evaluate k M1(dep*)
- Obtain given answer A1 [4]
- (ii) Separate variables correctly and integrate both sides M1
- Obtain terms $-\frac{1}{r}$ and $0.08t$, or equivalent A1 + A1
- Evaluate a constant or use limits $t = 0, r = 5$ with a solution containing terms of the form $\frac{a}{r}$ and bt M1
- Obtain solution $r = \frac{5}{(1-0.4t)}$, or equivalent A1 [5]
- (iii) State the set of values $0 \leq t < 2.5$, or equivalent B1 [1]
- [Allow $t < 2.5$ and $0 < t < 2.5$ to earn B1.]