UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2010 question paper for the guidance of teachers

9709 MATHEMATICS

9709/31

Paper 31, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	31

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0. B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking *g* equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	31

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
sos	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	31

1	EITHER:	quadratio	imply non-modular inequality $(x+3a)^2 > (2(x-2a))^2$, or corresponding c equation, or pair of linear equations $(x+3a) = \pm 2(x-2a)$ easonable solution attempt at a 3-term quadratic, or solve two linear	B1	
		equation		M1	
		Obtain c	ritical values $x = \frac{1}{3}a$ and $x = 7a$	A1	
		State ans	swer $\frac{1}{3}a < x < 7a$	A 1	
	OR:	solving a	he critical value $x = 7a$ from a graphical method, or by inspection, or by a linear equation or inequality he critical value $x = \frac{1}{3}a$ similarly	B1 B2	
			Swer $\frac{1}{3}a < x < 7a$	B1	[4]
			condone \leq for \leq ; accept 0.33 for $\frac{1}{3}$.]	D1	נין
2	Use corre	ect cos 2A 1	formula and obtain an equation in $\sin \theta$	M1	
	Obtain 4	$\sin^2 \theta + \sin^2 \theta$	$n\theta - 3 = 0$, or equivalent	A1	
			tempt to solve a 3-term quadratic in $\sin \theta$	M1	
		swer 48.6°		A1 A1 √	
			4° and no others in the given range and no others in the given range	AI V	[6]
			answers in radians as a misread. Ignore answers outside the given range.]	111	[~]
3	(i)	EITHER:	State or imply $n \ln x + \ln y = \ln C$	B1	
	(1)	BIIIIBI	Substitute <i>x</i> - and <i>y</i> -values and solve for <i>n</i>	M1	
			Obtain $n = 1.50$	A1	
			Solve for <i>C</i>	M1	
			Obtain $C = 6.00$	A1	
		OR:	Obtain two correct equations by substituting x- and y-values in $x^n y = C$	B1	
			Solve for n Obtain $n = 1.50$	M1	
			Solve for C	A1 M1	
			Obtain $C = 6.00$	A1	[5]
	(ii) State that the graph of $\ln y$ against $\ln x$ has equation $n \ln x + \ln y = \ln C$ whice				
			n y and ln x, or has equation of the form $nX + Y = \ln C$, where $X = \ln x$ and nd is thus a straight line	B1	[1]
4	(i)	State corre	ect expansion of $\cos(3x - x)$ or $\cos(3x + x)$	B1	
-	(-)		expansions in $\frac{1}{2}(\cos 2x - \cos 4x)$, or equivalent	M1	
			and obtain the given identity correctly	A1	[3]
	(ii)	Obtain int	egral $\frac{1}{4}\sin 2x - \frac{1}{8}\sin 4x$	B1	
			limits correctly in an integral of the form $a \sin 2x + b \sin 4x$	M1	
			ven answer following full, correct and exact working	A1	[3]

			GCE AS/A LEVEL – May/June 2010	9709	31	
5	Separate	varial	ples correctly		B1	
	_	Integrate and obtain term $\ln x$				
			$btain term \frac{1}{2} ln(y^2 + 4)$		B1	
			stant or use limits $y = 0$, $x = 1$ in a solution containing $a \ln x$	and $b\ln(y^2+4)$	M1	
	Obtain c	orrect	solution in any form, e.g. $\frac{1}{2}\ln(y^2 + 4) = \ln x + \frac{1}{2}\ln 4$		A1	
	Rearrang	ge as j	$y^2 = 4(x^2 - 1)$, or equivalent		A1	[6]
6	(i)	Using	g the formulae $\frac{1}{2}r^2\theta$ and $\frac{1}{2}r^2\sin\theta$, or equivalent, form an	equation	M1	
		Obta	in a correct equation in r and x and/or $x/2$ in any form		A1	
		Obta	in the given equation correctly		A1	[3]
	(ii)	Cons	ider the sign of $x - (\frac{3}{4}\pi - \sin x)$ at $x = 1.3$ and $x = 1.5$, or eq	uivalent	M1	
			plete the argument with correct calculations		A1	[2]
	(iii)	Use t	he iterative formula correctly at least once		M1	
	· /	Obta	in final answer 1.38		A1	
			v sufficient iterations to at least 4 d.p. to justify its accura is a sign change in the interval (1.375, 1.385)	cy to 2 d.p., or sho	A1	[3]
7	(i)	Obta	in modulus $\sqrt{8}$		B1	
		Obta	in argument $\frac{1}{4}\pi$ or 45°		B1	[2]
	(ii)	Show	v 1, i and u in relatively correct positions on an Argand diag	gram	B1	
			with perpendicular bisector of the line joining 1 and i		B1	
			<i>y</i> a circle with centre <i>u</i> and radius 1 e the correct region		B1 B1	[4]
						[.]
	(iii)		or imply relevance of the appropriate tangent from O to the out complete strategy for finding $ z $ for the critical point	e circle	B1 √ M1	
			in answer $\sqrt{7}$		A1	[3]
		Oota	in answer v		M	[5]
8	(i)	State	or imply the form $\frac{A}{x+1} + \frac{B}{x+3}$ and use a relevant method to	so find A or B	M1	
			x+1 $x+3$ in $A = 1, B = -1$		A1	[2]
	(**)	C			3.61	
	(ii)		re the result of part (i) and substitute the fractions of part (i) in the given answer correctly)	M1 A1	[2]
	(iii)		rate and obtain $-\frac{1}{x+1} - \ln(x+1) + \ln(x+3) - \frac{1}{x+3}$		В3	
			titute limits correctly in an integral containing at least two	terms of the corre		
		form Obta	in given answer following full and exact working		M1 A1	[5]
		Joia	in given answer following full and exact working		111	اریا

Mark Scheme: Teachers' version

Syllabus

Paper

Page 5

	Page 6	6	Mark Scheme: Teachers' version	Syllabus	Paper	•
			GCE AS/A LEVEL – May/June 2010	9709	31	
9	(i)	Hee and	otient or product rule to differentiate $(1-x)/(1+x)$		M1	
	(1)	_	correct derivative in any form		A1	
			· · · · · · · · · · · · · · · · · · ·			
		Use cha	ain rule to find $\frac{dy}{dx}$		M1	
			a correct expression in any form		A1	
			the gradient of the normal in the given form correctly		A1	[5]
	(4.8)	**			3.64	
	(ii)	_	oduct rule		M1	
			correct derivative in any form derivative to zero and solve for <i>x</i>		A1 M1	
		Obtain:			A1	[4]
		Obtain.	$x - \frac{1}{2}$		AI	[4]
10	(i)	Express	is general point of l or m in component form, e.g. $(1 + s)$	1 - s, 1 + 2s	or	
			6+2t, 1+t)		B1	
		_	at least two corresponding pairs of components and solve f	or s or t	M1	
			s = -1 or t = -2		A1	
		Verify t	that all three component equations are satisfied		A1	[4]
	(ii)	Carry o	out correct process for evaluating the scalar product of the	direction vectors	of	
	()	l and m			M1	
		Using the	he correct process for the moduli, divide the scalar produc	t by the product	of	
		the mod	duli and evaluate the inverse cosine of the result		M1	
		Obtain	answer 74.2° (or 1.30 radians)		A1	[3]
	(iii)	EITHE	R: Use scalar product to obtain $a - b + 2c = 0$ and $2a + 2b$	+ c = 0	B1	
	()		Solve and obtain one ratio, e.g. $a:b$		M1	
			Obtain $a:b:c=5:-3:-4$, or equivalent		A 1	
			Substitute coordinates of a relevant point and values	for a, b and c	in	
			general equation of plane and evaluate d		M1	
			Obtain answer $5x - 3y - 4z = -2$, or equivalent		A1	
		<i>OR</i> 1:	Using two points on <i>l</i> and one on <i>m</i> , or <i>vice versa</i> , state	e three equations		
			a, b, c and d		B1	
			Solve and obtain one ratio, e.g. <i>a</i> : <i>b</i>	5 2 4	M1	
			Obtain a ratio of three of the unknowns, e.g. $a:b:c=$ Use coordinates of a relevant point and found ratio		A1	
			unknown, e.g. d	to find the four	ш М1	
			Obtain answer $-5x + 3y + 4z = 2$, or equivalent		A1	
		<i>OR</i> 2:	Form a correct 2-parameter equation for the plane,		711	
			e.g. $\mathbf{r} = \mathbf{i} + \mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + \mu(2\mathbf{i} + 2\mathbf{j} + \mathbf{k})$		B1	
			State three equations in x, y, z, λ and μ		M1	
			State three correct equations		A1	
			Eliminate λ and μ		M1	
			Obtain answer $5x - 3y - 4z = -2$, or equivalent		A1	
		<i>OR</i> 3:	Attempt to calculate vector product of direction vectors	s of l and m	M1	
			Obtain two correct components of the product		A1	
			Obtain correct product, e.g. $-5i + 3j + 4k$		A1	
			Form a plane equation and use coordinates of a	relevant point		
			calculate d		M1	[<i>5</i>]
			Obtain answer $-5x + 3y + 4z = 2$, or equivalent		A1	[5]