# Paper Reference(s) 6683 Edexcel GCE

## **Statistics S1**

## **Advanced Subsidiary**

## Friday 14 January 2005 – Morning

### Time: 1 hour 30 minutes

<u>Materials required for examination</u> Mathematical Formulae (Lilac) Graph Paper (ASG2) **Items included with question papers** Nil

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

### **Instructions to Candidates**

In the boxes on the answer book, write the name of the examining body (Edexcel), your centre number, candidate number, the unit title (Statistics S1), the paper reference (6683), your surname, other name and signature.

Values from the statistical tables should be quoted in full. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

### **Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. This paper has seven questions. The total mark for this paper is 75.

#### **Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled. You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

N16741A

This publication may only be reproduced in accordance with London Qualifications Limited copyright policy. ©2005 London Qualifications Limited

- 1. A company assembles drills using components from two sources. Goodbuy supplies 85% of the components and Amart supplies the rest. It is known that 3% of the components supplied by Goodbuy are faulty and 6% of those supplied by Amart are faulty.
  - (a) Represent this information on a tree diagram.

An assembled drill is selected at random.

- (b) Find the probability that it is not faulty.
- 2. The number of caravans on Seaview caravan site on each night in August last year is summarised in the following stem and leaf diagram.

| Caravans |        |   |   | 1 0 means 10 |   |   |   |   | Tota | Totals |   |  |
|----------|--------|---|---|--------------|---|---|---|---|------|--------|---|--|
| 1        | 0      | 5 |   |              |   |   |   |   |      | (2)    | ) |  |
|          | 1      |   |   |              |   |   |   |   |      | (4)    | ) |  |
| 3        | 0<br>1 | 3 | 3 | 3            | 4 | 7 | 8 | 8 |      | (8)    | ) |  |
| 4        | 1      | 1 | 3 | 5            | 8 | 8 | 8 | 9 | 9    | (9)    | ) |  |
| 5        | 2      | 3 | 6 | 6            | 7 |   |   |   |      | (5)    | ) |  |
| 6        | 2      | 3 | 4 |              |   |   |   |   |      | (3)    | ) |  |

(a) Find the three quartiles of these data.

During the same month, the least number of caravans on Northcliffe caravan site was 31. The maximum number of caravans on this site on any night that month was 72. The three quartiles for this site were 38, 45 and 52 respectively.

(b) On graph paper and using the same scale, draw box plots to represent the data for both caravan sites. You may assume that there are no outliers.

|              |   | (6)          |
|--------------|---|--------------|
| ( <i>c</i> ) | Compare and contrast these two box plots.                                 | ( <b>2</b> ) |
| ( 1)         |   | (3)          |
| (d)          | Give an interpretation to the upper quartiles of these two distributions. | (2)          |
|              |   | (-)          |

(3)

(3)

(3)

 $( \cap$ 

3. The following table shows the height x, to the nearest cm, and the weight y, to the nearest kg, of a random sample of 12 students.

| x | 148 | 164 | 156 | 172 | 147 | 184 | 162 | 155 | 182 | 165 | 175 | 152 |
|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| у | 39  | 59  | 56  | 77  | 44  | 77  | 65  | 49  | 80  | 72  | 70  | 52  |

- (a) On graph paper, draw a scatter diagram to represent these data.
- (b) Write down, with a reason, whether the correlation coefficient between x and y is positive or negative.

The data in the table can be summarised as follows.

$$\Sigma x = 1962$$
,  $\Sigma y = 740$ ,  $\Sigma y^2 = 47746$ ,  $\Sigma xy = 122783$ ,  $S_{xx} = 1745$ .

(c) Find  $S_{xy}$ .

The equation of the regression line of y on x is y = -106.331 + bx.

- (d) Find, to 3 decimal places, the value of b.
- (e) Find, to 3 significant figures, the mean  $\overline{y}$  and the standard deviation s of the weights of this sample of students.
- (3) (f) Find the values of  $\overline{y} \pm 1.96s$ .
- (2)
- (g) Comment on whether or not you think that the weights of these students could be modelled by a normal distribution. (1)

(2)

(3)

(2)

(2)

### **4.** The random variable *X* has probability function

|                                    | $\mathbf{P}(X=x)=kx,$ | x = 1, 2,, 5. |     |
|------------------------------------|-----------------------|---------------|-----|
| (a) Show that $k = \frac{1}{15}$ . |                       |               | (2) |
| Find                               |                       |               |     |
| ( <i>b</i> ) $P(X < 4)$ ,          |                       |               | (2) |
| (c) $E(X)$ ,                       |                       |               |     |
| ( <i>d</i> ) $E(3X-4)$ .           |                       |               | (2) |
|                                    |                       |               | (2) |

5. Articles made on a lathe are subject to three kinds of defect, *A*, *B* or *C*. A sample of 1000 articles was inspected and the following results were obtained.

| <ul> <li>31 had a type A defect</li> <li>37 had a type B defect</li> <li>42 had a type C defect</li> <li>11 had both type A and type B defects</li> <li>13 had both type B and type C defects</li> <li>10 had both type A and type C defects</li> <li>6 had all three types of defect.</li> </ul> |             |
|---|-------------|
| ( <i>a</i> ) Draw a Venn diagram to represent these data.   |             |
| Find the probability that a randomly selected article from this sample had  | (6)         |
| (b) no defects,   | (1)         |
| (c) no more than one of these defects.  |             |
| An article selected at random from this sample had only one defect.   | (2)         |
| ( <i>d</i> ) Find the probability that it was a type <i>B</i> defect.   |             |
|   | (2)         |
| Two different articles were selected at random from this sample.  |             |
| ( <i>e</i> ) Find the probability that both had type <i>B</i> defects.  | (2)         |
| A discrete random variable is such that each of its values is assumed to be equally likely.   |             |
| ( <i>a</i> ) Write down the name of the distribution that could be used to model this random varial   | ble.<br>(1) |
| (b) Give an example of such a distribution.   |             |

(c) Comment on the assumption that each value is equally likely.
(d) Suggest how you might refine the model in part (a).

6.


7. The random variable *X* is normally distributed with mean 79 and variance 144.

Find

(a) 
$$P(X < 70)$$
, (3)

(b) 
$$P(64 < X < 96)$$
.

It is known that  $P(79 - a \le X \le 79 + b) = 0.6463$ . This information is shown in the figure below.



Given that  $P(X \ge 79 + b) = 2P(X \le 79 - a)$ ,

(c) show that the area of the shaded region is 0.1179.

(d) Find the value of b.

**TOTAL FOR PAPER:75 MARKS** 

(3)

(3)

(4)

END