


Question Number	Scheme	Marks
1.	<p>(a)</p> <p>Shape unchanged Point</p>	B1 B1 (2)
(b)	<p>Shape Point</p>	B1 B1 (2)
(c)	<p>Shape (2, 4) (-2, 4)</p>	B1 B1 B1 [7]

Question Number	Scheme	Marks
2.	$x^2 - x - 2 = (x-2)(x+1)$ $\frac{2x^2 + 3x}{(2x+3)(x-2)} = \frac{x(2x+3)}{(2x+3)(x-2)} = \frac{x}{x-2}$ $\frac{2x^2 + 3x}{(2x+3)(x-2)} - \frac{6}{x^2 - x - 2} = \frac{x(x+1) - 6}{(x-2)(x+1)}$ $= \frac{x^2 + x - 6}{(x-2)(x+1)}$ $= \frac{(x+3)(x-2)}{(x-2)(x+1)}$ $= \frac{x+3}{x+1}$	At any stage B1 B1 M1 A1 M1 A1 A1 (7) [7]
	Alternative method	
	$x^2 - x - 2 = (x-2)(x+1)$ $(2x+3) \text{ appearing as a factor of the numerator at any stage}$ $\frac{2x^2 + 3x}{(2x+3)(x-2)} - \frac{6}{(x-2)(x+1)} = \frac{(2x^2 + 3x)(x+1) - 6(2x+3)}{(2x+3)(x-2)(x+1)}$ $= \frac{2x^3 + 5x^2 - 9x - 18}{(2x+3)(x-2)(x+1)}$ $= \frac{(x-2)(2x^2 + 9x + 9)}{(2x+3)(x-2)(x+1)} \quad \text{or} \quad \frac{(2x+3)(x^2 + x - 6)}{(2x+3)(x-2)(x+1)} \quad \text{or} \quad \frac{(x+3)(2x^2 - x - 6)}{(2x+3)(x-2)(x+1)}$ $= \frac{(2x+3)(x-2)(x+3)}{(2x+3)(x-2)(x+1)}$ $= \frac{x+3}{x+1}$	At any stage B1 B1 M1 A1 can be implied M1 A1 A1 (7)

Question Number	Scheme	Marks
3.	$\frac{dy}{dx} = \frac{1}{x}$ accept $\frac{3}{3x}$ At $x = 3$ , $\frac{dy}{dx} = \frac{1}{3} \Rightarrow m' = -3$ Use of $mm' = -1$ $y - \ln 1 = -3(x - 3)$ $y = -3x + 9$ Accept $y = 9 - 3x$ $\frac{dy}{dx} = \frac{1}{3x}$ leading to $y = -9x + 27$ is a maximum of M1 A0 M1 M1 A0 = 3/5	M1 A1 M1 M1 A1 (5) [5]
4.	(a) (i) $\frac{d}{dx}(e^{3x+2}) = 3e^{3x+2}$ (or $3e^2 e^{3x}$ ) At any stage $\frac{dy}{dx} = 3x^2 e^{3x+2} + 2x e^{3x+2}$ Or equivalent (ii) $\frac{d}{dx}(\cos(2x^3)) = -6x^2 \sin(2x^3)$ At any stage $\frac{dy}{dx} = \frac{-18x^3 \sin(2x^3) - 3 \cos(2x^3)}{9x^2}$ Alternatively using the product rule for second M1 A1 $y = (3x)^{-1} \cos(2x^3)$ $\frac{dy}{dx} = -3(3x)^{-2} \cos(2x^3) - 6x^2 (3x)^{-1} \sin(2x^3)$ Accept equivalent unsimplified forms (b) $1 = 8 \cos(2y+6) \frac{dy}{dx}$ or $\frac{dx}{dy} = 8 \cos(2y+6)$ $\frac{dy}{dx} = \frac{1}{8 \cos(2y+6)}$ $\frac{dy}{dx} = \frac{1}{8 \cos\left(\arcsin\left(\frac{x}{4}\right)\right)}$ $\left(= (\pm) \frac{1}{2\sqrt{16-x^2}}\right)$	B1 M1 A1+A1 (4) M1 A1 M1 A1 (4) M1 A1 M1 A1 (5) [13]

Question Number	Scheme	Marks
5.	<p>(a) <math>2x^2 - 1 - \frac{4}{x} = 0</math> Dividing equation by <math>x</math>  <math>x^2 = \frac{1}{2} + \frac{4}{2x}</math> Obtaining <math>x^2 = \dots</math>  <math>x = \sqrt{\left(\frac{1}{2} + \frac{4}{2x}\right)} *</math> cso</p> <p>(b) <math>x_1 = 1.41, x_2 = 1.39, x_3 = 1.39</math>  If answers given to more than 2 dp, penalise first time then accept awrt above.</p> <p>(c) Choosing <math>(1.3915, 1.3925)</math> or a tighter interval  <math>f(1.3915) \approx -3 \times 10^{-3}, f(1.3925) \approx 7 \times 10^{-3}</math> Both, awrt  Change of sign (and continuity) <math>\Rightarrow \alpha \in (1.3915, 1.3925)</math>  <math>\Rightarrow \alpha = 1.392</math> to 3 decimal places * cso</p>	M1 M1 A1 (3) B1, B1, B1 (3) M1 A1 A1 (3) [9]
6.	<p>(a) <math>R \cos \alpha = 12, R \sin \alpha = 4</math>  <math>R = \sqrt{(12^2 + 4^2)} = \sqrt{160}</math> Accept if just written down, awrt 12.6  <math>\tan \alpha = \frac{4}{12}, \Rightarrow \alpha \approx 18.43^\circ</math> awrt <math>18.4^\circ</math></p> <p>(b) <math>\cos(x + \text{their } \alpha) = \frac{7}{\text{their } R} (\approx 0.5534)</math>  <math>x + \text{their } \alpha = 56.4^\circ</math> awrt <math>56^\circ</math>  <math>= \dots, 303.6^\circ</math> <math>360^\circ - \text{their principal value}</math>  <math>x = 38.0^\circ, 285.2^\circ</math> Ignore solutions out of range  If answers given to more than 1 dp, penalise first time then accept awrt above.</p> <p>(c)(i) minimum value is <math>-\sqrt{160}</math> ft their <math>R</math>  (ii) <math>\cos(x + \text{their } \alpha) = -1</math>  <math>x \approx 161.57^\circ</math> cao</p>	M1 A1 M1, A1(4) M1 A1 M1 A1, A1 (5) B1ft M1 A1 (3) [12]

Question Number	Scheme	Marks
7.	(a) (i) Use of $\cos 2x = \cos^2 x - \sin^2 x$ in an attempt to prove the identity. $\frac{\cos 2x}{\cos x + \sin x} = \frac{\cos^2 x - \sin^2 x}{\cos x + \sin x} = \frac{(\cos x - \sin x)(\cos x + \sin x)}{\cos x + \sin x} = \cos x - \sin x *$ cso	M1 A1 (2)
	(ii) Use of $\cos 2x = 2\cos^2 x - 1$ in an attempt to prove the identity. Use of $\sin 2x = 2\sin x \cos x$ in an attempt to prove the identity. $\frac{1}{2}(\cos 2x - \sin 2x) = \frac{1}{2}(2\cos^2 x - 1 - 2\sin x \cos x) = \cos^2 x - \cos x \sin x - \frac{1}{2} *$ cso	M1 M1 A1 (3)
(b)	$\cos \theta (\cos \theta - \sin \theta) = \frac{1}{2}$ $\cos^2 \theta - \cos \theta \sin \theta - \frac{1}{2} = 0$ $\frac{1}{2}(\cos 2\theta - \sin 2\theta) = 0$ $\cos 2\theta = \sin 2\theta *$	Using (a)(i) M1 Using (a)(ii) A1 (3)
(c)	$\tan 2\theta = 1$ $2\theta = \frac{\pi}{4}, \left( \frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4} \right)$ any one correct value of $2\theta$ $\theta = \frac{\pi}{8}, \frac{5\pi}{8}, \frac{9\pi}{8}, \frac{13\pi}{8}$ Obtaining at least 2 solutions in range The 4 correct solutions If decimals (0.393, 1.963, 3.534, 5.105) or degrees (22.5°, 112.5°, 202.5°, 292.5°) are given, but all 4 solutions are found, penalise one A mark only. Ignore solutions out of range.	M1 A1 M1 A1 (4) [12]

Question Number	Scheme	Marks
8.	(a) $\begin{aligned} gf(x) &= e^{2(2x+\ln 2)} \\ &= e^{4x}e^{2\ln 2} \\ &= e^{4x}e^{\ln 4} \\ &= 4e^{4x} \end{aligned}$ <p style="text-align: right;">Give mark at this point, cso</p> <p>(Hence <math>gf : x \mapsto 4e^{4x}, x \in \mathbb{R}</math>)</p>	M1 M1 M1 A1 (4)
(b)	 <p style="text-align: right;">Shape and point</p>	B1 (1)
(c)	Range is $\mathbb{R}_+$	Accept $gf(x) > 0, y > 0$ B1 (1)
(d)	$\begin{aligned} \frac{d}{dx}[gf(x)] &= 16e^{4x} \\ e^{4x} &= \frac{3}{16} \\ 4x &= \ln \frac{3}{16} \\ x &\approx -0.418 \end{aligned}$	M1 A1 M1 A1 (4) [10]