







2.




Figure 1

Figure 1 shows a ladder  $AB$ , of mass 25 kg and length 4 m, resting in equilibrium with one end  $A$  on rough horizontal ground and the other end  $B$  against a smooth vertical wall. The ladder is in a vertical plane perpendicular to the wall. The coefficient of friction between the ladder and the ground is  $\frac{11}{25}$ . The ladder makes an angle  $\beta$  with the ground. When Reece, who has mass 75 kg, stands at the point  $C$  on the ladder, where  $AC = 2.8$  m, the ladder is on the point of slipping. The ladder is modelled as a uniform rod and Reece is modelled as a particle.

- (a) Find the magnitude of the frictional force of the ground on the ladder. (3)
- (b) Find, to the nearest degree, the value of  $\beta$ . (6)
- (c) State how you have used the modelling assumption that Reece is a particle. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



















5.




Figure 2

A uniform lamina  $ABCD$  is made by joining a uniform triangular lamina  $ABD$  to a uniform semi-circular lamina  $DBC$ , of the same material, along the edge  $BD$ , as shown in Figure 2. Triangle  $ABD$  is right-angled at  $D$  and  $AD = 18$  cm. The semi-circle has diameter  $BD$  and  $BD = 12$  cm.

- (a) Show that, to 3 significant figures, the distance of the centre of mass of the lamina  $ABCD$  from  $AD$  is 4.69 cm. (4)

Given that the centre of mass of a uniform semicircular lamina, radius  $r$ , is at a distance  $\frac{4r}{3\pi}$  from the centre of the bounding diameter,

- (b) find, in cm to 3 significant figures, the distance of the centre of mass of the lamina  $ABCD$  from  $BD$ . (4)

The lamina is freely suspended from  $B$  and hangs in equilibrium.

- (c) Find, to the nearest degree, the angle which  $BD$  makes with the vertical. (4)

---

---

---

---

---

---

---

---

---

---












Figure 3

A cricket ball is hit from a point  $A$  with velocity of  $(p\mathbf{i} + q\mathbf{j}) \text{ m s}^{-1}$ , at an angle  $\alpha$  above the horizontal. The unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are respectively horizontal and vertically upwards. The point  $A$  is 0.9 m vertically above the point  $O$ , which is on horizontal ground.

The ball takes 3 seconds to travel from  $A$  to  $B$ , where  $B$  is on the ground and  $OB = 57.6 \text{ m}$ , as shown in Figure 3. By modelling the motion of the cricket ball as that of a particle moving freely under gravity,

- (a) find the value of  $p$ , (2)
- (b) show that  $q = 14.4$ , (3)
- (c) find the initial speed of the cricket ball, (2)
- (d) find the exact value of  $\tan \alpha$ . (1)
- (e) Find the length of time for which the cricket ball is at least 4 m above the ground. (6)
- (f) State an additional physical factor which may be taken into account in a refinement of the above model to make it more realistic. (1)

---

---

---

---

---

---

---

---

---

---



















**BLANK PAGE**

