Surname	Other na	mes
Surrume	Otherna	
Pearson	Centre Number	Candidate Number
Edexcel GCE		
Further F Mathema	atics FP3	
Advanced/Advanced	led Subsidiary	
Monday 27 June 2016 – Time: 1 hour 30 minute	Morning	Paper Reference 6669/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

PEARSON

Turn over ▶



(4)

1. $\mathbf{A} = \begin{pmatrix} -2 & 1 & -3 \\ k & 1 & 3 \\ 2 & -1 & k \end{pmatrix}, \text{ where } k \text{ is a constant}$

Given that the matrix A	A is singular	, find the pos	sible values of k .

	blank
Question 1 continued	
	Q1
(Total 4 marks)	



2. The curve C has equation

$$y = \frac{x^2}{8} - \ln x, \quad 2 \leqslant x \leqslant 3$$

Find the length of the curve <i>C</i>	giving your	answer in	the form I	$p + \ln q$,	where p	and q	are
rational numbers to be found.							

	(1)



Question 2 continued	blank
	Q2
(Total 7 marks)	



3. (a) Prove that

$$\frac{d(\operatorname{arcoth} x)}{\mathrm{d}x} = \frac{1}{1 - x^2}$$

(3)

Given that $y = (\operatorname{arcoth} x)^2$,

(b) show that

$$(1 - x^2)\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} = \frac{k}{1 - x^2}$$

where k is a constant to be determined.

(5)

	Leave blank
Question 3 continued	



Question 3 continue	ed	

Question 3 continued	blank
	Q3
(Total 8 marks)	



4. (i) Find, without using a calculator,

$$\int_{3}^{5} \frac{1}{\sqrt{15 + 2x - x^2}} \, \mathrm{d}x$$

giving your answer as a multiple of π .

(5)

(ii)

(a) Show that

$$5\cosh x - 4\sinh x = \frac{e^{2x} + 9}{2e^x}$$

(3)

(b) Hence, using the substitution $u = e^x$ or otherwise, find

$$\int \frac{1}{5\cosh x - 4\sinh x} \, \mathrm{d}x$$

(4)

	Leave
	blank
Question 4 continued	
Question 4 continued	
	1



Question 4 continued	

Question 4 continued	blank
	0.4
	Q4
(Total 12 marks)	



5. The hyperbola H has equation

$$\frac{x^2}{16} - \frac{y^2}{9} = 1$$

The point P (4 sec θ , 3 tan θ), $0 < \theta < \frac{\pi}{2}$, lies on H.

(a) Show that an equation of the normal to H at the point P is

$$3y + 4x\sin\theta = 25\tan\theta$$

(5)

The line l is the directrix of H for which x > 0

The normal to H at P crosses the line l at the point Q. Given that $\theta = \frac{\pi}{4}$

(b) find the y coordinate of Q, giving your answer in the form $a + b\sqrt{2}$, where a and b are rational numbers to be found.

•	•		
	n		
	v	,	

	Leave
	blank
Question 5 continued	
Question 3 continued	



	Leave
Question 5 continued	blank
Question 3 continued	

Question 5 continued	blank
	Q5
(Tota	l 11 marks)



6.
$$\mathbf{M} = \begin{pmatrix} p & -2 & 0 \\ -2 & 6 & -2 \\ 0 & -2 & q \end{pmatrix}$$

where p and q are constants.

Given that $\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$ is an eigenvector of the matrix **M**,

(a) find the eigenvalue corresponding to this eigenvector,

(3)

(b) find the value of p and the value of q.

(3)

Given that 6 is another eigenvalue of M,

(c) find a corresponding eigenvector.

(2)

Given that $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ is a third eigenvector of **M** with eigenvalue 3

(d) find a matrix **P** and a diagonal matrix **D** such that

$$\mathbf{P}^{\mathrm{T}}\mathbf{M}\mathbf{P} = \mathbf{D}$$

(3)



	Le bl
Question 6 continued	



Question 6 continued	

uestion 6 continued	



Question 6 continued	

Question 6 continued	blank
	Q6
(Total 11 marks)	



(3)

7. Given that

$$I_n = \int \frac{\sin nx}{\sin x} \, \mathrm{d}x, \quad n \geqslant 1$$

(a) prove that, for $n \ge 3$

$$I_n - I_{n-2} = \int 2\cos(n-1)x \,\mathrm{d}x$$

(b) Hence, showing each step of your working, find the exact value of

$$\int_{\frac{\pi}{12}}^{\frac{\pi}{6}} \frac{\sin 5x}{\sin x} \, \mathrm{d}x$$

giving your answer in the form $\frac{1}{12}(a\pi + b\sqrt{3} + c)$, where a, b and c are integers to be found. (7)

	Leave
	blank
Question 7 continued	



Question 7 continued		

	Leave blank
Question 7 continued	
	Q7
(Total 10 marks)	



8. The plane Π_1 has equation

$$x - 5y - 2z = 3$$

The plane Π_2 has equation

$$\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}) + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$$

where λ and μ are scalar parameters.

(a) Show that Π_1 is perpendicular to Π_2

(4)

(b) Find a cartesian equation for Π_2

(2)

(c) Find an equation for the line of intersection of Π_1 and Π_2 giving your answer in the form $(\mathbf{r} - \mathbf{a}) \times \mathbf{b} = \mathbf{0}$, where \mathbf{a} and \mathbf{b} are constant vectors to be found.

(6)

	_
	_
	_
	_
	_

	Leave
	blank
Question 8 continued	



Question 8 continued

	Leave
	blank
Question 8 continued	

