


June 2005
6673 Pure P3
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) Finding $f(\pm 2)$, and obtaining $16 - 32 + 10 + 6 = 0$ Or uses division and obtains $2x^2 - kx\dots$, obtaining $2x^2 - 4x - 3$ and concluding remainder = 0</p> <p>(b) Finding $f(\pm \frac{1}{2})$, and obtaining $-\frac{1}{4} - 2 - \frac{5}{2} + 6 = 1\frac{1}{4}$ Or uses division and obtains $x^2 - kx\dots$, obtaining $x^2 - \frac{9}{2}x + \frac{19}{4}$ and concluding remainder = $\frac{5}{4}$</p> <p>(c) $x = 2$ (also allow $\frac{2 \pm \sqrt{10}}{2}$ or $\frac{4 \pm \sqrt{40}}{4}$)</p>	<p>M1, A1 M1 A1</p> <p>M1, A1</p> <p>B1</p> <p style="text-align: right;">(5)</p>
2.	<p>(a) Writes down binomial expansion up to and including term in x^3, allow nC_r notation $1 + na + n(n-1)\frac{a^2x^2}{2} + \frac{n(n-1)(n-2)}{6}a^3x^3$ (condone errors in powers of a)</p> <p>States $na = 15$</p> <p>Puts $\frac{n(n-1)a^2}{2} = \frac{n(n-1)(n-2)a^3}{6}$ (condone errors in powers of a)</p> <p>$3 = (n-2)a$</p> <p>Solves simultaneous equations in n and a to obtain $a = 6$, and $n = 2.5$</p> <p>[n.b. Just writes $a = 6$, and $n = 2.5$ following no working or following errors allow the last M1 A1 A1]</p> <p>(b) Coefficient of $x^3 = 2.5 \times 1.5 \times 0.5 \times 6^3 \div 6 = 67.5$ (or equals coefficient of $x^2 = 2.5 \times 1.5 \times 6^2 \div 2 = 67.5$)</p>	<p>M1</p> <p>B1</p> <p>dM1</p> <p>M1 A1 A1 (6)</p> <p>B1 (1)</p> <p>[7]</p>

Question Number	Scheme	Marks
3. (a)	<p>Attempt at integration by parts, i.e. $kx \sin 2x \pm \int k \sin 2x dx$, with $k = 2$ or $\frac{1}{2}$ $= \frac{1}{2} x \sin 2x - \int \frac{1}{2} \sin 2x dx$</p> <p>Integrates $\sin 2x$ correctly, to obtain $\frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + c$ (penalise lack of constant of integration first time only)</p>	<p>M1 A1 M1, A1 (4)</p>
(b)	<p>Hence method : Uses $\cos 2x = 2 \cos^2 x - 1$ to connect integrals</p> <p>Obtains $\int x \cos^2 x dx = \frac{1}{2} \left\{ \frac{x^2}{2} + \text{answer to part(a)} \right\} = \frac{x^2}{4} + \frac{x}{4} \sin 2x + \frac{1}{8} \cos 2x + k$</p> <p>Otherwise method $\int x \cos^2 x dx = x \left(\frac{1}{4} \sin 2x + \frac{x}{2} \right) - \int \frac{1}{4} \sin 2x + \frac{x}{2} dx$ B1 for $(\frac{1}{4} \sin 2x + \frac{x}{2})$ $= \frac{x^2}{4} + \frac{x}{4} \sin 2x + \frac{1}{8} \cos 2x + k$</p>	<p>B1 M1 A1 (3) B1, M1 A1 (3)</p>
4 (a)	<p>$r = 3$ (both circles)</p> <p>Centres are at (2, 0) and (5, 0)</p>	<p>B1 B1, B1 (3)</p>
(b)	 <p>1st circle correct quadrants centre on x axis</p> <p>2nd circle correct quadrants centre on x axis</p> <p>circles same size and passing through centres of other circle</p>	<p>B1 B1 B1 (3)</p>
(c)	<p>Finds circles meet at $x = 3.5$, by mid point of centres or by solving algebraically</p> <p>Establishes $y = \pm \frac{3\sqrt{3}}{2}$, and thus distance is $3\sqrt{3}$.</p> <p>Or uses trig or Pythagoras with lengths 3, angles 60 degrees, or 120 degrees. Complete and accurate method to find required distance Establishes distance is $3\sqrt{3}$.</p>	<p>M1 M1, A1 (3) M1 M1 A1 (3)</p>

Question Number	Scheme	Marks
5.	<p>(a) Substitutes $t = 4$ to give $V, = 1975.31$ or 1975.30 or 1975 or 1980 (3 s.f)</p> <p>(b) $\frac{dV}{dt} = -\ln 1.5 \times V ; = -800.92$ or -800.9 or -801 M1 needs $\ln 1.5$ term</p> <p>(c) rate of decrease in value on 1st January 2005</p>	<p>M1 , A1 (2)</p> <p>M1 A1; A1 (3)</p> <p>B1 (1)</p>
6,	<p>(a) $\overline{AB} = \begin{pmatrix} c \\ d-5 \\ 10 \end{pmatrix} = k \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix}$ or $11+5\lambda = 21, \Rightarrow \lambda = 2$, $\therefore c = 4$ $d = 7$</p> <p>(b) $\begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 2\lambda \\ 5+\lambda \\ 11+5\lambda \end{pmatrix} = 0$ $\therefore 4\lambda + 5 + \lambda + 55 + 25\lambda = 0$ $\therefore \lambda = -2$</p> <p>Substitutes to give the point $P, -4\mathbf{i} + 3\mathbf{j} + \mathbf{k}$ (Accept $(-4, 3, 1)$)</p> <p>(c) Finds the length of OA, or OB or OP or AB as $\sqrt{146}$ or $\sqrt{506}$ or $\sqrt{26}$ or $\sqrt{120}$ resp. Uses area formula- either Area = $\frac{1}{2} \mathbf{AB} \times \mathbf{OP}$ or $= \frac{1}{2} \mathbf{OA} \times \mathbf{OB} \sin \angle AOB$ or $= \frac{1}{2} \mathbf{OA} \times \mathbf{AB} \sin \angle OAB$ or $= \frac{1}{2} \mathbf{AB} \times \mathbf{OB} \sin \angle ABO$ $= \frac{1}{2}\sqrt{120}\sqrt{26}$ or $\frac{1}{2}\sqrt{146}\sqrt{506} \sin 11.86$ or $\frac{1}{2}\sqrt{146}\sqrt{120} \sin 155.04$ or $\frac{1}{2}\sqrt{120}\sqrt{506} \sin 13.10$ $= 27.9$</p>	<p>M1 , A1 A1 (3)</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1, A1 (6)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1 (4)</p>

Question Number	Scheme	Marks
7 (a)	<p>As $V = \frac{4}{3}\pi r^3$, then $\frac{dV}{dr} = 4\pi r^2$</p> <p>Using chain rule $\frac{dr}{dt} = \frac{dV}{dt} \div \frac{dV}{dr} = \frac{k}{\frac{4}{3}\pi r^3} \times \frac{1}{4\pi r^2}$</p> $= \frac{B}{r^5} \quad *$ <p>(b) $\int r^5 dr = \int B dt$</p> $\therefore \frac{r^6}{6} = Bt + c \quad (\text{allow mark at this stage, does not need } r =)$ <p>(c) Use $r = 5$ at $t = 0$ to give $c = \frac{5^6}{6}$ or 2604 or 2600</p> <p>Use $r = 6$ at $t = 2$ to give $B = \frac{6^5}{2} - \frac{5^6}{12}$ or 2586 or 2588 or 2590</p> <p>Put $t = 4$ to obtain r^6 (approx 78000)</p> <p>Then take sixth root to obtain $r = 6.53$ (cm)</p>	<p>M1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>B1</p> <p>M1 A1 (3)</p> <p>M1</p> <p>M1</p> <p>M1 A1</p> <p>A1 (5)</p>

Question Number	Scheme	Marks
8. (a)	$\frac{dx}{dt} = -\frac{1}{(1+t)^2} \quad \text{and} \quad \frac{dy}{dt} = \frac{1}{(1-t)^2}$ $\therefore \frac{dy}{dx} = \frac{-(1+t)^2}{(1-t)^2} \quad \text{and at } t = \frac{1}{2}, \text{ gradient is } -9$ <p>M1 requires their dy/dt / their dx/dt and substitution of t.</p> <p>At the point of contact $x = \frac{2}{3}$ and $y = 2$</p> <p>Equation is $y - 2 = -9(x - \frac{2}{3})$</p>	<p>B1, B1</p> <p>M1 A1cao</p> <p>B1</p> <p>M1 A1</p> <p>(7)</p>
(b)	<p>Either obtain t in terms of x and y i.e, $t = \frac{1}{x} - 1$ or $t = 1 - \frac{1}{y}$ (or both)</p> <p>Then substitute into other expression $y = f(x)$ or $x = g(y)$ and rearrange (or put $\frac{1}{x} - 1 = 1 - \frac{1}{y}$ and rearrange)</p> <p>To obtain $y = \frac{x}{2x-1}$ *</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p>
	<p>Or Substitute into $\frac{x}{2x-1} = \frac{(1+t)}{\frac{2}{1+t} - 1}$</p> $= \frac{1}{2-(1+t)} = \frac{1}{1-t}$ $= y *$	<p>M1</p> <p>A1</p> <p>M1</p> <p>(3)</p>
(c)	<p>Area = $\int_{\frac{2}{3}}^1 \frac{x}{2x-1} dx$</p> $= \int \frac{u+1}{2u} \frac{du}{2} = \frac{1}{4} \int 1 + \frac{1}{u} du$ <p>putting into a form to integrate</p> $= \left[\frac{1}{4}u + \frac{1}{4} \ln u \right]_{\frac{2}{3}}^1$ $= \frac{1}{4} - \left(\frac{1}{12} + \frac{1}{4} \ln \frac{1}{3} \right)$ $= \frac{1}{6} + \frac{1}{4} \ln 3 \text{ or any correct equivalent.}$	<p>B1</p> <p>M1</p> <p>M1 A1</p> <p>M1</p> <p>A1</p> <p>(6)</p>

