



Pearson
Edexcel

Mark Scheme (Provisional)

Summer 2021

Pearson Edexcel International Advanced Level
In Statistics S2 Paper WST02/01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2021

Question Paper Log number P63151A

Publications Code WST02_01_2106_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
 7. Ignore wrong working or incorrect statements following a correct answer

Question Number	Scheme	Marks						
Throughout the paper the candidates may use different letters to the ones given in the mark scheme.								
1.	<p>(a) [$X \sim$ the number of pansy seeds that do not germinate or $Y =$ the number...that <u>do</u> germinate] $X \sim B(20, 0.05)$ <u>or</u> $Y \sim B(20, 0.95)$</p> <p>(i) $P(X \leq 4) - P(X \leq 2) = 0.9974 - 0.9245$ <u>or</u> $\binom{20}{3} 0.05^3 \times 0.95^{17} + \binom{20}{4} 0.05^4 \times 0.95^{16} = 0.05958... + 0.01332...$ $= 0.072909...$ awrt 0.0729</p> <p>(ii) $P(X \leq 1)$ <u>or</u> $P(Y \geq 19) = 20 \times (0.95)^{19} (0.05) + (0.95)^{20}$ $= 0.7358$ <u>or</u> $= 0.735839...$ awrt 0.736</p> <p>(b) [Let $W =$ no. of packets where $Y > 18$] $P(W = 5) = ("0.7358...")^5$ $= 0.21573...$ awrt 0.216</p> <p>(c) $H_0 : p = 0.05$ $H_1 : p > 0.05$</p> <p>(d) [$V =$ no. of seeds that do not germinate $V \sim B(100, 0.05)$ approximates to] $V \sim Po(5)$</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">CR for 1-tail in (c)</th> <th style="text-align: center;">CR for 2-tail in (c)</th> </tr> </thead> <tbody> <tr> <td>$P(V \geq 8) = 1 - P(V \leq 7)$ $= 1 - 0.8666$ $= 0.1334$</td> <td>$P(V \geq 9) = 0.0681$ $P(V \geq 10) = 0.0318$ CR $V \geq 10$ oe</td> <td>$P(V \geq 10) = 0.0318$ $P(V \geq 11) = 0.0137$ CR $V \geq 11$ oe</td> </tr> </tbody> </table> <p>Accept H_0 <u>or</u> not significant <u>or</u> 8 does not lie in the critical region Data consistent with <i>Spany's claim</i> <u>or</u> Insufficient evidence for <i>Jem's belief</i></p>		CR for 1-tail in (c)	CR for 2-tail in (c)	$P(V \geq 8) = 1 - P(V \leq 7)$ $= 1 - 0.8666$ $= 0.1334$	$P(V \geq 9) = 0.0681$ $P(V \geq 10) = 0.0318$ CR $V \geq 10$ oe	$P(V \geq 10) = 0.0318$ $P(V \geq 11) = 0.0137$ CR $V \geq 11$ oe	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 (5)</p> <p>M1</p> <p>A1 (2)</p> <p>B1</p> <p>(1)</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>dM1</p> <p>A1cso</p> <p>(6)</p> <p>Total 14</p>
	CR for 1-tail in (c)	CR for 2-tail in (c)						
$P(V \geq 8) = 1 - P(V \leq 7)$ $= 1 - 0.8666$ $= 0.1334$	$P(V \geq 9) = 0.0681$ $P(V \geq 10) = 0.0318$ CR $V \geq 10$ oe	$P(V \geq 10) = 0.0318$ $P(V \geq 11) = 0.0137$ CR $V \geq 11$ oe						
Notes								
(a)	B1: writing or using $B(20, 0.05)$ [Allow $Y \sim B(20, 0.95)$ if Y is clearly defined]. Implied by 1 correct prob.							
(i)	M1: for $P(X \leq 4) - P(X \leq 2)$ <u>and</u> one correct prob. <u>or</u> $P(X = 3) + P(X = 4)$ <u>and</u> 1 correct prob.							
(ii)	M1: for $P(X \leq 1)$ <u>or</u> $[20] \times (0.95)^{19} (0.05) + (0.95)^{20}$ - condone missing 20							
(b)	M1: for $(\text{their(a)(ii)})^5$							
(c)	B1: both hypotheses correct with p or π							
(d)	1st M1: for realising a Poisson approximation is appropriate. NB $Po(95)$ is M0A0 1st A1: writing or using $V \sim Po(5)$ i.e correct mean for the Poisson. 2nd M1: for writing or using $1 - P(V \leq 7)$ <u>or</u> $P(V \leq 7) = 0.8666$ <u>or</u> writing $P(V \geq 10) = 0.0318$ <u>or</u> $P(V \geq 9) = 0.0681$ <u>or</u> $P(V \geq 11) = 0.0137$ leading to a CR. Implied by correct CR <u>or</u> probability = awrt 0.133 2nd A1: for awrt 0.133 <u>or</u> $V \geq 10$ oe (e.g. $V > 9$) <u>or</u> $V \geq 11$ oe allow any letter but CR must match part(c) 3rd dM1: dep on 2 nd M1. ft their CR or probability. A correct statement based on comparing 8 with their CR <u>or</u> their prob with 0.05 or 0.025 [condone $0.866 < 0.95$]– contradicting non-contextual comments M0 3rd A1 cso: all previous marks must be awarded. A correct statement in context. Need Bold words. NB award M1A1 for a correct contextual statement on its own. If there are no hypotheses or they are the wrong way around, then 3 rd M0 3 rd A0							
SC1	Normal approximation: Award marks in pairs with 2, 4 or 6 marks available Sight of $N(5 \text{ or } 95, \sqrt{4.75^2})$ M1A1; probability awrt 0.125/6 M1A1; Correct contextual concl' dM1A1							
SC2	No approximation: Use of $B(100, 0.05)$ M0A0; probability awrt 0.128 <u>or</u> $CR \geq 10$ M1A1; then M0A0							

Question Number	Scheme	Marks
<p>2. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>[$X =$ number of faults in 4 m² so $X \sim \text{Po}(3)$]</p> <p>$P(X = 5) = P(X \leq 5) - P(X \leq 4) [= 0.9161 - 0.8153]$ <u>or</u> $\frac{e^{-3}3^5}{5!}$ (allow λ instead of 3)</p> <p>$= 0.1008$ <u>or</u> 0.100818... awrt 0.101</p> <p>[$Y =$ number of faults in 6 m² so] $Y \sim \text{Po}(4.5)$ <u>and</u> [$P(Y > 5)$] = $1 - P(Y \leq 5)$ [= $1 - 0.7029$]</p> <p>$= 0.2971$ <u>or</u> (calc) 0.29706956... awrt 0.297</p> <p>0.101 (or ft their answer to (a))</p> <p>Faults occur independently/ randomly</p> <p>[$F =$ number of faults in a small rug] $F \sim \text{Po}(0.9)$</p> <p>$e^{-0.9n}n \times 80 + (1 - e^{-0.9n})n \times 60 \geq 4000$ <u>or</u> (awrt 0.407)$n \times 80 +$ (awrt 0.593)$n \times 60 \geq 4000$</p> <p>$n \geq \frac{4000}{20e^{-0.9} + 60} = 58.71\dots$</p> <p>$n = \underline{59}$</p> <p>$H_0 : \lambda = 9$ $H_1 : \lambda > 9$</p> <p>$R \sim \text{Po}(0.9 \times 10)$ <u>and</u> [$P(R \geq 13)$] = $1 - P(R \leq 12)$ [= $1 - 0.8758$]</p> <p>$P(R \leq 13) = 0.9261$ <u>or</u> $P(R \geq 14) = 0.0739$ <u>or</u> $P(R \leq 14) = 0.9585$ <u>or</u> $P(R \geq 15) = 0.0415$</p> <p>[$P(R \geq 13)$] = 0.1242 awrt 0.124 <u>or</u> CR $R \geq 15$ (oe)</p> <p>so insufficient evidence to reject H_0 /not significant/ not in critical region</p> <p>There is insufficient evidence that the rate at which faults occur is higher for Rhiannon</p>	<p>M1</p> <p>A1</p> <p>(2)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>B1ft</p> <p>B1</p> <p>(2)</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(5)</p> <p>Total 15</p>
Notes		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>M1: for using or writing $P(X \leq 5) - P(X \leq 4)$ <u>or</u> $\frac{e^{-\lambda}\lambda^5}{5!}$ (Accept letter λ or any value of λ)</p> <p>M1: writing or using Po(4.5) <u>and</u> sight of [$P(Y > 5)$] = $1 - P(Y \leq 5)$ Implied by sight of $1 - 0.7029$</p> <p>2nd B1: for a comment about faults occurring randomly/independently <u>or</u> Poisson has “no memory”</p> <p>B1: writing or using Po(0.9) May be implied by sight of 0.407 or 0.593</p> <p>1st M1: for $e^{-\lambda}n \times 80 + (1 - e^{-\lambda})n \times 60 > 4000$ any value for λ. Allow = 4000</p> <p>2nd M1: for solving their equation leading to a positive value of n. Allow any value of λ and allow $n = \dots$</p> <p>A1: for an answer of 59 only</p> <p>B1: both hypotheses correct with λ or μ. Allow 3 or 0.75 or 0.9 instead of 9</p> <p>1st M1: for writing or using Po(“9”) and writing or using $1 - P(R \leq 12)$ (implied by $1 - 0.8758$) <u>or</u> one of: $P(R \leq 13) = 0.9261$, $P(R \geq 14) = 0.0739$, $P(R \leq 14) = 0.9585$, $P(R \geq 15) = 0.0415$ leading to a CR</p> <p>1st A1: for probability = awrt 0.124 <u>or</u> CR of $R \geq 15$ oe e.g. $R > 14$</p> <p>2nd M1: for a correct conclusion based on their prob & 0.05 <u>or</u> their CR & 13. Assume correct hypotheses. Do not allow contradicting conclusions</p> <p>2nd A1: dep on both Ms for a correct contextual comment including the words in bold.</p>	

Question Number	Scheme	Marks									
<p>4.</p> <p>(i)</p> <p>Possible samples with a range of 4 are: (1,1,5) (1,2,5) (1,5,5)</p> <p>(1,1,5) $\frac{2}{9} \times \frac{2}{9} \times \frac{4}{9} \times 3 = \frac{16}{243}$ <u>or</u> (1,5,5) $\frac{2}{9} \times \frac{4}{9} \times \frac{4}{9} \times 3 = \frac{32}{243}$</p> <p>(1,2,5) $\frac{2}{9} \times \frac{1}{3} \times \frac{4}{9} \times 6 = \frac{16}{81}$</p> <p>$P(B=4) = \frac{16}{243} + \frac{32}{243} + \frac{16}{81} = \frac{32}{81}$</p> <p>(ii)</p> <p>$P(B=0) = \left(\frac{2}{9}\right)^3 + \left(\frac{1}{3}\right)^3 + \left(\frac{4}{9}\right)^3 = \frac{11}{81}$</p> <p>$P(B=1) = 3 \times \frac{2}{9} \times \left(\frac{1}{3}\right)^2 + 3 \times \frac{1}{3} \times \left(\frac{2}{9}\right)^2 = \frac{10}{81}$ <u>or</u> $P(B=3) = 3 \times \frac{1}{3} \times \left(\frac{4}{9}\right)^2 + 3 \times \frac{4}{9} \times \left(\frac{1}{3}\right)^2 = \frac{28}{81}$</p> <p>$1 - \frac{11}{81} - \frac{10}{81} - \frac{32}{81} = \frac{28}{81}$ <u>or</u> $1 - \frac{11}{81} - \frac{28}{81} - \frac{32}{81} = \frac{10}{81}$</p> <table border="1" data-bbox="240 891 1370 1003"> <tr> <td>b</td> <td>0</td> <td>1</td> <td>3</td> <td>4</td> </tr> <tr> <td>$P(B=b)$</td> <td>$\frac{11}{81}$</td> <td>$\frac{10}{81}$</td> <td>$\frac{28}{81}$</td> <td>$\frac{32}{81}$</td> </tr> </table>	b	0	1	3	4	$P(B=b)$	$\frac{11}{81}$	$\frac{10}{81}$	$\frac{28}{81}$	$\frac{32}{81}$	<p>B1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>B1</p> <p>A1</p> <p>(10)</p> <p>Total 10</p>
b	0	1	3	4							
$P(B=b)$	$\frac{11}{81}$	$\frac{10}{81}$	$\frac{28}{81}$	$\frac{32}{81}$							
Notes											
<p>(i)</p> <p>1st M1: for identifying the 3 possible samples</p> <p>2nd M1: for $p \times p \times q \times 3$ <u>or</u> $p \times q \times q \times 3$ where p and q are probabilities with $(p + q) < 1$</p> <p>3rd M1: for $p \times q \times r \times 6$ where p, q and r are probabilities with $(p + q + r) = 1$</p> <p>A1: for $\frac{32}{81}$ <u>or</u> awrt 0.395 [Calc: 0.3950617...]</p> <p>(ii)</p> <p>1st M1: for $p^3 + q^3 + r^3$ (for their p, q and r)</p> <p>2nd M1: for $3 \times p \times (q)^2 + 3 \times q \times (p)^2$ <u>or</u> $3 \times q \times (r)^2 + 3 \times r \times (q)^2$ (for their p, q and r)</p> <p>3rd M1: for use of all probabilities of $P(B=b)$ adding to 1 [Must have 3, 4 or 5 values for b]</p> <p>B1: for ranges 0, 1, 3 and 4 with none omitted and no extras. Allow extras if assigned probability of 0</p> <p>A1: for a fully correct probability distribution.</p> <p>SC A0 in (i) If A0 scored in (i) <u>and</u> all other marks scored in (ii) <u>and</u> correct prob's for 2 values of b : award A1 in (ii)</p>											

Question Number	Scheme	Marks
<p>5 (a)(i)</p> <p>(ii)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>If $y = 0$ then $1 - (\alpha + \beta y^2) = 0 \quad \therefore \alpha = 1$ *</p> <p>If $y = 5$ then $1 - (\alpha + \beta y^2) = 1$</p> $1 + 25\beta = 0 \quad \therefore \beta = -\frac{1}{25} \quad *$ <p>$F(y) = \frac{1}{25}y^2$ so $f(y) = \frac{dF(y)}{dy} = \frac{2}{25}y$</p> $\therefore [f(y) =] \begin{cases} \frac{2}{25}y & 0 \leq y \leq 5 \\ 0 & \text{otherwise} \end{cases}$ <p>$\left[P\left(R > \frac{11}{5}\right) = P\left(Y > \frac{5}{3}\right) = 1 - \frac{1}{25} \times \left(\frac{5}{3}\right)^2 = \right] \frac{8}{9}$ oe</p> <p>$\frac{3d - \frac{11}{5}}{3d - d} = \frac{8}{9}$ oe or $\frac{\frac{11}{5} - d}{3d - d} = \frac{1}{9}$ oe</p> $d = \frac{9}{5}$ oe <p>$P\left(Y < \frac{11}{5}\right) = \frac{121}{625}$ or 0.1936</p> <p>[Let G = the number of spins with distance < 2.2 m]</p> <p>[$P(G \geq 5) =$</p> $\left(\frac{1}{9}\right)^3 \times \left(\frac{121}{625}\right)^3 + 3 \times \left(\frac{1}{9}\right)^2 \times \left(\frac{8}{9}\right) \times \left(\frac{121}{625}\right)^3 + 3 \times \left(\frac{1}{9}\right) \times \left(\frac{121}{625}\right)^2 \times \left(\frac{504}{625}\right)$ <p>= 0.000 373226 awrt 0.000 373</p>	<p>B1cso</p> <p>B1cso</p> <p>(2)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>B1</p> <p>M1, M1</p> <p>A1</p> <p>(4)</p> <p>Total 11</p>
Notes		
<p>(a) (i)</p> <p>(ii)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>B1: for stating or using the fact that when $y = 0$ then $\alpha + \beta y^2 = 1$</p> <p>B1: for stating or using that when $y = 5$ then $\alpha + \beta y^2 = 0$ and setting up the equation leading to $\beta = -\frac{1}{25}$</p> <p>M1: for differentiating. Implied by $\pm \frac{2}{25}y$ can fit their value of β</p> <p>A1: for a fully correct $f(y)$ defined for the whole range.</p> <p>B1: for using $F(y)$ and $\frac{5}{3}$ to find $P(Y > \frac{5}{3})$. Allow $\frac{8}{9}$ or any exact equivalent.</p> <p>M1: for LHS = p where $0 < p < 1$</p> <p>A1: for $\frac{9}{5}$ or any exact equivalent e.g. 1.8</p> <p>B1: for $\frac{121}{625}$ or awrt 0.194 This mark could be implied by a correct answer.</p> <p>1st M1: for $p^3q^3 + np^2(1-p)q^3 + np^3q^2(1-q)$ where p and q are probabilities and n is an integer > 0</p> <p>2nd M1: for $p^3q^3 + 3p^2(1-p)q^3 + 3p^3q^2(1-q)$ where p and q are probabilities.</p> <p>A1: for awrt 0.000 373</p>	

Question Number	Scheme	Marks
6. (i)	$z = 1.25$ $\frac{187.5 - \mu}{\sigma} = 1.25$ $187.5 - \mu = 1.25\sigma$ $\mu = 225p$ $\sigma = \sqrt{225p(1-p)}$ $(187.5 - 225p)^2 = (1.25)^2 \times 225p(1-p) \text{ or } (150 - 180p)^2 = 225p(1-p) \text{ (o.e.)}$ $\text{e.g. } 900(5 - 6p)^2 = 225(p - p^2) \Rightarrow 4(25 - 60p + 36p^2) = p - p^2$ <p>Leading to $145p^2 - 241p + 100 = 0^*$</p>	B1 M1 M1 A1 M1 M1 M1 A1* M1 A1 (10)
	Notes	Total 10
(i)	<p>B1: for 1.25 or better (calculator gives: 1.25027...)</p> <p>1st M1: for attempting to use a continuity correction i.e. for sight of 188 ± 0.5</p> <p>2nd M1: for standardising using μ and σ or np and $\sqrt{np(1-p)}$ (Condone letter n or any integer > 0)</p> <p>1st A1: for a correct equation with compatible signs, allow 1.250... If using a value for n it must be 225</p> <p>3rd M1: for $\mu = 225p$ seen at any stage in the working.</p> <p>4th M1: for $\sigma = \sqrt{225p(1-p)}$ seen at any stage in the working.</p> <p>5th M1: for squaring to get a quadratic equation in p</p> <p>2nd A1*: dep on all previous Ms and use of 1.25 for at least 1 correct intermediate step from a correct quadratic equation e.g one of those in scheme for 5th M1</p>	
(ii)	<p>M1: for solving the quadratic correctly—leading to $p = \dots$ or implied by 0.8 or awrt 0.862</p> <p>A1: for 0.8 <u>and</u> a correct reason to eliminate 0.862</p>	

