

GCE Examinations  
Advanced Subsidiary

## **Core Mathematics C3**

Paper F

Time: 1 hour 30 minutes

### *Instructions and Information*

---

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration.

Full marks may be obtained for answers to ALL questions.

Mathematical formulae and statistical tables are available.

This paper has eight questions.

### *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner.  
Answers without working may gain no credit.



*Written by Shaun Armstrong*

© Solomon Press

*These sheets may be copied for use solely by the purchaser's institute.*

1. Solve the equation

$$3 \operatorname{cosec} \theta^\circ + 8 \cos \theta^\circ = 0$$

for  $\theta$  in the interval  $0 \leq \theta \leq 180$ , giving your answers to 1 decimal place. **(6)**

---

2. The functions  $f$  and  $g$  are defined by

$$f : x \rightarrow 1 - ax, \quad x \in \mathbb{R},$$

$$g : x \rightarrow x^2 + 2ax + 2, \quad x \in \mathbb{R},$$

where  $a$  is a constant.

- (a) Find the range of  $g$  in terms of  $a$ . **(3)**

Given that  $gf(3) = 7$ ,

- (b) find the two possible values of  $a$ . **(4)**
- 

3. (a) Solve the equation

$$\ln(3x + 1) = 2$$

giving your answer in terms of  $e$ . **(3)**

- (b) Prove, by counter-example, that the statement

$$“\ln(3x^2 + 5x + 3) \geq 0 \text{ for all real values of } x”$$

is false. **(5)**

---

4. A curve has the equation  $x = y\sqrt{1-2y}$ .

- (a) Show that

$$\frac{dy}{dx} = \frac{\sqrt{1-2y}}{1-3y}. \quad \mathbf{(5)}$$

The point  $A$  on the curve has  $y$ -coordinate  $-1$ .

- (b) Show that the equation of tangent to the curve at  $A$  can be written in the form


$$\sqrt{3}x + py + q = 0$$

where  $p$  and  $q$  are integers to be found. **(3)**

---

5. (a) Sketch the graph of  $y = 2 + \sec\left(x - \frac{\pi}{6}\right)$  for  $x$  in the interval  $0 \leq x \leq 2\pi$ .  
 Show on your sketch the coordinates of any turning points and the equations of any asymptotes. (5)
- (b) Find, in terms of  $\pi$ , the  $x$ -coordinates of the points where the graph crosses the  $x$ -axis. (5)
- 

6.



**Figure 1**

Figure 1 shows the curve  $y = f(x)$  which has a minimum point at  $\left(-\frac{3}{2}, 0\right)$ , a maximum point at  $(3, 6)$  and crosses the  $y$ -axis at  $(0, 4)$ .

Sketch each of the following graphs on separate diagrams. In each case, show the coordinates of any turning points and of any points where the graph meets the coordinate axes.

- (a)  $y = f(|x|)$  (3)
- (b)  $y = 2 + f(x + 3)$  (4)
- (c)  $y = \frac{1}{2}f(-x)$  (4)
- 

**Turn over**

7. 
$$f(x) = 1 + \frac{4x}{2x-5} - \frac{15}{2x^2 - 7x + 5}, \quad x \in \mathbb{R}, \quad x < 1.$$

(a) Show that

$$f(x) = \frac{3x+2}{x-1}. \quad (5)$$

(b) Find an expression for the inverse function  $f^{-1}(x)$  and state its domain. (5)

(c) Solve the equation  $f(x) = 2$ . (2)

---

8. A curve has the equation  $y = x^2 - \sqrt{4 + \ln x}$ .

(a) Show that the tangent to the curve at the point where  $x = 1$  has the equation

$$7x - 4y = 11. \quad (5)$$

The curve has a stationary point with  $x$ -coordinate  $\alpha$ .

(b) Show that  $0.3 < \alpha < 0.4$  (3)

(c) Show that  $\alpha$  is a solution of the equation

$$x = \frac{1}{2}(4 + \ln x)^{-\frac{1}{4}}. \quad (2)$$

(d) Use the iteration formula

$$x_{n+1} = \frac{1}{2}(4 + \ln x_n)^{-\frac{1}{4}},$$

with  $x_0 = 0.35$ , to find  $x_1, x_2, x_3$  and  $x_4$ , giving your answers to 5 decimal places. (3)

---

**END**