

GCE Examinations
Advanced Subsidiary

Core Mathematics C3

Paper A

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.





Written by Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

C3 Paper A – Marking Guide

1. $\frac{dx}{dy} = 2 \sec y \times \sec y \tan y + \sec^2 y = \sec^2 y(2 \tan y + 1) = \frac{2 \tan y + 1}{\cos^2 y}$ M1 A1
 $\frac{dy}{dx} = 1 \div \frac{dx}{dy} = \frac{\cos^2 y}{2 \tan y + 1}$ M1 A1 (4)
-
2. (a) $= f\left(\frac{1}{2}\right) = -\frac{5}{2}$ M1 A1
 (b) $gf(x) = \frac{2}{(3x-4)+3} = \frac{2}{3x-1}$ M1 A1
 $\therefore \frac{2}{3x-1} = 6$
 $2 = 6(3x - 1)$ M1
 $x = \frac{4}{9}$ A1 (6)
-
3. $e^{2y} - x + 2 = 0 \Rightarrow e^{2y} = x - 2$
 $2y = \ln(x - 2)$ M1
 sub. $\Rightarrow \ln(x + 3) - \ln(x - 2) - 1 = 0$ A1
 $\ln \frac{x+3}{x-2} = 1$ M1
 $\frac{x+3}{x-2} = e$ A1
 $x + 3 = e(x - 2)$ M1
 $3 + 2e = x(e - 1)$ M1
 $x = \frac{2e+3}{e-1} = 4.91$ (2dp), $y = \frac{1}{2} \ln\left(\frac{2e+3}{e-1} - 2\right) = 0.53$ (2dp) A2 (8)
-
4. (a) $\frac{d}{dx}(\tan x) = \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right)$
 $= \frac{\cos x \times \cos x - \sin x \times (-\sin x)}{\cos^2 x}$ M1 A1
 $= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x$ M1 A1
 (b) $\frac{dy}{dx} = 2 \times \tan x + 2x \times \sec^2 x = 2 \tan x + 2x \sec^2 x$ M1 A1
 $x = \frac{\pi}{4}, y = \frac{\pi}{2}, \text{grad} = 2 + \pi$ B1
 $\therefore y - \frac{\pi}{2} = (2 + \pi)(x - \frac{\pi}{4})$ M1
 at P, $x = 0$
 $\therefore y = \frac{\pi}{2} - \frac{\pi}{4}(2 + \pi) = -\frac{1}{4}\pi^2$ M1 A1 (10)
-
5. (a) $3 \cos x + \sin x = R \cos x \cos \alpha + R \sin x \sin \alpha$
 $R \cos \alpha = 3, R \sin \alpha = 1$
 $\therefore R = \sqrt{3^2 + 1^2} = \sqrt{10}$ M1 A1
 $\tan \alpha = \frac{1}{3}, \alpha = 18.4$ (3sf) M1 A1
 $\therefore 3 \cos x^\circ + \sin x^\circ = \sqrt{10} \cos(x - 18.4)^\circ$
 (b) $6 \cos^2 x + 2 \sin x \cos x = 0$ M1
 $2 \cos x(3 \cos x + \sin x) = 0$ M1
 $\cos x = 0$ or $3 \cos x + \sin x = \sqrt{10} \cos(x - 18.4) = 0$ A1
 $x = 90, 270$ or $x - 18.4 = 90, 270$ B1 M1
 $x = 90, 108.4$ (1dp), $270, 288.4$ (1dp) A1 (10)

6. (a) (i)  (ii) 

(b) $y = 0 \Rightarrow 2x - 1 = 0 \Rightarrow x = \frac{1}{2} \therefore p = \frac{1}{2}$ M1 A1
 $x = 0 \Rightarrow y = 1 \therefore q = 1$ B1

(c) $y = \frac{2x-1}{x-1}, \quad y(x-1) = 2x-1$ M1
 $x(y-2) = y-1, \quad x = \frac{y-1}{y-2}$ M1
 $\therefore f^{-1}(x) = \frac{x-1}{x-2}$ A1 (12)


7. (a) (i) LHS = $\sin x \cos 30 + \cos x \sin 30 + \sin x \cos 30 - \cos x \sin 30$ M1 A1
 $= 2 \sin x \cos 30 = \sqrt{3} \sin x \quad [a = \sqrt{3}]$ A1

(ii) let $x = 45, \sin 75 + \sin 15 = \sqrt{3} \sin 45 = \sqrt{3} \times \frac{1}{\sqrt{2}} = \frac{1}{2}\sqrt{6}$ M2 A1

(b) $2(\operatorname{cosec}^2 y - 1) + 5 \operatorname{cosec} y + \operatorname{cosec}^2 y = 0$ M1
 $3 \operatorname{cosec}^2 y + 5 \operatorname{cosec} y - 2 = 0, \quad (3 \operatorname{cosec} y - 1)(\operatorname{cosec} y + 2) = 0$ M1
 $\operatorname{cosec} y = -2$ or $\frac{1}{3}$ (no solutions) A1
 $\sin y = -\frac{1}{2}$
 $y = 180 + 30, 360 - 30$ B1 M1
 $y = 210, 330$ A1 (12)

8. (a)
$$\begin{array}{r} x^2 + 0x + 1 \\ x^2 + x - 6 \overline{) x^4 + x^3 - 5x^2 + 0x - 9} \\ \underline{x^4 + x^3 - 6x^2} \\ x^2 + 0x - 9 \\ \underline{x^2 + x - 6} \\ -x - 3 \end{array}$$
 M1 A1

$\therefore f(x) = x^2 + 1 + \frac{-x-3}{x^2+x-6}$ A1
 $= x^2 + 1 - \frac{x+3}{(x-2)(x+3)} = x^2 + 1 - \frac{1}{x-2} \quad [A = 1, B = -1, C = -2]$ M1 A1

(b)  B2

$f(x) = 0 \Rightarrow x^2 + 1 = \frac{1}{x-2},$ graphs intersect once \therefore exactly one real root B1

(c) e.g. $x_0 = 3, \quad x_1 = 2.1, \quad x_2 = 2.1848, \quad x_3 = 2.1732, \quad x_4 = 2.1747$ M1 A1
 \therefore root = 2.17 (3sf) A1
 $f(2.165) = -0.37, \quad f(2.175) = 0.016$ M1
 sign change, $f(x)$ continuous \therefore root A1 (13)

Total (75)

Performance Record – C3 Paper A

Question no.	1	2	3	4	5	6	7	8	Total
Topic(s)	differentiation	functions	exponentials and logarithms	differentiation	trigonometry	functions	trigonometry	rational expressions, numerical methods	
Marks	4	6	8	10	10	12	12	13	75
Student									