	<b>EXAMINATION PAPER 1</b>	Matching the syllabus written by	
	Calculators Allowed	EDEXCEL Curriculum 2004+	
W	here appropriate give your answers to 3 s.f.		
	© ZigZag Education 2004	Core Mathematics – C4	
	Time Allowe	ed:-1 hour 30 minutes	
der	. 2		
uy/c	$dx = \sin^2 x$		
Wh	en $x = 0.1$ , $y = 0.2$ . [Note: x and y are]	given in radians.]	
Fine	d y = $f(x)$ and evaluate any constants.		
		$\frac{1}{2}$	
a)	Expand the first four terms in the bin	iomial expansion of $(1 - x)^{-2}$ , $ x  < 1$ , in ascending	
1 \	powers of x.		
<b>b</b> )	Letting $x = 10$ in the above expansion	on, <u>nence</u> find VII correct to 5 significant figures.	
<u> </u>			
a)	Calculate J $xe^{-1} dx$		
b)	Hence or otherwise calculate $\int x^2 e^x dx$	x	
<u> </u>	A has position vestor : : : : ! ! or d D	has position vootor i + ai + ly where a is a constant	
aj	If $\Omega A$ and $\Omega R$ are perpendicular then	find $c$	
	n on and ob are perpendicular then		
L <sub>1</sub> h	has equation $\mathbf{r} = (\mathbf{i} + 2\mathbf{j} + \mathbf{k}) + \mu(3\mathbf{i} + 2\mathbf{j} - \mathbf{k})$	+ 3 <b>k</b> ), where $\mu$ is a scalar.	
$L_2 h$	has equation $\mathbf{r} = (\mathbf{i} + \mathbf{k}) + \lambda(\mathbf{i} + \mathbf{i} + \mathbf{k})$ .	here $\lambda$ is a scalar.	
b)	Calculate where the 2 lines cross.		
c)	Calculate the acute angle between the	e lines. Give your answer in degrees.	
f(x)	$=\frac{x+1}{(x-1)(x+2)}$		
	$(\lambda - 1)(\lambda + 2)$		
a)	Express $f(x)$ as partial fractions		
b)	Calculate $f'(2)$ .	$\mathbf{A} = \mathbf{A} + $	
c)	Calculate the area A, which is bound lines $x = 4$ and $x = 5$ . Leave your and	eq by the curve $y = f(x)$ and the x-axis and the	
	x = 4 and $x = 5$ . Leave your and	swei exactiy.	
$x^{2} +$	$-y^2 + 2x + 4y + 1 = 0$		
a)	Determine the coordinates of the poi	nts of this circle when $x = 1/2$	
c)	Calculate $dy/a$ , in terms of r and y	-	
-) -)	Coloulate the gradients of the $2$ terms	ants to the surve when $x = 1/2$	
u)	Calculate the gradients of the 2 tange	This to the curve when $x - \frac{1}{2}$	
e)	Calculate the equation of the tangent	with a positive gradient.	
The	e sketch shows the finite region bounded	1 by	
the	curve $y = x \sin^{\frac{1}{2}} x$ and the x-axis, $0 \le x \le x$	$\leq \pi$ .	
This	s region is rotated $2\pi$ radians about the	x-axis,	
Fine	d the exact value of volume of the gene	rated. 0 $\pi \times x$	

[75]

	EXAMINATION PAPER 2	Matching the syllabus written by	
	Calculators Allowed	EDEXCEL Curriculum 2004+	
W	here appropriate give your answers to 3 s.f.		
	© ZigZag Education 2004	Core Mathematics – C4	
	Time Allowe	ed:-1 hour 30 minutes	
$\mathbf{v} =$	$3e^{x}$		
y a)	Sketch this curve stating where the c	curve crosses the v-axis	[2]
b)	Find the area bounded by the curve,	the x-axis, the y-axis and the line $x = \ln 10$ .	[4]
a)	Calculate $\int_{1}^{4} \frac{1}{1+2x} dx$ . <i>Give your answe</i>	er exactly.	[3]
b)	Estimate $\int_{1}^{4} \frac{x}{1+2x} dx$ by using the trapez	zium rule with 3 intervals.	[5]
a)	Expand the first four terms in the bin	nomial expansion of $(1 + x)^{-3}$ , $ x  < 1$ , in ascending	[5]
b)	Hence expand the first four terms in powers of $x$ .	the binomial expansion of $(4 + 4x)^{-3}$ , $ x  < 1$ , in asc	ending [2]
A pa a)	air of parametric equations are $x = t + 1$ Formulate an equivalent Cartesian ec	and $y = t^2 + 2$ . quation.	[3]
A cub)	urve's parametric equations are $x = \sin^3$ Find $\frac{dy}{dx}$ as a function of t.	t and $y = t + cost$	[4]
c)	i) Find the gradient of the curve y	when $t = \pi/A$ .	[2]
,	ii) Find the equation of the norma	l to the curve when $t = \pi/4$ .	[5]
$\overline{\mathbf{f}(x)}$	$=$ $\frac{7}{(1+1)^2}$		
	(x+6)(2x+1)		
a) b)	Express $f(x)$ as partial fractions Calculate $f'(x)$ and hence find $f'(1)$ le	aving your answer as a fraction.	[7] [4]
A ha	as position vector $-\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ and <i>B</i> has	position vector $\mathbf{i} + 2\mathbf{j} - 5\mathbf{k}$ .	
a)	Calculate the position vector $\overrightarrow{AB}$ .		[3]
h)	Calculate the unit vector of $\overrightarrow{AB}$		[2]
c)	Calculate the vector equation of the s	straight line that passes through $A$ and $B$	[ <del>2</del> ]
A se	econd straight line also passes through	<i>B</i> and through the origin 0	[-]
d)	Find the acute angle between the line	e in c) and this 2 <sup>nd</sup> line. <i>Give your answer in degre</i>	es. <b>[5]</b>
a)	Calculate $\int_{1}^{\frac{\pi}{4}} \sin x \cos x  dx$		[4]
b)	Calculate $\int_{0}^{1} e^{x} \sin x  dx$ by using integr	ration by parts twice.	

A curve is defined implicitly as  $x \sin x = y \cos y$ .

[5]

	<b>EXAMINATION PAPER 3</b>	Matching the syllabus written by	
	Calculators Allowed	EDEXCEL Curriculum 2004+	
WI	ere appropriate give your answers to 3	S.f. Core Mathematics – C4	
	Time A	llowed:-1 hour 30 minutes	
	×	nowed. I nour so minutes	
y = 1	$3 - 2e^{x}$	the sum is an again the wavis and the wavis	
a) b)	Sketch this curve, stating where Find the area bounded by the cu	the curve crosses the x-axis and the y-axis. urve, the x-axis and the lines $r = 2$ and $r = 3$	
0)	Find the area bounded by the cu	inve, the x-axis and the times $x - 2$ and $x - 3$ .	
a)	Expand the first four terms in th	the binomial expansion of $(1 - 4x)^{1/2}$ , $ x  < 1/4$ , in ascend	in
1-)	powers of x.	amost to 5 significant figures	
D)	Let $x = 10$ and hence find v6 c	orrect to 5 significant figures.	
A cu	urve's parametric equations are x =	$=\sin^2 t$ and $y = 1 + \cos t$	
a)	Find $\frac{dy}{dx}$ as a function of t.		
b)	Find A where, the gradient of th	e curve is $A\sqrt{2}$ , when $t = \pi/4$ .	
c)	Eliminate t from the parametric	equations and therefore find an equivalent Cartesian	
,	equation of the curve.	-	
 a)	Calculate $\int \sin x \cos^3 x  dx$ using t	the substitution $u = \cos x$ . Show your working to gain full marks	
u) b)	Finding A and B; write 2sin 6x c	$\cos 5x$ in the form $\sin Ax + \sin Bx$	
c)	Using your result from b) or oth	herwise integrate $\int \sin 6x \cos 5x  dx$	
a) b)	$\overline{(x-1)^2(x+2)}$ Express f(x) as partial fractions Calculate f'(2).	bounded by the curve $y = f(r)$ the r-axis and the lines $r = r^2$	_
	and $x = 5$ . Leave your answer e	exactly.	
a)	Calculate $\int_{1}^{4} e^{x} dx$		
b)	i) Estimate $\int_{1}^{4} e^{-x} dx$ by using t	the trapezium rule with 3 intervals.	
	ii) Explain with the aid of a c estimation of the exact int trapezium rule.	diagram explain whether your answer is an over or unde egral and give a method of improving the accuracy usin	r
c) W	What is the relationship between th	e graphs of $f(x)$ and $g(x)$ , where $f(x) = e^x$ and $g(x) = e^{-x}f(x)$	)
A ha	as position vector $-\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ and $B$	B has position vector $\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ .	
a)	Calculate the position vector $\overrightarrow{AB}$		
b)	Calculate the unit vector of $\overrightarrow{OA}$ .	(O the origin).	
c)	Calculate the distance between A	A and B.	
$L_1$ h	as equation $\mathbf{r} = (\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}) + \mu(\mathbf{i})$	$3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ ), where $\mu$ is a scalar quantity.	
L <sub>2</sub> h	as equation $\mathbf{r} = (3\mathbf{i} + 2\mathbf{j} + \mathbf{k}) + \lambda(\mathbf{i}$	$\mathbf{i} + 2\mathbf{j} + A\mathbf{k}$ ), where A is a constant and $\lambda$ is a scalar quan	ti
d)	Given that the 2 lines intersect f	ind the value of A.	
e)	Calculate the acute angle betwee	en the lines. <i>Give your answer in degrees</i> .	

[75]

	EXAMINATION PAPER 4	Matching the syllabus written by	
	Calculators Allowed	EDEXCEL Curriculum 2004+	
WI	here appropriate give your answers to 3 s.f.	Coro Mathamatica C4	
	© ZigZag Education 2004	d: 1 hour 20 minutes	
A n	air of parametric equations are $r = 3cor$	$s_{2t}$ and $v = 3sin_{2t}$	
For	mulate an equivalent Cartesian equation	and describe the nature of this curve.	
dy/	$d_x = x + \sin x$ and $x = 0.1$ when $y = 0.2$ .	The value of x is given in radians.	
Finc	d y = f(x) and evaluate any constants.		
a)	Expand the first four terms in the bin powers of $x$ .	omial expansion of $(1 - 2x)^{1/2}$ , $ x  < 1/2$ , in ascende	lir
b)	By letting $x = 10^{-2}$ in the above expanding figures.	sion, find an approximation of $\sqrt{2}$ correct to 5 sig	ŗn
a)	Calculate $\int x \sin x  dx$		
b)	Hence or otherwise calculate $\int x^2 \cos x$	c dx	
A cu	urve's parametric equations are $x = \cos \theta$	St and $y = \cos^2 t$	
a)	Find $dy/dx$ as a function of t.		
b)	Find the gradient of the curve when t	$=\pi/_4$	
c)	Find the equation of the normal to the	e curve when $t = \frac{\pi}{4}$ .	
a)	$\int \frac{x^3}{x^4+1} dx$		
b)	Calculate $\int_{1}^{5} x^{\frac{1}{2}} + e^{x} dx$ . <i>Give your answe</i>	er to 1 decimal place.	
c)	Estimate $\int_{1}^{5} \frac{1}{x^{\frac{1}{2}} + e^{x}} dx$ by using the tra	apezium rule with 4 intervals. $x \neq y$	
	A sketch of $y = \frac{1}{x^{\frac{1}{2}} + e^x}$ is shown, sket	tch the area	
	represented by this integral and state trapezium rule gives an over or under of the integral Shade onto your sket	ch this	
	over or under estimate between $x = 1$	and $x = 2$ .	
$\frac{1}{A}$ ha	as position vector $-\mathbf{i} + \mathbf{j} + \mathbf{k}$		-
C ha	as position vector $c\mathbf{i} + \mathbf{j} + 2\mathbf{k}$ as position vector $c\mathbf{i} + \mathbf{j} + 5\mathbf{k}$ where c is	a constant. $\rightarrow \rightarrow \rightarrow$	
a) ( Givi	Calculate the acute angle between the period of the period	osition vectors $OA$ and $OB$ , where $O$ is the origin.	
b) (	Calculate the length between point $A$ an	d <i>B</i> .	
	Given that the position vectors $\overrightarrow{AC}$ and	$\overrightarrow{AB}$ are perpendicular, then find <i>c</i> .	
c) (	<b>a. 1 . . . . . . . . . .</b>		

a) Express f(x) in partial fractions.

[3]

b)	Calculate $f'(x)$ .	[2]
c)	Expand $f(x)$ as a series of ascending powers of x, up to and including the term involving	$x^2$ .
Clear	ly indicate in your answer to c) the range of values for which the expansion is valid.	[6]

[75]

	Matching the syllabus written by	<b>EXAMINATION PAPER 5</b>	
	EDEXCEL Curriculum 2004+	Calculators Allowed	
		<i>here appropriate give your answers to 3 s.f.</i>	Whe
	Core Mathematics – C4	© ZigZag Education 2004	
	:d:-1 hour 30 minutes	Time Allow	
50 <sup>0</sup> ał I	<i>x</i> -axis and the lines $x = 2$ and $x = 3$ is rotated 36 ly, and using integration, the volume formed.	e area bounded by the curve $y = e^{x/2}$ the <i>x</i> -axis. Find, leaving your answer exact	The an he <i>x</i> -a
		$d_x = x + e^x$ and $x = 1$ when $y = 2$ .	ly/dr
onstan	r answer in the form $y = f(x)$ & evaluate any co	ve the differential equation & write you	Solve
	and $y = \sin 3t$ .	curve's parametric equations are $x = sint$	A cur
I		Find $\frac{dy}{dx}$ as a function of t.	ı)
I	$=\pi/4.$	Find the gradient of the curve when	))
l	e curve when $t = \pi/4$ .	Find the equation of the tangent to the	:)
	$x \equiv \frac{1}{2} [3\sin 6x - \sin 2x] \text{ find,}$	Given that $2\sin 2x\cos 4x + \cos 2x\sin^2 4x$	a)
[		$\int (4\sin 2x\cos 4x + 2\cos 2x\sin 4x)  \mathrm{d}x$	
	$\int_{0}^{\frac{\pi}{2}} 16\cos x \sin^{4} x  dx$	<b>Using</b> the substitution $u = \sin x$ , find	))
I	0 <i>S</i> .	Show your working to gain full mark	
	his a constant	as position vector $\mathbf{i} + \mathbf{j} + \mathbf{k}$	4 has
	) is a constant.	has position vector $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$	C has
	a 14 units.	e distance between the points A and B i	The d
[		Find the possible values of <i>b</i> .	ı)
origin	e position vectors OA and OC, where O is the o	Calculate the acute angle between th	))
ĺ		ve your answer in degrees.	Give y
	at line that passes through A and C the point $17i$ , $31i$ , $dk$ and find d	Find the vector equation of the straig	:)
I	The point $1/1 - 31j - ak$ and find $a$ .	and snow that the fine passes through	
	st and $y = 3 \sin t$ .	pair of parametric equations are, $x = 2cc$	A pair
I	an equation of this curve.	1) Formulate an equivalent Cartes	1)
	nd a formula for $\frac{dy}{dx}$ in terms of x and y.	ii) Using implicit differentiation f	
I	terms of t.	i) Using the chain rule find $\frac{dy}{dx}$ in	))
	f <i>x</i> and y and show that this is the same as	ii) Hence determine $\frac{dy}{dr}$ in terms of	

 $\frac{2}{(2x+1)(x+2)}$ 

a)

Express f(x) as partial fractions. [6] Expand f(x) as a series of ascending powers of *x*, up to and including the term involving  $x^2$ . b)