

# GCE Examinations

# Further Pure Mathematics Module FP2

Advanced Subsidiary / Advanced Level

## Paper A

Time: 1 hour 30 minutes

### *Instructions and Information*

---

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 8 questions.

### *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner.  
Answers without working will gain no credit.



*Written by Rosemary Smith & Shaun Armstrong*

© Solomon Press

*These sheets may be copied for use solely by the purchaser's institute.*

1. A curve has the equation

$$y = x + 2x^2 + 5x^3.$$

Show that the radius of curvature of the curve at the origin is  $\frac{1}{\sqrt{2}}$ . **(5 marks)**

---

2. Show that

$$\int_0^{\ln 2} x \operatorname{sech}^2 x \, dx = \frac{3}{5} \ln 2 - \ln\left(\frac{5}{4}\right). \quad \text{(8 marks)}$$

---

3. (a) Prove that

$$\frac{d}{dx} (\arcsin 2x) = \frac{2}{\sqrt{1-4x^2}}. \quad \text{(3 marks)}$$

Given that

$$f(x) = 2x \arcsin 2x + \sqrt{1-4x^2},$$

- (b) show that

$$f''(x)[f(x) - xf'(x)] = 4. \quad \text{(6 marks)}$$

---

4.




Fig. 1

The parametric coordinates of the curve  $C$  shown in Figure 1 are

$$x = t^2, \quad y = t - \frac{1}{3}t^3, \quad 0 \leq t \leq a.$$

The curve  $C$  meets the  $x$ -axis at the point  $A$  where  $t = a$ .

(a) Find the value of  $a$ . **(2 marks)**

The curve  $C$  is rotated through  $2\pi$  about  $Ox$ .

(b) Find the surface area of the solid generated. **(8 marks)**

---

5. (a) Using the definitions of  $\cosh x$  and  $\sinh x$  in terms of  $e^x$  and  $e^{-x}$ , prove that

$$\cosh 2x = 2 \cosh^2 x - 1. \quad \text{(3 marks)}$$

(b) Solve the equation

$$2 \cosh 2x = 13 \cosh x - 12,$$

giving your answers in terms of natural logarithms. **(7 marks)**

---

*Turn over*

6.  $x^2 - 10x + 41 \equiv (x + a)^2 + b.$

(a) Find the values of the constants  $a$  and  $b$ . **(2 marks)**

(b) Show that

$$\int_5^9 \frac{x}{\sqrt{x^2 - 10x + 41}} dx = p(\sqrt{2} - 1) + q \ln r,$$

stating your values of  $p$ ,  $q$  and  $r$ . **(8 marks)**

---

7.  $I_n = \int_0^{\frac{\pi}{2}} x^n \cos x dx, \quad n \geq 0.$

(a) Prove that

$$I_n = \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2} \quad n \geq 2. \quad \text{(5 marks)}$$

(b) Hence find the value of  $I_4$ , giving your answer in terms of  $\pi$ . **(6 marks)**

---

8. The rectangular hyperbola  $C$  has equation  $xy = c^2$ , where  $c$  is a positive constant.

(a) Show that an equation of the tangent to  $C$  at the point  $P\left(cp, \frac{c}{p}\right)$  is

$$x + yp^2 = 2cp. \quad \text{(4 marks)}$$

The tangent to  $C$  at  $P$  meets the  $x$ -axis at the point  $X$ .

The point  $Q$  on  $C$  has coordinates  $\left(cq, \frac{c}{q}\right)$ ,  $q \neq p$  such that  $QX$  is parallel to the  $y$ -axis.

(b) Show that  $q = 2p$ . **(3 marks)**

$M$  is the mid-point of  $PQ$ .

(c) Find, in Cartesian form, an equation of the locus of  $M$  as  $p$  varies. **(5 marks)**

---

**END**