

**ADVANCED GCE**  
**MATHEMATICS**  
Core Mathematics 3

**4723**

Candidates answer on the Answer Booklet

**OCR Supplied Materials:**

- 8 page Answer Booklet
- List of Formulae (MF1)

**Other Materials Required:**

None

**Thursday 15 January 2009**  
**Morning**

**Duration:** 1 hour 30 minutes



**INSTRUCTIONS TO CANDIDATES**

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- This document consists of **4** pages. Any blank pages are indicated.

1 Find

(i)  $\int 8e^{-2x} dx,$

(ii)  $\int (4x + 5)^6 dx.$

[5]

2 (i) Use Simpson's rule with four strips to find an approximation to

$$\int_4^{12} \ln x dx,$$

giving your answer correct to 2 decimal places.

[4]

(ii) Deduce an approximation to  $\int_4^{12} \ln(x^{10}) dx.$

[1]

3 (i) Express  $2 \tan^2 \theta - \frac{1}{\cos \theta}$  in terms of  $\sec \theta.$

[3]

(ii) Hence solve, for  $0^\circ < \theta < 360^\circ,$  the equation

$$2 \tan^2 \theta - \frac{1}{\cos \theta} = 4.$$

[4]

4 For each of the following curves, find  $\frac{dy}{dx}$  and determine the exact  $x$ -coordinate of the stationary point:

(i)  $y = (4x^2 + 1)^5,$

[3]

(ii)  $y = \frac{x^2}{\ln x}.$

[4]

5 The mass,  $M$  grams, of a certain substance is increasing exponentially so that, at time  $t$  hours, the mass is given by

$$M = 40e^{kt},$$

where  $k$  is a constant. The following table shows certain values of  $t$  and  $M.$

$t$	0	21	63
$M$		80	

(i) In either order,

(a) find the values missing from the table,

[3]


(b) determine the value of  $k.$

[2]

(ii) Find the rate at which the mass is increasing when  $t = 21.$

[3]

6



The function  $f$  is defined for all real values of  $x$  by

$$f(x) = \sqrt[3]{\frac{1}{2}x + 2}.$$


The graphs of  $y = f(x)$  and  $y = f^{-1}(x)$  meet at the point  $P$ , and the graph of  $y = f^{-1}(x)$  meets the  $x$ -axis at  $Q$  (see diagram).

- (i) Find an expression for  $f^{-1}(x)$  and determine the  $x$ -coordinate of the point  $Q$ . [3]
- (ii) State how the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$  are related geometrically, and hence show that the  $x$ -coordinate of the point  $P$  is the root of the equation

$$x = \sqrt[3]{\frac{1}{2}x + 2}. \quad [2]$$

- (iii) Use an iterative process, based on the equation  $x = \sqrt[3]{\frac{1}{2}x + 2}$ , to find the  $x$ -coordinate of  $P$ , giving your answer correct to 2 decimal places. [4]


7



The diagram shows the curve  $y = e^{kx} - a$ , where  $k$  and  $a$  are constants.

- (i) Give details of the pair of transformations which transforms the curve  $y = e^x$  to the curve  $y = e^{kx} - a$ . [3]
- (ii) Sketch the curve  $y = |e^{kx} - a|$ . [2]
- (iii) Given that the curve  $y = |e^{kx} - a|$  passes through the points  $(0, 13)$  and  $(\ln 3, 13)$ , find the values of  $k$  and  $a$ . [4]

8



The diagram shows the curve with equation

$$y = \frac{6}{\sqrt{x}} - 3.$$

The point  $P$  has coordinates  $(0, p)$ . The shaded region is bounded by the curve and the lines  $x = 0$ ,  $y = 0$  and  $y = p$ . The shaded region is rotated completely about the  $y$ -axis to form a solid of volume  $V$ .

(i) Show that  $V = 16\pi \left( 1 - \frac{27}{(p+3)^3} \right)$ . [6]

(ii) It is given that  $P$  is moving along the  $y$ -axis in such a way that, at time  $t$ , the variables  $p$  and  $t$  are related by

$$\frac{dp}{dt} = \frac{1}{3}p + 1.$$

Find the value of  $\frac{dV}{dt}$  at the instant when  $p = 9$ . [4]

9 (i) By first expanding  $\cos(2\theta + \theta)$ , prove that

$$\cos 3\theta \equiv 4 \cos^3 \theta - 3 \cos \theta. \quad [4]$$

(ii) Hence prove that

$$\cos 6\theta \equiv 32 \cos^6 \theta - 48 \cos^4 \theta + 18 \cos^2 \theta - 1. \quad [3]$$

(iii) Show that the only solutions of the equation

$$1 + \cos 6\theta = 18 \cos^2 \theta$$

are odd multiples of  $90^\circ$ . [5]