

GCE Examinations
Advanced / Advanced Subsidiary

Core Mathematics C1

Paper E

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for using a valid method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.





Written by Shaun Armstrong

© *Solomon Press*


These sheets may be copied for use solely by the purchaser's institute.

C1 Paper E – Marking Guide

1.	(i) $= \frac{21}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} = 3\sqrt{7}$ (ii) $= \frac{1}{\sqrt[3]{8}} = \frac{1}{2}$	M1 A1 M1 A1	(4)
<hr/>			
2.	(i) $= 1 - 4x$ (ii) $= -6x^{-3}$	M1 A1 M1 A1	(4)
<hr/>			
3.	(a) $= (x - 5)^2 - 25 + 27$ $= (x - 5)^2 + 2$ (b) (i) (ii) 	M1 A2 B3	 (6)
<hr/>			
4.	(i) $y + 5 = 2(x - 4)$ $y = 2x - 13$ (ii) $3x - y = 4 \Rightarrow y = 3x - 4 \therefore \text{grad} = 3$ $\text{grad } l_2 = \frac{-1}{3} = -\frac{1}{3}$ $\therefore y - 0 = -\frac{1}{3}(x - 3) \quad [y = -\frac{1}{3}x + 1]$ (iii) $2x - 13 = -\frac{1}{3}x + 1$ $x = 6$ $\therefore (6, -1)$	M1 A1 M1 A1 A1 M1 A1 A1	 (8)
<hr/>			
5.	(i) real roots $\therefore b^2 - 4ac \geq 0$ $(-k)^2 - [4 \times 4 \times (k - 3)] \geq 0$ $k^2 - 16k + 48 \geq 0$ (ii) $(k - 4)(k - 12) \geq 0$  $k \leq 4$ or $k \geq 12$ (iii) $k = 4$ $4x^2 - 4x + 1 = 0$ $(2x - 1)^2 = 0$ $x = \frac{1}{2}$	M1 A1 M1 M1 A1 B1 M1 A1	 (8)
<hr/>			
6.	(i) $= (-\frac{2+4}{2}, \frac{6-1}{2}) = (1, \frac{5}{2})$ (ii) radius = dist. $(-2, 6)$ to $(1, \frac{5}{2}) = \sqrt{9 + \frac{49}{4}} = \sqrt{\frac{85}{4}}$ $\therefore (x - 1)^2 + (y - \frac{5}{2})^2 = (\sqrt{\frac{85}{4}})^2$ $x^2 - 2x + 1 + y^2 - 5y + \frac{25}{4} = \frac{85}{4}$ $x^2 + y^2 - 2x - 5y - 14 = 0$ (iii) $(2, 7)$, LHS = $4 + 49 - 4 - 35 - 14 = 0 \therefore R$ lies on circle $\angle PRQ = 90^\circ$	M1 A1 M1 A1 M1 A1 A1 B1 B1	 (9)

7. (i) translation by 1 unit in the positive x -direction B2

(ii)



B3

(iii) $\frac{1}{x-1} = 2 + \frac{1}{x}$
 $x = 2x(x-1) + (x-1)$ M1
 $2x^2 - 2x - 1 = 0$ A1
 $x = \frac{2 \pm \sqrt{4+8}}{4}$ M1
 $x = \frac{2 \pm 2\sqrt{3}}{4}$ M1
 $x = \frac{1}{2} \pm \frac{1}{2}\sqrt{3}$ A1 (10)

8. (i) $x(x^2 + 3x - 4) = 0$ M1
 $x(x+4)(x-1) = 0$ M1
 $x = 0$ (at O), -4 , 1
 $\therefore (-4, 0), (1, 0)$ A1

(ii) $\frac{dy}{dx} = 3x^2 + 6x - 4$ M1 A1
 $\text{grad} = -4$ M1
 $\therefore y = -4x$ A1

(iii) $x^3 + 3x^2 - 4x = -4x$ M1
 $x^3 + 3x^2 = 0$
 $x^2(x+3) = 0$ M1
 $x = 0$ (at O), -3 A1
 $\therefore (-3, 12)$ A1 (11)

9. (i) $\frac{dy}{dx} = 3x^{\frac{1}{2}} - 4x^{-\frac{1}{2}}$ M1 A2

(ii) $3x^{\frac{1}{2}} - 4x^{-\frac{1}{2}} = 0$ M1
 $x^{-\frac{1}{2}}(3x - 4) = 0$ M1
 $x = \frac{4}{3}$ A1

(iii) $x = 2 \therefore y = 2(2\sqrt{2}) - 8(\sqrt{2}) = -4\sqrt{2}$ M1 A1
 $\text{grad} = 3\sqrt{2} - \frac{4}{\sqrt{2}} = 3\sqrt{2} - 2\sqrt{2} = \sqrt{2}$ M1 A1
 $\therefore y + 4\sqrt{2} = \sqrt{2}(x - 2)$ M1
 $y = \sqrt{2}x - 6\sqrt{2}$ A1 (12)

Total (72)

