

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education**

MEI STRUCTURED MATHEMATICS

4753/1

Methods for Advanced Mathematics (C3)

Thursday

8 JUNE 2006

Morning

1 hour 30 minutes

Additional materials:

8 page answer booklet

Graph paper

MEI Examination Formulae and Tables (MF2)

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is 72.

This question paper consists of 5 printed pages and 3 blank pages.

Section A (36 marks)

1 Solve the equation $|3x - 2| = x$. [3]

2 Show that $\int_0^{\frac{1}{6}\pi} x \sin 2x \, dx = \frac{3\sqrt{3} - \pi}{24}$. [6]

3 Fig. 3 shows the curve defined by the equation $y = \arcsin(x - 1)$, for $0 \leq x \leq 2$.




Fig. 3

(i) Find x in terms of y , and show that $\frac{dx}{dy} = \cos y$. [3]

(ii) Hence find the exact gradient of the curve at the point where $x = 1.5$. [4]

- 4 Fig. 4 is a diagram of a garden pond.




Fig. 4

The volume $V \text{ m}^3$ of water in the pond when the depth is h metres is given by

$$V = \frac{1}{3}\pi h^2(3 - h).$$

- (i) Find $\frac{dV}{dh}$. [2]

Water is poured into the pond at the rate of 0.02 m^3 per minute.

- (ii) Find the value of $\frac{dh}{dt}$ when $h = 0.4$. [4]

- 5 Positive integers a , b and c are said to form a Pythagorean triple if $a^2 + b^2 = c^2$.

- (i) Given that t is an integer greater than 1, show that $2t$, $t^2 - 1$ and $t^2 + 1$ form a Pythagorean triple. [3]

- (ii) The two smallest integers of a Pythagorean triple are 20 and 21. Find the third integer.

Use this triple to show that not all Pythagorean triples can be expressed in the form $2t$, $t^2 - 1$ and $t^2 + 1$. [3]

- 6 The mass M kg of a radioactive material is modelled by the equation

$$M = M_0 e^{-kt},$$

where M_0 is the initial mass, t is the time in years, and k is a constant which measures the rate of radioactive decay.

- (i) Sketch the graph of M against t . [2]

- (ii) For Carbon 14, $k = 0.000121$. Verify that after 5730 years the mass M has reduced to approximately half the initial mass. [2]

The half-life of a radioactive material is the time taken for its mass to reduce to exactly half the initial mass.

- (iii) Show that, in general, the half-life T is given by $T = \frac{\ln 2}{k}$. [3]

- (iv) Hence find the half-life of Plutonium 239, given that for this material $k = 2.88 \times 10^{-5}$. [1]

Section B (36 marks)

- 7 Fig. 7 shows the curve $y = \frac{x^2 + 3}{x - 1}$. It has a minimum at the point P. The line l is an asymptote to the curve.




Fig. 7

- (i) Write down the equation of the asymptote l . [1]
- (ii) Find the coordinates of P. [6]
- (iii) Using the substitution $u = x - 1$, show that the area of the region enclosed by the x -axis, the curve and the lines $x = 2$ and $x = 3$ is given by

$$\int_1^2 \left(u + 2 + \frac{4}{u} \right) du.$$

Evaluate this area exactly. [7]

- (iv) Another curve is defined by the equation $e^y = \frac{x^2 + 3}{x - 1}$. Find $\frac{dy}{dx}$ in terms of x and y by differentiating implicitly. Hence find the gradient of this curve at the point where $x = 2$. [4]

- 8 Fig. 8 shows part of the curve $y = f(x)$, where $f(x) = e^{-\frac{1}{5}x} \sin x$, for all x .




Fig. 8

- (i) Sketch the graphs of

(A) $y = f(2x)$,

(B) $y = f(x + \pi)$.

[4]

- (ii) Show that the x -coordinate of the turning point P satisfies the equation $\tan x = 5$.

Hence find the coordinates of P.

[6]

- (iii) Show that $f(x + \pi) = -e^{-\frac{1}{5}\pi} f(x)$. Hence, using the substitution $u = x - \pi$, show that

$$\int_{\pi}^{2\pi} f(x) dx = -e^{-\frac{1}{5}\pi} \int_0^{\pi} f(u) du.$$

Interpret this result graphically. [You should *not* attempt to integrate $f(x)$.]

[8]

6
BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.