

**ADVANCED GCE
MATHEMATICS (MEI)**

4756/01

Further Methods for Advanced Mathematics (FP2)

WEDNESDAY 9 JANUARY 2008

Afternoon

Time: 1 hour 30 minutes

Additional materials: Answer Booklet (8 pages)
Graph paper
MEI Examination Formulae and Tables (MF2)

INSTRUCTIONS TO CANDIDATES

- Write your name in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions in Section A and **one** question from Section B.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.

This document consists of 4 printed pages.

Section A (54 marks)

Answer all the questions

- 1 (a) Fig. 1 shows the curve with polar equation $r = a(1 - \cos 2\theta)$ for $0 \leq \theta \leq \pi$, where a is a positive constant.




Fig. 1

Find the area of the region enclosed by the curve. [7]

- (b) (i) Given that $f(x) = \arctan(\sqrt{3} + x)$, find $f'(x)$ and $f''(x)$. [4]

(ii) Hence find the Maclaurin series for $\arctan(\sqrt{3} + x)$, as far as the term in x^2 . [4]

(iii) Hence show that, if h is small, $\int_{-h}^h x \arctan(\sqrt{3} + x) dx \approx \frac{1}{6}h^3$. [3]

- 2 (a) Find the 4th roots of $16j$, in the form $re^{j\theta}$ where $r > 0$ and $-\pi < \theta \leq \pi$. Illustrate the 4th roots on an Argand diagram. [6]

- (b) (i) Show that $(1 - 2e^{j\theta})(1 - 2e^{-j\theta}) = 5 - 4 \cos \theta$. [3]

Series C and S are defined by

$$C = 2 \cos \theta + 4 \cos 2\theta + 8 \cos 3\theta + \dots + 2^n \cos n\theta,$$

$$S = 2 \sin \theta + 4 \sin 2\theta + 8 \sin 3\theta + \dots + 2^n \sin n\theta.$$

- (ii) Show that $C = \frac{2 \cos \theta - 4 - 2^{n+1} \cos(n+1)\theta + 2^{n+2} \cos n\theta}{5 - 4 \cos \theta}$, and find a similar expression for S . [9]

3 You are given the matrix $\mathbf{M} = \begin{pmatrix} 7 & 3 \\ -4 & -1 \end{pmatrix}$.

(i) Find the eigenvalues, and corresponding eigenvectors, of the matrix \mathbf{M} . [8]

(ii) Write down a matrix \mathbf{P} and a diagonal matrix \mathbf{D} such that $\mathbf{P}^{-1}\mathbf{M}\mathbf{P} = \mathbf{D}$. [2]

(iii) Given that $\mathbf{M}^n = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, show that $a = -\frac{1}{2} + \frac{3}{2} \times 5^n$, and find similar expressions for b , c and d . [8]

Section B (18 marks)

Answer one question

Option 1: Hyperbolic functions

4 (i) Given that $k \geq 1$ and $\cosh x = k$, show that $x = \pm \ln(k + \sqrt{k^2 - 1})$. [5]

(ii) Find $\int_1^2 \frac{1}{\sqrt{4x^2 - 1}} dx$, giving the answer in an exact logarithmic form. [5]

(iii) Solve the equation $6 \sinh x - \sinh 2x = 0$, giving the answers in an exact form, using logarithms where appropriate. [4]

(iv) Show that there is no point on the curve $y = 6 \sinh x - \sinh 2x$ at which the gradient is 5. [4]

[Question 5 is printed overleaf.]

Option 2: Investigation of curves

This question requires the use of a graphical calculator.

5 A curve has parametric equations $x = \frac{t^2}{1+t^2}$, $y = t^3 - \lambda t$, where λ is a constant.

(i) Use your calculator to obtain a sketch of the curve in each of the cases

$$\lambda = -1, \quad \lambda = 0 \quad \text{and} \quad \lambda = 1.$$

Name any special features of these curves. [5]

(ii) By considering the value of x when t is large, write down the equation of the asymptote. [1]

For the remainder of this question, assume that λ is positive.

(iii) Find, in terms of λ , the coordinates of the point where the curve intersects itself. [3]

(iv) Show that the two points on the curve where the tangent is parallel to the x -axis have coordinates

$$\left(\frac{\lambda}{3+\lambda}, \pm \sqrt{\frac{4\lambda^3}{27}} \right). \quad [4]$$

Fig. 5 shows a curve which intersects itself at the point $(2, 0)$ and has asymptote $x = 8$. The stationary points A and B have y -coordinates 2 and -2 .




Fig. 5

(v) For the curve sketched in Fig. 5, find parametric equations of the form $x = \frac{at^2}{1+t^2}$, $y = b(t^3 - \lambda t)$, where a , λ and b are to be determined. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.