



Physics A

PHYA2

Unit 2 Mechanics, Materials and Waves

Data and Formulae Booklet

DATA

FUNDAMENTAL CONSTANTS AND VALUES

| <i>Quantity</i> | <i>Symbol</i> | <i>Value</i> | <i>Units</i> |
|--|---------------|---------------------------|-----------------------------------|
| speed of light in vacuo | c | 3.00×10^8 | m s^{-1} |
| permeability of free space | μ_0 | $4\pi \times 10^{-7}$ | H m^{-1} |
| permittivity of free space | ϵ_0 | 8.85×10^{-12} | F m^{-1} |
| charge of electron | e | -1.60×10^{-19} | C |
| the Planck constant | h | 6.63×10^{-34} | J s |
| gravitational constant | G | 6.67×10^{-11} | $\text{N m}^2 \text{kg}^{-2}$ |
| the Avogadro constant | N_A | 6.02×10^{23} | mol^{-1} |
| molar gas constant | R | 8.31 | $\text{J K}^{-1} \text{mol}^{-1}$ |
| the Boltzmann constant | k | 1.38×10^{-23} | J K^{-1} |
| the Stefan constant | σ | 5.67×10^{-8} | $\text{W m}^{-2} \text{K}^{-4}$ |
| the Wien constant | α | 2.90×10^{-3} | m K |
| electron rest mass (equivalent to 5.5×10^{-4} u) | m_e | 9.11×10^{-31} | kg |
| electron charge/mass ratio | e/m_e | 1.76×10^{11} | C kg^{-1} |
| proton rest mass (equivalent to 1.00728 u) | m_p | $1.67(3) \times 10^{-27}$ | kg |
| proton charge/mass ratio | e/m_p | 9.58×10^7 | C kg^{-1} |
| neutron rest mass (equivalent to 1.00867 u) | m_n | $1.67(5) \times 10^{-27}$ | kg |
| gravitational field strength | g | 9.81 | N kg^{-1} |
| acceleration due to gravity | g | 9.81 | m s^{-2} |
| atomic mass unit (1u is equivalent to 931.3 MeV) | u | 1.661×10^{-27} | kg |

GEOMETRICAL EQUATIONS

| | |
|---------------------------------|------------------------|
| <i>arc length</i> | $= r\theta$ |
| <i>circumference of circle</i> | $= 2\pi r$ |
| <i>area of circle</i> | $= \pi r^2$ |
| <i>surface area of cylinder</i> | $= 2\pi rh$ |
| <i>volume of cylinder</i> | $= \pi r^2h$ |
| <i>area of sphere</i> | $= 4\pi r^2$ |
| <i>volume of sphere</i> | $= \frac{4}{3}\pi r^3$ |

ASTRONOMICAL DATA

| <i>Body</i> | <i>Mass/kg</i> | <i>Mean radius/m</i> |
|-------------|-----------------------|----------------------|
| Sun | 1.99×10^{30} | 6.96×10^8 |
| Earth | 5.98×10^{24} | 6.37×10^6 |

AS FORMULAE

PARTICLE PHYSICS

Rest energy values

| class | name | symbol | rest energy /MeV |
|---------|-------------|-----------|------------------|
| photon | photon | γ | 0 |
| lepton | neutrino | ν_e | 0 |
| | | ν_μ | 0 |
| | electron | e^\pm | 0.510999 |
| | muon | μ^\pm | 105.659 |
| mesons | π meson | π^\pm | 139.576 |
| | | π^0 | 134.972 |
| | K meson | K^\pm | 493.821 |
| | | K^0 | 497.762 |
| baryons | proton | p | 938.257 |
| | neutron | n | 939.551 |

Properties of quarks

antiquarks have opposite signs

| type | charge | baryon number | strangeness |
|----------|-----------------|----------------|-------------|
| u | $+\frac{2}{3}e$ | $+\frac{1}{3}$ | 0 |
| d | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | 0 |
| s | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | -1 |

Properties of Leptons

| | lepton number |
|---|---------------|
| particles: $e^-, \nu_e; \mu^-, \nu_\mu$ | +1 |
| antiparticles: $e^+, \bar{\nu}_e; \mu^+, \bar{\nu}_\mu$ | -1 |

Photons and Energy Levels

photon energy $E = hf = hc/\lambda$

photoelectricity $hf = \phi + E_{K(\max)}$

energy levels $hf = E_1 - E_2$

de Broglie Wavelength $\lambda = \frac{h}{p} = \frac{h}{mv}$

ELECTRICITY

current and pd $I = \frac{\Delta Q}{\Delta t}$ $V = \frac{W}{Q}$ $R = \frac{V}{I}$

emf $\varepsilon = \frac{E}{Q}$ $\varepsilon = I(R + r)$

resistors in series $R = R_1 + R_2 + R_3 + \dots$

resistors in parallel $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$

resistivity $\rho = \frac{RA}{L}$

power $P = VI = I^2R = \frac{V^2}{R}$

alternating current $I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$ $V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$

MECHANICS

moments moment = Fd

velocity and acceleration $v = \frac{\Delta s}{\Delta t}$ $a = \frac{\Delta v}{\Delta t}$

equations of motion $v = u + at$ $s = \frac{(u+v)t}{2}$

$v^2 = u^2 + 2as$ $s = ut + \frac{at^2}{2}$

force $F = ma$

work, energy and power $W = Fs \cos \theta$ $E_K = \frac{1}{2}mv^2$ $\Delta E_p = mg\Delta h$

$P = \frac{\Delta W}{\Delta t}$, $P = Fv$

efficiency = $\frac{\text{useful output power}}{\text{input power}}$

MATERIALS

density $\rho = \frac{m}{V}$ Hooke's law $F = k \Delta L$

Young modulus = $\frac{\text{tensile stress}}{\text{tensile strain}}$ tensile stress = $\frac{F}{A}$
tensile strain = $\frac{\Delta L}{L}$

energy stored $E = \frac{1}{2}F\Delta L$

WAVES

wave speed $c = f\lambda$ period $T = \frac{1}{f}$

fringe spacing $w = \frac{\lambda D}{s}$ diffraction grating $d \sin \theta = n\lambda$

refractive index of a substance $n = \frac{c}{c_s}$

for two different substances of refractive indices n_1 and n_2 ,

law of refraction $n_1 \sin \theta_1 = n_2 \sin \theta_2$

critical angle $\sin \theta_c = \frac{n_2}{n_1}$ for $n_1 > n_2$