

Physics (B): Physics in Context


Data and Formulae Booklet

FUNDAMENTAL CONSTANTS AND OTHER NUMERICAL DATA

| Quantity | Symbol | Value | Units |
|------------------------------|--------------|--------------------------------|-----------------------------------|
| speed of light in vacuo | c | 3.00×10^8 | m s^{-1} |
| Planck constant | h | 6.63×10^{-34} | J s |
| gravitational constant | G | 6.67×10^{-11} | $\text{N m}^2 \text{kg}^{-2}$ |
| gravitational field strength | g | 9.81 | N kg^{-1} |
| acceleration due to gravity | g | 9.81 | m s^{-2} |
| electron rest mass | m_e | 9.11×10^{-31} | kg |
| | m_e | $5.5 \times 10^{-4} \text{ u}$ | |
| electron charge | e | -1.60×10^{-19} | C |
| proton rest mass | m_p | $1.67(3) \times 10^{-27}$ | kg |
| | m_p | 1.00728 u | |
| neutron rest mass | m_n | $1.67(5) \times 10^{-27}$ | kg |
| | m_n | 1.00867 u | |
| permittivity of free space | ϵ_0 | 8.85×10^{-12} | F m^{-1} |
| molar gas constant | R | 8.31 | $\text{J K}^{-1} \text{mol}^{-1}$ |
| Boltzmann constant | k | 1.38×10^{-23} | J K^{-1} |
| Avogadro constant | N_A | 6.02×10^{23} | mol^{-1} |
| Wien constant | α | 2.90×10^{-3} | m K |

GEOMETRICAL EQUATIONS

| | |
|--------------------------|-----------------------------|
| arc length | $r\theta$ |
| circumference of circle | $2\pi r$ |
| area of circle | πr^2 |
| surface area of sphere | $4\pi r^2$ |
| volume of sphere | $\frac{4}{3}\pi r^3$ |
| surface area of cylinder | $2\pi rh$ |
| volume of cylinder | $\pi r^2 h$ |
| | $\sin \theta = \frac{a}{c}$ |
| | $\cos \theta = \frac{b}{c}$ |
| | $\tan \theta = \frac{a}{b}$ |
| | $c^2 = a^2 + b^2$ |



Unit Conversions

| | |
|------------------------|------------------------------------|
| 1 atomic mass unit (u) | $1.661 \times 10^{-27} \text{ kg}$ |
| 1 year (y) | $3.15 \times 10^7 \text{ s}$ |
| 1 parsec (pc) | $3.08 \times 10^{16} \text{ m}$ |
| 1 parsec | 3.26 ly |
| 1 light year (ly) | $9.46 \times 10^{15} \text{ m}$ |

Particle Properties

Properties of quarks *antiquarks have opposite signs*

| type | charge | Baryon number | strangeness |
|----------|-----------------|----------------|-------------|
| u | $+\frac{2}{3}e$ | $+\frac{1}{3}$ | 0 |
| d | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | 0 |
| s | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | -1 |

Properties of Leptons

| | Lepton Number |
|---|---------------|
| <i>particles:</i> $e^-, \nu_e; \mu^-, \nu_\mu; \tau^-, \nu_\tau$ | +1 |
| <i>antiparticles:</i> $e^+, \bar{\nu}_e; \mu^+, \bar{\nu}_\mu; \tau^+, \bar{\nu}_\tau$ | -1 |

AS FORMULAE

| | Waves | | Quantum Physics and Astrophysics |
|--|---|-------------------------------------|--|
| wave speed | $c = f\lambda$ | photon energy | $E = hf$ |
| period | $T = \frac{1}{f}$ | Einstein equation | $hf = \phi + E_{k(\max)}$ |
| intensity | $I = \frac{P}{A}$ | line spectrum equation | $hf = E_1 - E_2$ |
| stretched string frequency | $f = \frac{1}{2L} \sqrt{\frac{T}{\mu}}$ | de Broglie wavelength | $\lambda = \frac{h}{p} = \frac{h}{mv}$ |
| beat frequency | $f = f_1 - f_2$ | Doppler shift for $v \ll c$ | $\frac{\Delta f}{f} = -\frac{\Delta \lambda}{\lambda} = \frac{v}{c}$ |
| fringe spacing | $w = \frac{\lambda D}{s}$ | Wien's law | $\lambda_{\max} T = 0.0029 \text{ m K}$ |
| diffraction grating | $n\lambda = d \sin \theta$ | Hubble law | $v = H d$ |
| half beam width | $\sin \theta = \frac{\lambda}{a}$ | intensity for a point source | $I = \frac{P}{4\pi r^2}$ |
| refractive index of a substance (s) | $n = \frac{c}{c_s}$ | | |
| for two different substances of refractive indices n_1 and n_2 | $n_1 \sin i_1 = n_2 \sin i_2$ | | |
| critical angle | $\sin \theta_c = \frac{n_2}{n_1}$ for $n_1 > n_2$ | | |
| | Mechanics | | Electricity |
| speed or velocity | $v = \frac{\Delta s}{\Delta t}$ | current | $I = \frac{\Delta Q}{\Delta t}$ |
| acceleration | $a = \frac{\Delta v}{\Delta t}$ | electromotive force (emf) | $\varepsilon = \frac{E}{Q}$ |
| equations of motion | $v = u + at$ $s = \frac{(u+v)}{2} t$ $v^2 = u^2 + 2as$ $s = ut + \frac{1}{2} at^2$ | resistance | $\varepsilon = I(R+r)$ |
| force | $F = ma$ | resistors in series | $R = \frac{V}{I}$ |
| change in potential energy | $\Delta E_p = mg\Delta h$ | resistors in parallel | $R = R_1 + R_2 + R_3 + \dots$ $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$ |
| kinetic energy | $E_k = \frac{1}{2} mv^2$ | resistivity | $\rho = \frac{RA}{L}$ |
| momentum | $p = mv$ | power | $P = VI = I^2 R = \frac{V^2}{R}$ |
| impulse | $F\Delta t = \Delta(mv)$ | potential divider formula | $V_o = \left(\frac{R_1}{R_1 + R_2} \right) \times V_i$ |
| spring stiffness | $k = \frac{F}{\Delta L}$ | energy | $E = VIt$ |
| energy stored for $F \propto L$ | $E = \frac{1}{2} F\Delta L$ | efficiency | $\frac{\text{useful output power}}{\text{input power}}$ |
| work done | $W = Fs \cos \theta$ | | |
| power | $P = \frac{\Delta W}{\Delta t} = Fv$ | | |
| density | $\rho = \frac{m}{V}$ | | |
| | | | Energy production and transmission |
| | | rate of heat transfer by conduction | $= UA \Delta \theta$ |
| | | maximum power for a wind turbine | $= \frac{1}{2} \pi r^2 \rho v^3$ |

| Circular Motion | | Gases and Thermal Physics | |
|-----------------------------|---|--|--|
| angular velocity | $\omega = \frac{v}{r}$ | pressure | $p = \frac{F}{A}$ |
| angular acceleration | $\alpha = \frac{\Delta\omega}{\Delta t}$ | gas law (N is number of atoms) | $pV = NkT$ |
| angular frequency | $\omega = 2\pi f$ | gas law (n is quantity in mol) | $pV = nRT$ |
| centripetal force | $F = \frac{mv^2}{r} = m\omega^2 r$ | kinetic theory model | $pV = \frac{1}{3}Nm\langle c^2 \rangle$ |
| centripetal acceleration | $a = \frac{v^2}{r} = r\omega^2$ | kinetic energy of gas molecule | $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$ |
| angular momentum | $L = I\omega$ | energy to change temperature | $E = mc\Delta\theta$ |
| angular kinetic energy | $E_k = \frac{1}{2}I\omega^2$ | first law of thermodynamics | $\Delta U = Q + W$ $W = \text{work done on the system}$ |
| moment of inertia | $I = \frac{Tr}{\alpha}$ | entropy change | $\Delta S = \frac{Q}{T}$ |
| torque | $T = Fd$ | maximum thermal efficiency | $\eta = \frac{T_H - T_C}{T_H}$ |
| equations of angular motion | $\omega_2 = \omega_1 + \alpha t$ $\omega_2^2 = \omega_1^2 + 2\alpha\theta$ $\theta = \frac{(\omega_1 + \omega_2)}{2}t$ $\theta = \omega_1 t + \frac{1}{2}\alpha t^2$ | work done | $W = p\Delta V$ |
| power | $P = T\omega$ | | |
| Oscillations | | Radioactivity and nuclear physics | |
| acceleration | $a = -(2\pi f)^2 x$ | absorption of radiation | $I = I_0 e^{-\mu x}$ |
| displacement | $x = A \cos(2\pi f t)$ | radioactive decay | $N = N_0 e^{-\lambda t}$ |
| maximum speed | $v_{\max} = 2\pi f A$ | half-life | $T_{1/2} = \frac{\ln 2}{\lambda}$ |
| maximum acceleration | $a_{\max} = (2\pi f)^2 A$ | radioactive change represented by | $dN/dt = -\lambda N$ |
| for a mass-spring system | $T = 2\pi\sqrt{\frac{m}{k}}$ | activity | $A = \lambda N$ |
| for a simple pendulum | $T = 2\pi\sqrt{\frac{l}{g}}$ | mass-energy equivalence | $\Delta E = \Delta mc^2$ |