

Teacher Resource Bank

GCE Physics B: Physics in Context Sample A2 ISA: Damped Harmonic Motion

• Mark Scheme



Copyright © 2009 AQA and its licensors. All rights reserved.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334). Registered address: AQA, Devas Street, Manchester M15 6EX. *Dr Michael Cresswell*, Director General.

AQA Physics Sample A2 ISA, Damped Harmonic Motion, Mark Scheme

Stage 1		Mark	Additional guidance notes
(a)	table with column headings showing all recorded results and correct units \checkmark	1	column headings can be either in words or standard symbols
			units can be in words or the correct abbreviation e.g. time/seconds, T/s
			alternative acceptable labelling includes time(s), time in s etc
(b)	significant figures correct for all readings \checkmark	1	distances quoted to nearest mm, e.g. 3.8 cm and timings compatible with precision of stopclock or stopwatch
(c)	at least two readings for the time of the oscillations and each scale reading \checkmark	1	e.g. records time for 10 oscillations (or 20 oscillations) etc and shows at least one repeat
(d)	correct computation of the amplitude and $t \checkmark$	1	sample of calculations should be checked
(e)	correct computation of $\ln A \checkmark$	1	no sf penalty on these values, sample of values checked
(f)	suitably large graph scale (do not award if scale on axis could have been doubled) \checkmark	1	the plotted points should occupy at least half of each axis
(g)	graph heading and correctly labelled axes with units \checkmark	1	alternative method of labelling axes as in (a) above table headings
(h)	most points accurately plotted to within 1 mm (no more than one point > ± 1 mm) and line of best fit drawn ✓ 1	1	this mark is independent of mark (g), i.e. if candidates have used an unsuitable scale they can still achieve marks for accurately plotting the points
			markers should check a proportion of plotted points
			'most points' would typically allow one incorrect plot in a sample of 6 or 7 plotted points
			the line should be a straight line with approximately an equal number of points on either side of the line
			points which are obviously anomalous should not unduly influence the line
			if the plotted points suggest a curve line, the mark can be awarded for a suitable smooth curve
	Total	8	



Section A		Mark	Additional guidance notes
Question 1			
(a)	amplitude ✓	1	
(b)	mass and spring (or force) constant \checkmark	1	
(c)	reaction time may vary ✓ miscounting oscillations ✓	2	do not accept 'human error'
(d)	realistic estimate from spread of repeat times \checkmark	1	
(e)	% uncertainty = $\frac{\text{uncertainty}}{\text{largest time}} \times 100 \checkmark$	1	
(f) (i) (ii) (iii)	take logs \checkmark In $A = \ln A_0 - \lambda t \checkmark$ - $\lambda \checkmark$ correct comment on validity of a straight line graph for their plotted points eg points lie close to a straight line \checkmark	4	
(g)	for example: use a position sensor ✓	1	accept other sensible suggestions
	Total	11	

Section B		Mark	Additional guidance notes
Question 2			
(a)	all 5 correct average <i>t</i> _{1/2} values: 4.4, 12.2, 22.6, 34.7, 42.2 ✓		
	all 5 correct <i>λ</i> ⁻¹ values: 6.21, 17.5, 32.6, 50.0, 61.0 ✓	3	must not be to more significant figures
	correct significant figures ✓		
(b)	correct scales and axes ✓	3	
	all plotted points to nearest mm ✓		the line should be a straight line with approximately an equal number of points on either side of the line
	best line of fit to points \checkmark		
(C)	large triangle drawn (at least 8 cm \times 8 cm) \checkmark	3	
	correct values read from graph \checkmark		
	correct answer for gradient = 0.69 \checkmark		
	(allow 0.68 to 0.70 or calculation based on ecf from incorrectly read scale value(s))		
(d)	difference or % difference quoted \checkmark	2	
	discusses scatter of points ✓		
	Total	11	

Question 3		Mark	Additional guidance notes
(a)	0.2s (half the spread) ✓	1	
(b)	random ✓	1	
(C)	closeness of repeat readings ✓	1	
(d) (i)	uncertainty in λ^{-1} is also 0.2% \checkmark		
(ii)	uncertainty = $\frac{0.20}{42.2}$ × 100 = 0.47% ✓	3	no sf penalty
(iii)	0.2 + 0.47 = 0.67% ✓		
(e)	$t_{\rm 1/2}$ is inversely proportional to λ \checkmark	1	
	Total	7	

Question 4		Mark	Additional guidance notes
	keeping card horizontal ✓	max 4	
	ensuring oscillations are vertical \checkmark		
	arranging the metre rule close enough \checkmark		
	card colliding with metre ruler \checkmark		
	$t_{ m 1/2}$ not occurring at the end of an oscillation \checkmark		
	card wobbles \checkmark		
	Total	4	