

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

Advanced GCE

PHYSICS A 2826/01

Unifying Concepts in Physics

Thursday 15 JUNE 2006 Morning 1 hour 15 minutes

Candidates answer on the question paper. Additional materials: Electronic calculator

Candidate Name	 Centr	e Nı	umbe	er	and Num	idate ber	!

TIME 1 hour 15 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer all the questions.
- Write your answers in the spaces provided on the question paper.
- Read the questions carefully and make sure you know what you have to do before starting your answer.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You may use a scientific calculator.
- You are advised to show all the steps in any calculations.

FOR EXAMINER'S USE					
Qu.	Max.	Mark			
1	10				
2	7				
3	14				
4	12				
5	17				
TOTAL	60				

Data

speed of light in free space,	$c = 3.00 \times 10^8 \mathrm{ms^{-1}}$
permeability of free space,	$\mu_0 = 4\pi \times 10^{-7}~{\rm Hm^{-1}}$
permittivity of free space,	$\epsilon_0 = 8.85 \times 10^{-12} \mathrm{F m^{-1}}$
elementary charge,	$e = 1.60 \times 10^{-19} \mathrm{C}$
the Planck constant,	$h = 6.63 \times 10^{-34} \mathrm{J}\mathrm{s}$
unified atomic mass constant,	$u = 1.66 \times 10^{-27} \text{ kg}$
rest mass of electron,	$m_{\rm e} = 9.11 \times 10^{-31} \rm kg$
rest mass of proton,	$m_{\rm p} = 1.67 \times 10^{-27} \rm kg$
molar gas constant,	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
the Avogadro constant,	$N_{\rm A} = 6.02 \times 10^{23} {\rm mol}^{-1}$
gravitational constant,	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
acceleration of free fall,	$g = 9.81 \text{ m s}^{-2}$

Formulae

uniformly accelerated motion,
$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

refractive index,
$$n = \frac{1}{\sin C}$$

capacitors in series,
$$\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} + \dots$$

capacitors in parallel,
$$C = C_1 + C_2 + \dots$$

capacitor discharge,
$$x = x_0 e^{-t/CR}$$

pressure of an ideal gas,
$$p = \frac{1}{3} \frac{Nm}{V} < c^2 >$$

radioactive decay,
$$X = X_0 e^{-\lambda t}$$

$$t_{\frac{1}{2}} = \frac{0.693}{\lambda}$$

critical density of matter in the Universe,
$$\rho_0 = \frac{3H_0^2}{8\pi G}$$

relativity factor,
$$= \sqrt{(1 - \frac{v^2}{c^2})}$$

current,
$$I = nAve$$

nuclear radius,
$$r = r_0 A^{1/3}$$

sound intensity level,
$$= 10 \lg \left(\frac{I}{I_0}\right)$$

2826 Jun06 [Turn over


Answer all the questions.


1	avai app	ilable aratı	measurements are quite easy to take accurately if laboratory instruments are a. They can also be done accurately – to 2 or 3 significant figures – using simple as such as a 3-metre tape measure, kitchen scales and a watch. Outline, using these ly, how you could determine the following.
	(a)	The	thickness of a sheet of paper in a paperback book.
			[2]
	(b)	The	mass of a paperclip.
	(c)	(i)	The diameter of a large cylindrical stone column in a cathedral.
			[2]
		(ii)	Explain the problem in measuring the diameter directly by holding the tape measure stretched alongside of the column.


(d) (i)	The length of a pavement in a road outside	a school.
		[2]
(ii)	The average speed of a car travelling alon the pavement.	
		[2]
		[Total: 10]
For eac	h of the following, give the full name of the SI	unit used.
(a) ele	ctric charge	
(b) cap	pacitance	
(c) free	quency	
(d) stre	ess	
(e) gra	vitational field strength	
(f) ma	gnetic flux	
(g) rad	lioactive activity	[7]
		[Total: 7]
		• •


2


3 Data can be displayed in graphical form in many different ways. Sometimes it is nece change from one way of displaying data to another. Four graphs are drawn on page			
	(a)	(i)	Calculate the total distance travelled from the velocity-time graph A .
			distance = m [3]
		(ii)	Using graph A , draw the corresponding distance-time graph. [3]
	(b)		ph B shows how the current <i>I</i> in a circuit varies with the total circuit resistance <i>R</i> en the e.m.f. of the supply is kept constant.
		(i)	Draw the corresponding graph of 1/ <i>I</i> against <i>R</i> . [2]
		(ii)	What is the e.m.f. of the supply?
			e.m.f. = V [1]
		(iii)	How is the gradient of the graph you have drawn related to your answer to (b)(ii)?
	(c)		ph C shows how g , the acceleration due to gravity, varies with r , the distance from centre of the Earth. A log-log graph showing the same data has been drawn on new s.
		(i)	Calculate the gradient of the log-log graph.
			gradient =[2]
		(ii)	What can be deduced from the value of the gradient?
			[2]
			[Total: 14]














- 4 (a) The three states of matter, solid, liquid and gas, are characterised by having different separations and patterns of movement of the molecules. Discuss the changes that take place to the following list of quantities when melting and boiling occur. Assume that no chemical changes occur.
 - · separation of molecules
 - speed of molecules
 - · pattern of movement of molecules
 - internal energy of molecules

In your answer, comment on similarities and differences between melting and boiling.

	[9]
(b)	Suggest and explain two differences which might exist between hydrogen gas at 200 K and hydrogen at 200 000 000 K.
	[3]
	[Total: 12]

2826 Jun06

[Turn over

Read	d the following paragraph and answer the questions about it which follow.
	An extract of text has been removed due to third party copyright restriction
	Details: An extract of text adapted from an article by Ian Robinson in Physics Worl May 2004 about the kilogram
(a)	(Adapted from an article by Ian Robinson in Physics World, May 2004.) State two circumstances which could very slightly change the mass of the standard kilogram when it is being used for a mass determination.
	2
(b)	What percentage uncertainty is there in a mass measurement accurate to 1 μg in 1
	percentage uncertainty =% [2]
(c)	Silicon is used in the suggested re-definition because the technology is available to produce large, very pure single crystals of silicon from which the sphere of mass one kilogram and diameter 94.0 mm can be made.
(c)	produce large, very pure single crystals of silicon from which the sphere of mass one

(ii) Calculate the uncertainty in measuring the diameter of the silicon sphere if the

		uncertainty in the volume is to be only 2 parts in 10 ⁸ .
		uncertainty =[2]
(d)	Kno acc	spacing of atoms in silicon can be measured very accurately using X-ray diffraction. wing this spacing, the number of silicon atoms in the sphere can be calculated urately providing the silicon is pure and the crystal structure is regular. Suggest and ain a problem with obtaining high accuracy if one of these conditions is not met.
		[2]
(e)	Silio	con has 3 stable isotopes.
	(i)	What is meant by the term isotope?
		[1]
	(ii)	Why is it important to know accurately the relative proportions of these isotopes in the silicon sphere?
		[2]
	(iii)	It would be preferable to have the sphere made entirely of one isotope. Suggest why this is difficult to achieve.
		[2]

[Total: 17]

