Please check the examination details below before entering your candidate information

Candidate surname	Other names

Pearson Edexcel Level 3 GCE

Centre Number | | | | | |

Candidate Number | | | | |

Thursday 14 May 2020

Afternoon (Time: 1 hour 30 minutes) | Paper Reference **8PS0/02**

Psychology

Advanced Subsidiary

Paper 2: Biological Psychology and Learning Theories

You do not need any other materials.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*

Information

- The total mark for this paper is 70.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- The list of formulae and statistical tables are printed at the start of this paper.
- Candidates may use a calculator.

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶

P62572A

1/1/1/

P 6 2 5 7 2 A 0 1 3 2

Pearson

FORMULAE AND STATISTICAL TABLES

Standard deviation (sample estimate)

$$\sqrt{\left(\frac{\sum(x-\bar{x})^2}{n-1}\right)}$$

Spearman's rank correlation coefficient

$$1-\frac{6\sum d^2}{n(n^2-1)}$$

Critical values for Spearman's rank

	Level of significance for a one-tailed test				
	0.05	**0.025**	**0.01**	**0.005**	**0.0025**
	Level of significance for a two-tailed test				
N	**0.10**	**0.05**	**0.025**	**0.01**	**0.005**
5	0.900	1.000	1.000	1.000	1.000
6	0.829	0.886	0.943	1.000	1.000
7	0.714	0.786	0.893	0.929	0.964
8	0.643	0.738	0.833	0.881	0.905
9	0.600	0.700	0.783	0.833	0.867
10	0.564	0.648	0.745	0.794	0.830
11	0.536	0.618	0.709	0.755	0.800
12	0.503	0.587	0.678	0.727	0.769
13	0.484	0.560	0.648	0.703	0.747
14	0.464	0.538	0.626	0.679	0.723
15	0.446	0.521	0.604	0.654	0.700
16	0.429	0.503	0.582	0.635	0.679
17	0.414	0.485	0.566	0.615	0.662
18	0.401	0.472	0.550	0.600	0.643
19	0.391	0.460	0.535	0.584	0.628
20	0.380	0.447	0.520	0.570	0.612
21	0.370	0.435	0.508	0.556	0.599
22	0.361	0.425	0.496	0.544	0.586
23	0.353	0.415	0.486	0.532	0.573
24	0.344	0.406	0.476	0.521	0.562
25	0.337	0.398	0.466	0.511	0.551
26	0.331	0.390	0.457	0.501	0.541
27	0.324	0.382	0.448	0.491	0.531
28	0.317	0.375	0.440	0.483	0.522
29	0.312	0.368	0.433	0.475	0.513
30	0.306	0.362	0.425	0.467	0.504

The calculated value must be equal to or exceed the critical value in this table for significance to be shown.

Chi-squared distribution formula

$$X^2 = \sum \frac{(O-E)^2}{E} \qquad df = (r-1)(c-1)$$

Critical values for chi-squared distribution

	Level of significance for a one-tailed test					
	0.10	**0.05**	**0.025**	**0.01**	**0.005**	**0.0005**
	Level of significance for a two-tailed test					
df	**0.20**	**0.10**	**0.05**	**0.025**	**0.01**	**0.001**
1	1.64	2.71	3.84	5.02	6.64	10.83
2	3.22	4.61	5.99	7.38	9.21	13.82
3	4.64	6.25	7.82	9.35	11.35	16.27
4	5.99	7.78	9.49	11.14	13.28	18.47
5	7.29	9.24	11.07	12.83	15.09	20.52
6	8.56	10.65	12.59	14.45	16.81	22.46
7	9.80	12.02	14.07	16.01	18.48	24.32
8	11.03	13.36	15.51	17.54	20.09	26.12
9	12.24	14.68	16.92	19.02	21.67	27.88
10	13.44	15.99	18.31	20.48	23.21	29.59
11	14.63	17.28	19.68	21.92	24.73	31.26
12	15.81	18.55	21.03	23.34	26.22	32.91
13	16.99	19.81	22.36	24.74	27.69	34.53
14	18.15	21.06	23.69	26.12	29.14	36.12
15	19.31	22.31	25.00	27.49	30.58	37.70
16	20.47	23.54	26.30	28.85	32.00	39.25
17	21.62	24.77	27.59	30.19	33.41	40.79
18	22.76	25.99	28.87	31.53	34.81	42.31
19	23.90	27.20	30.14	32.85	36.19	43.82
20	25.04	28.41	31.41	34.17	37.57	45.32
21	26.17	29.62	32.67	35.48	38.93	46.80
22	27.30	30.81	33.92	36.78	40.29	48.27
23	28.43	32.01	35.17	38.08	41.64	49.73
24	29.55	33.20	36.42	39.36	42.98	51.18
25	30.68	34.38	37.65	40.65	44.31	52.62
26	31.80	35.56	38.89	41.92	45.64	54.05
27	32.91	36.74	40.11	43.20	46.96	55.48
28	34.03	37.92	41.34	44.46	48.28	56.89
29	35.14	39.09	42.56	45.72	49.59	58.30
30	36.25	40.26	43.77	46.98	50.89	59.70
40	47.27	51.81	55.76	59.34	63.69	73.40
50	58.16	63.17	67.51	71.42	76.15	86.66
60	68.97	74.40	79.08	83.30	88.38	99.61
70	79.72	85.53	90.53	95.02	100.43	112.32

The calculated value must be equal to or exceed the critical value in this table for significance to be shown.

Mann-Whitney U test formulae

$$U_a = n_a n_b + \frac{n_a(n_a+1)}{2} - \sum R_a$$

$$U_b = n_a n_b + \frac{n_b(n_b+1)}{2} - \sum R_b$$

(U is the smaller of U_a and U_b)

Critical values for the Mann-Whitney U test

	N_b															
	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
N_a																
***p* ≤ 0.05 (one-tailed), *p* ≤ 0.10 (two-tailed)**																
5	4	5	6	8	9	11	12	13	15	16	18	19	20	22	23	25
6	5	7	8	10	12	14	16	17	19	21	23	25	26	28	30	32
7	6	8	11	13	15	17	19	21	24	26	28	30	33	35	37	39
8	8	10	13	15	18	20	23	26	28	31	33	36	39	41	44	47
9	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54
10	11	14	17	20	24	27	31	34	37	41	44	48	51	55	58	62
11	12	16	19	23	27	31	34	38	42	46	50	54	57	61	65	69
12	13	17	21	26	30	34	38	42	47	51	55	60	64	68	72	77
13	15	19	24	28	33	37	42	47	51	56	61	65	70	75	80	84
14	16	21	26	31	36	41	46	51	56	61	66	71	77	82	87	92
15	18	23	28	33	39	44	50	55	61	66	72	77	83	88	94	100
16	19	25	30	36	42	48	54	60	65	71	77	83	89	95	101	107
17	20	26	33	39	45	51	57	64	70	77	83	89	96	102	109	115
18	22	28	35	41	48	55	61	68	75	82	88	95	102	109	116	123
19	23	30	37	44	51	58	65	72	80	87	94	101	109	116	123	130
20	25	32	39	47	54	62	69	77	84	92	100	107	115	123	130	138

N_a	N_b = 5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$p \leq 0.01$ (one-tailed), $p \leq 0.02$ (two-tailed)																
5	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
6	2	3	4	6	7	8	9	11	12	13	15	16	18	19	20	22
7	3	4	6	7	9	11	12	14	16	17	19	21	23	24	26	28
8	4	6	7	9	11	13	15	17	20	22	24	26	28	30	32	34
9	5	7	9	11	14	16	18	21	23	26	28	31	33	36	38	40
10	6	8	11	13	16	19	22	24	27	30	33	36	38	41	44	47
11	7	9	12	15	18	22	25	28	31	34	37	41	44	47	50	53
12	8	11	14	17	21	24	28	31	35	38	42	46	49	53	56	60
13	9	12	16	20	23	27	31	35	39	43	47	51	55	59	63	67
14	10	13	17	22	26	30	34	38	43	47	51	56	60	65	69	73
15	11	15	19	24	28	33	37	42	47	51	56	61	66	70	75	80
16	12	16	21	26	31	36	41	46	51	56	61	66	71	76	82	87
17	13	18	23	28	33	38	44	49	55	60	66	71	77	82	88	93
18	14	19	24	30	36	41	47	53	59	65	70	76	82	88	94	100
19	15	20	26	32	38	44	50	56	63	69	75	82	88	94	101	107
20	16	22	28	34	40	47	53	60	67	73	80	87	93	100	107	114

N_a	N_b = 5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$p \leq 0.025$ (one-tailed), $p \leq 0.05$ (two-tailed)																
5	2	3	5	6	7	8	9	11	12	13	14	15	17	18	19	20
6	3	5	6	8	10	11	13	14	16	17	19	21	22	24	25	27
7	5	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34
8	6	8	10	13	15	17	19	22	24	26	29	31	34	36	38	41
9	7	10	12	15	17	20	23	26	28	31	34	37	39	42	45	48
10	8	11	14	17	20	23	26	29	33	36	39	42	45	48	52	55
11	9	13	16	19	23	26	30	33	37	40	44	47	51	55	58	62
12	11	14	18	22	26	29	33	37	41	45	49	53	57	61	65	69
13	12	16	20	24	28	33	37	41	45	50	54	59	63	67	72	76
14	13	17	22	26	31	36	40	45	50	55	59	64	67	74	78	83
15	14	19	24	29	34	39	44	49	54	59	64	70	75	80	85	90
16	15	21	26	31	37	42	47	53	59	64	70	75	81	86	92	98
17	17	22	28	34	39	45	51	57	63	67	75	81	87	93	99	105
18	18	24	30	36	42	48	55	61	67	74	80	86	93	99	106	112
19	19	25	32	38	45	52	58	65	72	78	85	92	99	106	113	119
20	20	27	34	41	48	55	62	69	76	83	90	98	105	112	119	127

N_a	N_b 5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
***p* ≤ 0.005 (one-tailed), *p* ≤ 0.01 (two-tailed)**																
5	0	1	1	2	3	4	5	6	7	7	8	9	10	11	12	13
6	1	2	3	4	5	6	7	9	10	11	12	13	15	16	17	18
7	1	3	4	6	7	9	10	12	13	15	16	18	19	21	22	24
8	2	4	6	7	9	11	13	15	17	18	20	22	24	26	28	30
9	3	5	7	9	11	13	16	18	20	22	24	27	29	31	33	36
10	4	6	9	11	13	16	18	21	24	26	29	31	34	37	39	42
11	5	7	10	13	16	18	21	24	27	30	33	36	39	42	45	48
12	6	9	12	15	18	21	24	27	31	34	37	41	44	47	51	54
13	7	10	13	17	20	24	27	31	34	38	42	45	49	53	56	60
14	7	11	15	18	22	26	30	34	38	42	46	50	54	58	63	67
15	8	12	16	20	24	29	33	37	42	46	51	55	60	64	69	73
16	9	13	18	22	27	31	36	41	45	50	55	60	65	70	74	79
17	10	15	19	24	29	34	39	44	49	54	60	65	70	75	81	86
18	11	16	21	26	31	37	42	47	53	58	64	70	75	81	87	92
19	12	17	22	28	33	39	45	51	56	63	69	74	81	87	93	99
20	13	18	24	30	36	42	48	54	60	67	73	79	86	92	99	105

The calculated value must be equal to or less than the critical value in this table for significance to be shown.

Wilcoxon Signed Ranks test process

- Calculate the difference between two scores by taking one from the other
- Rank the differences giving the smallest difference Rank 1

  Note: do not rank any differences of 0 and when adding the number of scores, do not count those with a difference of 0, and ignore the signs when calculating the difference
- Add up the ranks for positive differences
- Add up the ranks for negative differences
- T is the figure that is the smallest when the ranks are totalled (may be positive or negative)
- N is the number of scores left, ignore those with 0 difference

Critical values for the Wilcoxon Signed Ranks test

	Level of significance for a one-tailed test		
	0.05	0.025	0.01
	Level of significance for a two-tailed test		
n	0.1	0.05	0.02
N=5	0	-	-
6	2	0	-
7	3	2	0
8	5	3	1
9	8	5	3
10	11	8	5
11	13	10	7
12	17	13	9

The calculated value must be equal to or less than the critical value in this table for significance to be shown.

Answer ALL questions.

SECTION A: BIOLOGICAL PSYCHOLOGY

1 When studying biological psychology you will have learned about different types of correlations.

(a) State what is meant by a positive and a negative correlation.

(2)

Positive correlation

Negative correlation

(b) Explain **one** strength and **one** weakness of the correlational research method.

(4)

Strength

Weakness

(Total for Question 1 = 6 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2 Max has been learning about evolution and natural selection. He has been asked by his psychology teacher to prepare a presentation to show how evolution and natural selection explain human aggressive behaviour. His teacher has suggested he includes:

- a reference to gene selection
- a short video
- a reference to inherited traits
- a research study.

Describe what Max could include in his presentation to show how evolution and natural selection explain human aggressive behaviour.

(Total for Question 2 = 4 marks)

BLANK PAGE

3 Isaac is a psychologist who has investigated whether alcohol affected the ability of participants to complete a task successfully. The task consisted of participants walking 20 steps along a straight line whilst balancing a book on their head after drinking alcohol.

Isaac gathered an opportunity sample from his local university. He asked them to drink 2 units of alcohol (units being a measurement of alcohol) on day one and 4 units on day two.

Table 1 shows the results for Isaac's investigation.

Participants	Number of mistakes made on day one (2 units of alcohol)	Number of mistakes made on day two (4 units of alcohol)
A	6	6
B	12	11
C	4	14
D	8	19

Table 1

(a) Calculate the percentage of mistakes made by participant C on day one from the total of mistakes made by all participants on day one.

You must give your answer to **one** decimal place.

(1)

SPACE FOR CALCULATIONS

Percentage .....

(b) Calculate the mean number of mistakes made on day two.

You must give your answer to **one** decimal place.

(1)

SPACE FOR CALCULATIONS

Mean number of mistakes made on day two .....

(c) Explain **one** strength of the opportunity sampling technique used by Isaac in his investigation.

(2)

.....

.....

.....

.....

.....

.....

(d) Explain **one** improvement that Isaac could make to his investigation.

(2)

.....

.....

.....

.....

.....

.....

(Total for Question 3 = 6 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

4 Amanda is struggling to control her aggression. Her doctor is considering sending Amanda for an fMRI scan to find out if her brain may be influencing her aggression.

(a) Explain **two** reasons for Amanda's doctor sending her for an fMRI scan.

(4)

1 .....

.....

.....

.....

.....

.....

.....

.....

2 .....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Name **one** brain scanning technique other than fMRI.

(1)

.....

.....

(Total for Question 4 = 5 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

5 Assess whether the role of hormones can explain human behaviour such as aggression.

(8)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 = 8 marks)

TOTAL FOR SECTION A = 29 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

SECTION B: LEARNING THEORIES

6 In learning theories, you will have learned about one of the following contemporary studies:

- Becker et al. (2002)
- Bastian et al. (2011)
- Capafóns et al. (1998).

(a) State **one** aim of your chosen contemporary study.

(1)

(b) Describe the results of your chosen contemporary study.

(2)

(c) Explain **one** strength and **one** weakness of your chosen contemporary study.

(4)

Strength

Weakness

(Total for Question 6 = 7 marks)

7 You were required to carry out a practical investigation when studying learning theories.

Describe the procedure for your practical investigation in learning theories.

(Total for Question 7 = 3 marks)

8 Bertie wanted to investigate the severity of symptoms for older and younger patients after treatment for a phobia.

He sampled two groups of patients who had received treatment and were recovering from a phobia:

- 20 patients aged between 20–35 years
- 20 patients aged between 40–55 years.

Bertie asked the patients to rate the severity of their symptoms after treatment. He used a rating scale ranging from 1–10, where 1–5 was judged as 'less severe' symptoms and 6–10 was judged as 'more severe' symptoms.

Bertie's results are shown in **Table 2** below.

Age group	Patients with 'less severe' symptoms after treatment	Patients with 'more severe' symptoms after treatment
20–35 years	5	15
40–55 years	11	9

Table 2

(a) Complete **Table 3** to calculate the chi-squared test for Bertie's data to **two** decimal places.

(4)

		Observed	Expected	O–E	$(O–E)^2$	$(O–E)^2/E$
20–35 years	**Patients with 'less severe' symptoms after treatment**	5	8			
	Patients with 'more severe' symptoms after treatment	15	12			
40–55 years	**Patients with 'less severe' symptoms after treatment**	11	8			
	Patients with 'more severe' symptoms after treatment	9	12			
					Chi-squared =	

Table 3

SPACE FOR CALCULATIONS

Chi-squared (X^2) .....

(b) Explain **one** strength of Bertie's study in terms of reliability.

(2)

(c) Explain **one** improvement that Bertie could make to his study in terms of validity.

(2)

(Total for Question 8 = 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

9 Bradley decided to use a content analysis to investigate how much violence is shown in children's television programmes.

Describe how Bradley could use a content analysis to investigate violence in children's television programmes.

(Total for Question 9 = 3 marks)

10 Evaluate social learning theory as an explanation of human behaviour.

(8)

(Total for Question 10 = 8 marks)

TOTAL FOR SECTION B = 29 MARKS

SECTION C

11 To what extent do classical conditioning, the process of synaptic transmission and neurotransmitter functioning explain human behaviour?

(12)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 11 = 12 marks)

TOTAL FOR SECTION C = 12 MARKS
TOTAL FOR PAPER = 70 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

BLANK PAGE

BLANK PAGE