



GCE MARKING SCHEME

**PHYSICS
AS/Advanced**

SUMMER 2014

INTRODUCTION

The marking schemes which follow were those used by WJEC for the Summer 2014 examination in GCE PHYSICS. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

	Page
PH1	1
PH2	6
PH3	10
PH4	13
PH5	20
PH6	36

Question			Marking details	Marks Available
3	(a)	(i)	J s^{-1}	[3×1]
		(ii)	V A^{-1}	
		(iii)	A s	
	(b)	(i)	$t = 2 \times 3\,600$ or $7\,200 \text{ s}$ (1) $Q = 0.15 \times 7\,200 = 1\,080 \text{ [C]}$ (1)	[2]
		(ii)	$\frac{6480}{1080} = 6 \text{ [V]}$ (ecf on Q)	[1]
		(iii)	$\frac{5832}{1080} = 5.4 \text{ [V]}$ (ecf on Q)	[1]
		(iv)	$6 - 5.4 = 0.6 \text{ [V]}$ (1) (ecf from (b)(ii) & (iii)) $\frac{0.6}{0.15} = 4 \text{ [\Omega]}$ (1) (ecf on 0.6 [V])	
			Or Correct substitution into $V = E - Ir$ (i.e. $5.4 = 6.0 - 0.15r$) (1) $r = 4 \text{ [\Omega]}$ (1) (ecf from (b)(ii) & (iii))	[2]
			Alternative Solution: $\frac{(6480 - 5832)}{7200} = 0.09 \text{ J s}^{-1}$ (Lost energy in cell per second) (1) $I^2 r = 0.09$ and $r = 4 \text{ [\Omega]}$ (1)	
			Question 3 Total	[9]

Question		Marking details	Marks Available
4	(a)	<p><u>Electrical energy (or work done) transferred [to other forms passing] between two points (1) per coulomb of charge (1)</u> Definition of 1 V award 1 mark only</p>	[2]
	(b)	<p>(i) $V_{\text{supply}} = V_1 + V_2 + V_3$</p> <p>(ii) Energy</p>	[1] [1]
	(c)	<p>(i) $R_1 + 12 = \frac{9}{0.5}$ (1) Clear manipulation seen to show $R_1 = 6[\Omega]$ (1)</p> <p>(ii) (I) $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$ to show effective parallel combination = 6Ω (1) this can be implied V across upper 6Ω resistor shown = 4.5 [V] (ecf on parallel combination) (1)</p> <p>(II) Total resistance = 12Ω (1) $I = \frac{9.0}{12} = 0.75\text{ [A]}$ (1) (accept $\frac{4.5}{6} = 0.75\text{ [A]}$)</p> <p>(III) $1.2 = \frac{9}{(6 + R_{\text{parallel}})}$ (1) $R_{\text{parallel}} = 1.5\text{ [}\Omega\text{]}$ (1) $n \times \left(\frac{1}{12}\right) = \frac{1}{1.5}$ (1) ecf on $1.5\text{ [}\Omega\text{]}$ $n = 8$ (1) Full marks for correct answer based on trial and error Alternative solution: $\frac{9}{1.2} = 7.5\text{ [}\Omega\text{]}$ (1) $7.5 - 6 = 1.5\text{ [}\Omega\text{]}$ (1) $\frac{12}{n} = 1.5\text{ [}\Omega\text{]}$ (1) $n = 8$ (1)</p>	[2] [2] [2] [4]
Question 4 Total			[14]

Question			Marking details	Marks Available
5	(a)	(i)	Ruler and wire shown and labelled (1) Moving pointer or jockey or crocodile clip indicated (1) Either: Correctly positioned ohmmeter with no power supply; or correctly positioned voltmeter and ammeter with power supply (1) [No labelling required for either method].	[3]
		(ii)	Diagonal line through origin	[1]
		(iii)	CSA from <u>diameter of wire</u> (1) Gradient from graph = (R/l) or (ρ/A) Or stated take a pair of R and l values from the graph (1) $\rho = \text{gradient} \times \text{CSA}$ or use of $\rho = RA/l$ (1)	[3]
	(b)	(i)	$R = \frac{144}{32} = 4.5 [\Omega]$ (1) Correct substitution into $R = \rho l/A$ (1) $l = 0.375 [\text{m}]$ (1) (ecf on R)	[3]
		(ii)	$I = 2.7 [\text{A}]$ (from V/R or P/V etc) (1) (ecf on I) Correct substitution into $I = nAve$ (1) $v = 1.24 \times 10^{-2} [\text{m s}^{-1}]$ (1) accept 0.01 m s^{-1}	[3]
			Question 5 Total	[13]

Question			Marking details	Marks Available	
6	(a)	(i)	Acceleration defined as rate of change of <u>velocity</u> [or equivalent] or $a = \frac{(v-u)}{t} \quad (1)$ <u>Clear manipulation</u> to show that $v=u+at$ (1)	[2]	
		(ii)	$v=u+at$ substituted into $x = (u+v)t/2$ (1) <u>Clear manipulation</u> shown (1)	[2]	
	(b)	(i)	A (1) Horizontal velocity (= 65 m s^{-1}) constant or same speed as plane or sack lands directly underneath plane (1) Vertical velocity increases or there is a vertical acceleration (1)	[3]	
		(ii)	(I)	Substitution into $v^2=u^2+2ax$ and $u = 0$ shown (1) x calculated = 45.9 [m] (1)	[2]
			(II)	Correct substitution into $v = at$ or $x=1/2at^2$ or $x = \frac{(u+v)t}{2}$ (1) $t=3.1 \text{ [s]}$ (1)	[2]
		(iii)	$v_R^2 = (65^2 + 30^2)$ (correct substitution into Pythagoras) (1) $v_R = 71.6 \text{ [m s}^{-1}\text{]}$ (1) Valid angle calculated <u>and shown</u> or described e.g. $\theta = 24.8^\circ$ below horizontal (1)	[3]	
Question 6 Total			[14]		
7	(a)	Replace <i>mass</i> with <i>force</i> (1) Don't accept weight Introduce <u>perpendicular distance to pivot</u> (1)	[2]		
	(b)	$(2 \times 700) - 1\,200$ (1) Weight of beam = 200 [N] (1) Alternative solution: Moment about A or B e.g. $(700 \times 5) = (1\,200 + W) \times 2.5$	[2]		
	(c)	(i)		[2]	
			Upward forces as shown and indicated (1) Downward forces as shown and indicated (1) N.B. $1\,200 \text{ [N]}$ force can be indicated anywhere between W and F_B		
			(ii)	Taking moments about A: $F_B \times 5.0$ (1) $(1\,200 \times 3.5) + (200 \times 2.5)$ (1) (ecf on 200) $F_B = 940 \text{ [N]}$ (1)	[3]
(iii)	$1\,400 - 940 = 460 \text{ [N]}$ (ecf from (b) and/or (c)(ii)) Accept answer based on moments calculated about B.	[1]			
Question 7 Total			[10]		

PH2

Question			Marking details	Marks Available
1	(a)	(i)	Attempt at sinusoid, right way up, passing within 1 mm of all dots	1
		(ii)	P and Q are in phase (1) Amplitude of P > amplitude of Q (1)	2
(iii)		Q and R are in antiphase / exactly out of phase (1) Amplitude of Q = amplitude of R (1)	2	
(iv)		$\frac{\lambda}{2} = 0.20$ [m] or $\lambda = 0.40$ [m] or by implication (1) $v = 96 \text{ m s}^{-1}$ UNIT ecf (1)	2	
	(b)		$\frac{\lambda}{2} = 0.15$ [m] (or $\lambda = 0.30$ [m]) or $v = 96$ [m s ⁻¹] ecf from (a)(iv) or $f = \left(\frac{4}{3}\right) 240$ [Hz] or by implication (1) $f = 320$ [Hz] but not by cancellation of errors, ecf on v from (a)(iv) (1)	2
			Question 1 total	[9]
2	(a)	(i)	$S_2Q = \sqrt{(350^2 + 120^2)}$ [mm] or equivalent (1) Therefore $S_2Q - S_1Q = (370 - 350)$ [mm] (1)	2
		(ii)	For any dot, path difference = $n\lambda$, or for P, path difference = 0 or any other remark relevant to the conclusion that ... (1) $\lambda = 10$ [mm] (1)	2
		(iii)	$\lambda = \left(\frac{120 \times 30}{350}\right)$ (1) $\lambda = 10 \text{ mm}$ or 10.3 mm UNIT (1)	2
	(b)	With sensor in front of source either rotate sensor [at least through 90°] or interpose array of metal rods /metal grille and rotate [at least through 90°] (1) Don't accept metal grid Signal strength changes (1) Accept in words or in diagram	2	
			Question 2 total	[8]

Question		Marking details	Marks Available	
3	(a)	[Flat, opaque] screen / sheet/ plate / material with slits / gaps (1) Slits are parallel / vertical or equally spaced or closely spaced or many / multiple (1)	2	
	(b)	(i) $\frac{1}{400000} = [2.5 \times 10^{-6} \text{ m}]$	1	
		(ii) $2\lambda = 2.5 \times 10^{-6} \sin 25.2^\circ$ even with the 2 missing or mishandled (1) Correct placing of the 2 (1) $\lambda = 532 \times 10^{-9} \text{ [m]}$ ecf on d only (1)	3	
		(iii) $3 \times 532 = 2500 \sin \theta$ or equivalent ecf on λ (1) $\theta = 39.7^\circ$ or 40° ecf on λ (1)	2	
		(iv) Young's slits much further apart than slits in grating Don't accept slits much narrower or gaps are much smaller	1	
		Question 3 Total	[9]	
4	(a)	(i) medium 1: $2.0 \times 10^8 \text{ [m s}^{-1}\text{]}$ and medium 2: $2.5 \times 10^8 \text{ [m s}^{-1}\text{]}$	1	
		(ii) Correct use of $\sin 30^\circ$ seen clearly (1) Rest of argument, including use of $t = \frac{d}{v}$ [ecf on v and on value of $\sin 30^\circ$, if failure to reach the stated time is noted]. (1)	2	
		(iii) $BD = 2.5 \times 10^8 \text{ ecf} \times 2.5 \times 10^{-11} \text{ [m]}$ [= 6.25 mm] or by implication (1) $\theta_2 = 38.7^\circ$ (or 39°) ecf on $v = 2.5 \times 10^8 \text{ [m s}^{-1}\text{]}$ (1)	2	
		(iv) $1.50 \sin 30^\circ = 1.20 \sin \theta_2$ (1) Therefore $\theta_2 = 38.7^\circ$ (or 39°) no ecf (1)	2	
		(b)	(i) Use of $v = 2.0 \times 10^8 \text{ [m s}^{-1}\text{]}$ (1) $t = \frac{1600}{2.0 \times 10^8} \text{ [s]}$ ecf on v (1)	2
	(ii) Critical angle = 76° or by implication (1) $n_{\text{clad}} [\times \sin 90^\circ] = 1.500 \sin 76^\circ$ ecf on 76° or by implication (1) $n_{\text{clad}} = 1.455$ or 1.46 do not accept 1.45 no ecf (1)	3		
	(iii) $\frac{AC}{AB} = \cos 14^\circ$ or equivalent or by implication (1) $\Delta t = 0.24 \mu\text{s}$ ecf on v (1)	2		
			Question 4 Total	[14]

Question		Marking details	Marks Available	
5	(a)	[Minimum] energy needed to release [or eject] electron from magnesium [or metal or surface or solid not atom]	1	
	(b)	$E_{k \max} = 6.63 \times 10^{-34} \times 1.16 \times 10^{15} \text{ [J]} - 5.9 \times 10^{-19} \text{ [J]}$ (1) $E_{k \max} = 1.79 \times 10^{-19} \text{ [J]}$ (1)	2	
	(c)	<u>Photon</u> energy < work function (1) don't accept photon energy in symbols. Accept not enough energy to liberate an electron. Don't accept $E_{k \max}$ can't be negative. $E_{\text{phot}} = 5.4 \times 10^{-19} \text{ [J]}$ accept $f_{\text{thresh}} = 8.9 \times 10^{14} \text{ [Hz]}$ (1) If negative energy award 1 mark only	2	
	(d)	(i)	Planck constant. Accept Planck's constant or h .	1
		(ii)	[-] work function. Accept [-] ϕ .	1
(iii)		f_0 or minimum frequency to eject electron or threshold frequency	1	
Question 5 Total			[8]	
6	(a)	(i)	Any 2 × (1) from: • Monochromatic or same frequency or same wavelength • Wavefronts continuous or light in phase across width of beam • <u>Photons</u> in phase	2
		(ii)	Use of $E = hf$ and $f = \frac{c}{\lambda}$ or $E = \frac{hc}{\lambda}$ (1) $1.87 \times 10^{-19} \text{ [J]}$ (1)	2
		(iii)	$1.3 \times 10^{20} \text{ [s}^{-1}\text{]}$ ecf	1
		(iv)	Downward arrow from U to L (1) $2.29 \times 10^{-19} \text{ J}$ (or $2.3 \times 10^{-19} \text{ J}$) (1) ecf	2
	(b)	[Passing] photon stimulates electron to drop <u>from U to L</u> (1) Emitting another photon (1) Any 2 × (1) from: • Process may happen repeatedly (or equivalent) as photons traverse cavity • Population inversion [between U and L] needed for stimulated emission to predominate over absorption • Pumping to P and drop to U brings about inversion • Level L self-emptying so less pumping needed or population inversion easier to accomplish • In phase with or travelling in the same direction as or polarised in the same direction as or identical to passing photon • Stimulated photon must have an energy of $1.87 \times 10^{-19} \text{ J}$ or equivalent	4	
Question 6 Total			[11]	

Question		Marking details	Marks Available	
7	(a)	(i) $\lambda_{\text{peak}} = \frac{2.90 \times 10^{-3}}{9900} \text{ [m]}$ or equivalent (1) $\lambda_{\text{peak}} = 293 \times 10^{-9} \text{ [m]}$ (1)	2	
		(ii) Peak between 280 and 300 nm (1) Curve goes through origin [with zero gradient at origin] and is consistent with approaching zero at very long wavelengths (1)	2	
		(iii) Blue accept white or violet or purple	1	
	(b)	$A = \frac{L}{\sigma T^4}$ with A as subject, with symbols or data or $1.84 \times 10^{19} \text{ m}^2$ (1) Attempt to use $A = 4\pi r^2$ and $d = 2r$ or $A = \pi I^2$ (1) $d = 2.4 \times 10^9 \text{ m}$ ecf on slips of 2^n or 10^n if already penalised (1)	3	
	(c)	(i) Absorption accept excitation Don't accept pumping	1	
		(ii) Dark / black lines crossing or missing wavelengths [continuous] spectrum or coloured background	1	
		(iii) B almost absent and any reference to populations of levels (1) First excited state not populated [so no transitions start here] or all electrons in ground state (1)	2	
	Question 7 Total			[12]
	8	(a)	(i) $uud + u\bar{u}d \rightarrow uud + u\bar{d}d$ (1) $+ u\bar{d}$ (1)	2
			(ii) $1 + 1 > 1 + 1 + 0$ (all numbers must be shown) or equivalent	1
(iii) Strong because no [photons (gammas) or] neutrinos or no flavour changes			1	
(iv) Charge or momentum or energy or strangeness Accept up quark number or down quark number			1	
(b)		(i) $0 + 0 > 0 + (-1) + 1$ (all numbers must be shown)	1	
		(ii) Weak interaction accept fusion (1) Takes place in the Sun [accept stars] (1) Part of the process whereby we get sunlight or energy or equivalent (1)	3	
Question 8 Total			[9]	

PH3

TEST 1 – Mark Scheme

SECTION A

A1.

- (a) Appropriate measurements taken to ± 1 mm. (1)
Volume calculated correctly with correct unit. Ignore sig figs. (1) [2]
- (b) Using correct instrument resolution for uncertainty (± 1 mm) Accept ($\pm \frac{1}{2}$ mm). (1)
All percentage uncertainties calculated correctly. **ecf** on resolution (1)
Percentage uncertainties added to give final value. (1) No sig fig penalty. [3]
(Allow **ecf** on incorrect % uncertainty; need to add 3 values)
- (c) (i) Absolute uncertainty calculated correctly. No sig fig or unit penalty. **ecf** [1]
- (ii) Value of the absolute uncertainty to 1 or 2 sig figs and
volume to the same precision **and unit quoted**
e.g. $152 \pm 11 \text{ cm}^3$ or $150000 \pm 10000 \text{ mm}^3$. **ecf** [1]
- (d) Measuring instrument with a lower instrument resolution (accept reference to precision
but not accuracy) OR use of calipers or micrometer [1]

Total [8]

A2.

- (a) Unit of extension in the table (accept mm or cm or m). (1)
Extension measured consistently to the nearest mm. (1)
Mean value of k calculated correctly. (1) No sig fig penalty. Answer must be seen in (a)
Unit of k correct. (1) (Accept: N m^{-1} or N mm^{-1} or kg s^{-2} or $\text{kg m s}^{-2} \text{ mm}^{-1}$.) [4]
- (b) Absolute uncertainty calculated correctly. (1) No sig fig or unit penalty.
Percentage uncertainty calculated correctly. **ecf** (1) No sig fig penalty. [2]
N.B. If all values for k are the same then candidate needs to state that the
uncertainty is 0 (1) and the % uncertainty is 0. (1)
Or if candidate calculates % uncertainty in extension (1) % uncertainty in $k =$ %
uncertainty in extension. (1)
- (c) Correct rearrangement of equation $m = \frac{kx}{g}$ or implied. (1)
Value of $m = 0.120$ kg. **Unit required.** (1) (Accept 0.11–0.13)
Accept answer in grammes. [2]

Total [8]

A3.

- (a) Value of correct time or velocity repeated and mean taken. (1)
Velocity calculated correctly including unit (m s^{-1}). **ecf** (1) No sig fig penalty. [2]
- (b) (i) All units correct. (1)
Columns for mean time and mean velocity correct. (1)
Values for v^2 correct. **ecf** (1)
Consistent use of 2 dps in each column of data. (1) [4]
- (ii) Valid conclusion made with some reference to the data. (1)
Ratio $\frac{n}{v^2}$ or $\frac{v^2}{n}$ calculated for all 3 rows. (1) [2]
- Total [8]

SECTION B**B4**

- (a) (i) Circuit diagram drawn with correct symbols (ignore positions of voltmeter and ammeter for this mark). (1)
Voltmeter and ammeter correctly positioned. (1) [2]
- (ii) Change length and measure V and I . (1)
Reference to $R = \frac{V}{I}$. (1)
Suitable intervals stated allow intervals up to a maximum of 0.2 m. (1) [3]
None of the above can be awarded from the table.
- (b) Clear headings (length or l / current or I / voltage or pd or V / resistance or R) and correct units on all columns. (1)
Values of voltage; current; given in sequential values of length. (1)
Resistance calculated correctly. (1)
All I and V values given to 2 dp (1) [4]
- (c) Graph of resistance against length plotted with axes labelled and correct units given on both axes. (1) **ecf from table**
Suitable scale chosen (don't allow multiples of 3) so that all data points occupy at least half the graph paper. (1)
All points plotted correctly to within $\pm \frac{1}{2}$ small square division. (1)
1 good line of best fit consistent with the data. (1)
2 good distinct lines of best fit drawn showing difference in gradient. (1) [5]
- (d) Distance correctly read from the graph. (allow $\pm \frac{1}{2}$ small square division) (1)
Unit and value given to the nearest cm or consistent with the scale used on the graph.
Apply **ecf** if distance read incorrectly. (1) [2]

- (e) (i) Large triangle used (1) (should be close to the extremities of the line of best fit for wire starting at X) [or 2 equivalent suitable points clearly indicated on the graph].
 Correct values used for gradient calculation. (1)
 Gradient calculated correctly. **ecf**. No unit penalty. (1) If wire starting at Y do not award this mark. [3]
- (ii) Measuring the diameter of the wire starting at X. (1) No unit or sig fig penalty.
 Accept in the range 0.28-0.33
 Correct calculation of the cross-sectional area. (1) No unit or sig fig penalty.
 Realising that gradient of the graph = $\frac{\rho}{A}$ (can be applied anywhere in the answer) or using data values from the line of the graph. (1)
 Calculating a value for resistivity in Ω m. (1) **ecf** for gradient and area.
 No unit or sig fig penalty. If data taken from the table and it is not on the graph line award 1 mark only.
 Correct conclusion consistent with their resistivity value. (1) [5]
 (Candidates need to calculate a value for resistivity before conclusion mark can be awarded.)

Total [24]


TEST 2 – Mark Scheme as TEST 1

Except:

A2(c) 2nd mark: Value of $m = 0.160$ kg. Unit required. (1) (Accept 0.15–0.17)

A3 (b)(ii) 2nd mark: Ratio $\frac{n}{v^2}$ or $\frac{v^2}{n}$ calculated for at least 3 rows of data or 2 sets of data. (1)


PH4

Question		Marking details	Marks Available
1	(a)	<p>Reasonable attempt at conservation of momentum (1) e.g. $330\,000m = \pm 10\,000m + 6.6 \times 10^{-27} \times v_1$</p> <p>conservation of momentum applied correctly and values substituted (1) e.g. $330\,000 \times 3.4 \times 10^{-25} = -10\,000 \times 3.3 \times 10^{-25} + 6.6 \times 10^{-27} \times v_1$</p> <p>correct answer = $1.75 \times 10^7 \text{ [m s}^{-1}\text{]}$ (no ecf) (1)</p>	3
	(b)	<p>(i) Any valid answer e.g. impulse (or force or acceleration or change in momentum) is vertical, gamma has no momentum in horizontal direction, perpendicular directions are independent etc. Accept: no horizontal force</p> <p>(ii) Attempt at using $p = \frac{h}{\lambda}$ (1)</p> <p>$E = hf$ and $c = f\lambda$ quoted (or equivalent $E = \frac{hc}{\lambda}$) (1)</p> <p>N.B. $p = \frac{E}{c}$ gains 2 marks</p> <p>Correct momentum = 6.33×10^{-22} (1)</p> <p>Answer = $= \frac{6.33 \times 10^{-22}}{3.3 \times 10^{-25}}$ [$1\,920 \text{ m s}^{-1}$] (1)</p>	1 4
	(iii)	<p>Method i.e. $\sqrt{10000^2 + 2000^2}$ (1)</p> <p>Answer = $10\,200 \text{ [m s}^{-1}\text{]}$ ecf on v from (b)(ii) (1)</p> <p>Method and correct indication of angle e.g. $\tan^{-1}\left(\frac{2000}{10000}\right)$ (1)</p> <p>Answer = 11.5° or 0.2 [rad] (or $90-11.5$ for other angle if indicated etc.) (1)</p> <div style="text-align: right;">  </div>	4
Question 1 Total			[12]

Question		Marking details	Marks Available
2	(a)	(i) (Number of moles) $n = 4.73$ (1) Mass = 4×4.73 or 0.004×4.73 (or implied) (1) Density = $0.004 \times 4.73 / 0.113$ [= 0.167] (1)	3
	(b)	(ii) Either $p = \frac{1}{3} \rho \overline{c^2}$ used or equivalent e.g. $\frac{3}{2} nRT = \frac{1}{2} M \overline{c^2}$ (1) 1 350 [m s^{-1}] (1) Density = $0.004 \times 4.73 / 0.212$ or $T = \frac{45000 \times 0.212}{4.73 \times 8.31}$ ecf (1) $p = \frac{1}{3} \rho \overline{c^2}$ used or $\frac{3}{2} nRT = \frac{1}{2} M \overline{c^2}$ used or equivalent (1) Answer = 1 230 [m s^{-1}] (1) Question 2 Total	2 [8]
3	(a)	Substitution into $v = \sqrt{\frac{GM}{r}}$ (1) Answer = 158 000 [m s^{-1}] (1)	2
	(b)	Measured velocity is greater (1) Which implies that the mass is greater (1) Suggests the existence of dark matter (1) Question 3 Total	3 [5]

Question		Marking details	Marks Available
4	(a)	Mass substituted into $T = 2\pi \sqrt{\frac{m}{k}}$ (1) $T = \frac{1}{f}$ used or implied (1) Answer = 152 N m ⁻¹ UNIT mark (1)	3
	(b)	$3.47 \times 2\pi [= 21.803]$	1
	(c)	(i) $v = \omega A [= 1.853]$ or max PE = max KE (1) $KE = \frac{1}{2}mv^2$ used or = $\frac{1}{2}kx^2$ (1) Answer = 0.55 [J] (1)	3
		(ii) Acceleration = $\omega^2 A$ or $F = kA$ Accept $F = kA - mg$ (1) Answer = 12.9 [N] (1)	2
	(d)	Substitution of values e.g. $-1.4 = 8.5\sin(21.8 \times 0.1 + \varepsilon)$ (1) $\sin^{-1}\left(\frac{-1.4}{8.5}\right) = -0.165$ (1) $\varepsilon = -2.35$ or equivalent in degree (-135°) or other quadrant (-5.16) ecf on minus sign (1)	3
		Question 4 total	[12]

Question		Marking details	Marks Available
5	(a)	(i) Force per unit mass (this minimalist answer is acceptable unless some contradiction)	1
		(ii) Work done per unit mass <u>from infinity</u> (this minimalist answer is acceptable unless some contradiction)	1
	(b)	(i) $F = \frac{GMm}{r^2}$ used (1) Answer = 22.8 [N] (1)	2
		(ii) $PE = [-] \frac{GMm}{r}$ used or equivalent (1) Answer = - 13.7 M[J] (1)	2
	(c)	$PE = [-] \frac{GMm}{r}$ used or equivalent (1) Answer = - 61.8 M[J] (ecf on – sign) (1)	2
	(d)	Difference in PE attempted (1) Correct answer = 48.1 M[J] ((b)(ii) – (c)) ecf (1) Answer must be consistent with their signs	2
	Question 5 Total		[10]

Question		Marking details	Marks Available
6	(a)	<p>All arrows correct ✓✓</p> <p>Directions in line with dotted lines but some (or all) directions inverted ✓</p> 	2
	(b)	$E = \frac{Q}{4\pi\epsilon_0 r^2}$ used (1) <p>Answer = 1 500 V m⁻¹ or NC⁻¹ or equivalent UNIT mark (1)</p>	2
	(c)	<p><u>Field of</u> 13 μC ×2 and ×12/13 (1)</p> <p>Answer = 222 [V m⁻¹] (1)</p> <p>To the left or implied clearly in the calculation (1)</p>	3
	(d)	$V = \frac{Q}{4\pi\epsilon_0 r}$ used for 3 charges with $r = 12$ or 13 (1) $V = \frac{1}{4\pi\epsilon_0} \left(2 \frac{13}{13} - \frac{24}{12} \right)$ as shown or equivalent (cm perfectly valid) (1)	2
	(e)	<p>Any 3 (×1) from:</p> <ul style="list-style-type: none"> • initial total energy is zero / initial and final PE is zero • final total energy is zero / initial and final KE is zero • initial force is to the right (has to be linked to the field and the negative charge) • later the force is to the left (but not a resistive force) <p>Question 6 Total</p>	3
			[12]

Question		Marking details	Marks Available
7	(a)	$T = 2\pi \sqrt{\frac{(3 \times 10^{10})^3}{6.67 \times 10^{-11} \times (7 \times 10^{29} + 4 \times 10^{28})}} \quad (1)$ <p>Answer = 4.65×10^6 [s] (1) (4.78×10^6 s scores 1/2 marks)</p>	2
	(b)	$r_1 = \frac{M_2}{M_1 + M_2} d \quad \text{used or } M_1 r_1 = M_2 r_2 \quad \text{used (1)}$ <p>Star orbit radius = 0.162×10^{10} [m] (1) (0.171×10^{10} scores 1/2 marks)</p>	2
	(c)	$v = \frac{2\pi r}{T} \quad \text{or } v = \omega r \quad \text{and } \omega = 2\pi f \quad \text{ecf on } T \text{ and } r \text{ (1)}$ $v = \frac{2\pi \times 0.162 \times 10^{10}}{4.65 \times 10^6} [= 2191] \text{ (1)}$ $\frac{\Delta\lambda}{\lambda} = \frac{v}{c} \quad \text{attempted or rearranged ecf on } v \text{ (1)}$ <p>Answer = 4.8×10^{-12} [m] (1)</p>	4
	(d)	<p>Hotter or the Earth is cooler or equivalent (1)</p> <p>Due to higher intensity [of e-m radiation] (1) Accept because $5^2 > 20$ or similar</p> <p>Question 7 Total</p>	2
			[10]

Question		Marking details	Marks Available																																			
8	(a)	(i) $T = \frac{pV}{nR}$ seen or equivalent or implied (1) $T = \frac{95000 \times 0.79}{28.9 \times 8.31}$ (= 312.5 K) (1)	2																																			
		(ii) $U = \frac{3}{2}nRT$ used or $3/2 pV$ (1) AB = -36 400[J] (1)	2																																			
	(b)	(i) 0	1																																			
		(ii) Valid method either stated or clearly implied (1) Accept area under the graph Answer = - 47 250 [J] (1)	2																																			
(c)	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">AB</th> <th style="text-align: center;">BC</th> <th style="text-align: center;">CA</th> <th style="text-align: center;">ABCA</th> <th></th> </tr> </thead> <tbody> <tr> <td>W</td> <td style="text-align: center;">0</td> <td style="text-align: center;">37.6 kJ</td> <td style="text-align: center;">-47.3 kJ</td> <td style="text-align: center;">-9.7 kJ</td> <td></td> </tr> <tr> <td>ΔU</td> <td style="text-align: center;">-36.4 kJ</td> <td style="text-align: center;">33.5 kJ</td> <td style="text-align: center;">2.9 kJ</td> <td style="text-align: center;">0</td> <td style="text-align: center;">4</td> </tr> <tr> <td>Q</td> <td style="text-align: center;">-36.4 kJ</td> <td style="text-align: center;">71.1 kJ</td> <td style="text-align: center;">-44.4 kJ</td> <td style="text-align: center;">-9.7 kJ</td> <td></td> </tr> <tr> <td></td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> <td></td> </tr> <tr> <td></td> <td style="text-align: center;">ecf on ΔU</td> <td style="text-align: center;">no ecf</td> <td style="text-align: center;">ecf on W</td> <td style="text-align: center;">ecf on all if $\Delta U \approx 0$ but must make sense</td> <td></td> </tr> </tbody> </table>		AB	BC	CA	ABCA		W	0	37.6 kJ	-47.3 kJ	-9.7 kJ		ΔU	-36.4 kJ	33.5 kJ	2.9 kJ	0	4	Q	-36.4 kJ	71.1 kJ	-44.4 kJ	-9.7 kJ			✓	✓	✓	✓			ecf on ΔU	no ecf	ecf on W	ecf on all if $\Delta U \approx 0$ but must make sense		
	AB	BC	CA	ABCA																																		
W	0	37.6 kJ	-47.3 kJ	-9.7 kJ																																		
ΔU	-36.4 kJ	33.5 kJ	2.9 kJ	0	4																																	
Q	-36.4 kJ	71.1 kJ	-44.4 kJ	-9.7 kJ																																		
	✓	✓	✓	✓																																		
	ecf on ΔU	no ecf	ecf on W	ecf on all if $\Delta U \approx 0$ but must make sense																																		
Question 8 Total			[11]																																			

PH5

Question		Marking details	Marks Available
1	(a)	(i) Attempt at $6n + 6p$ - mass of carbon nucleus (1) $\times 931$ and $\div 12$ or $E = mc^2$ and $\div 12$ (1) Correct answer = 7.7 [MeV/nucleon] or 1.23×10^{-12} [J] (1)	3
		(ii) Conversion of 7.16 MeV to mass = 0.00769 [u] (1) Conservation of mass-energy $4.0015 + 11.9967 - 0.00769$ (1) Answer approx correct e.g. 15.99 u accept (16.07 ± 0.08) [u] or $(2.67 \pm 0.02) \times 10^{-26}$ [kg] (1) 15.9905 (accuracy mark, also available for 16.0059 and without unit) (1)	4
		(b)	
	(i) Neutron absorber or high melting point (or other valid response)	1	
	(ii) Light nucleus or poor neutron absorber or slows down neutrons (or other valid response)	1	
	(iii) High heat capacity or poor neutron absorber or doesn't become radioactive (or other valid response) Don't accept must be a fluid or good ability to conduct heat away	1	
	Question 1 Total		[10]

Question		Marking details	Marks Available
2	(a)	(i) Mass number = 206 (1) Atomic number = 82 (1)	2
		(ii) Gas can be <u>inhaled</u> (1) (Don't accept if state both ingest and inhale) Any 2 (×1) from: <ul style="list-style-type: none"> • Dense gas or stays in basements • Alpha <u>highly</u> ionising • Multiple emissions i.e. more than 1 alpha (do not accept emits alpha and beta by itself) • Short half-life • Contaminates wells • Enters through cracks Don't accept high activity or contaminates water supply or alpha particles are breathed in or causing cancer	3
	(b)	Use of $\lambda = \frac{\ln 2}{T_{1/2}}$ e.g. $\lambda = 0.182$ [day ⁻¹] (2.11×10^{-6} s ⁻¹) or $t = nT_{1/2}$ (1) Logs taken correctly e.g. $\ln A = \ln A_0 - \lambda t$ or $\ln A = \ln A_0 - n \ln 2$ (1) Algebra correct e.g. $t = \frac{1}{\lambda} \ln \frac{A_0}{A}$ or $n = \frac{1}{\ln 2} \ln \frac{A_0}{A}$ or implied (1) Correct answer 13.2 [days] (1.14×10^6 [s]) (1)	4
(c)	Daughter nuclei give added activity	1	
		Question 2 Total	[10]

Question			Marking details	Marks Available
3	(a)	(i)	Values substituted into $C = \frac{\epsilon_0 A}{d}$ ($= 7.32 \times 10^{-9}$ F) (1) $Q = CV$ (or implied) note $C = \frac{Q}{V}$ not good enough (1) Answer = 9.37×10^{-7} [C] (1)	3
		(ii)	Answer = 6.0×10^{-5} [J] (ecf)	1
		(iii)	$E = \frac{V}{d}$ (1) Answer = 2 170 000 [V m ⁻¹] (1)	2
	(b)	(i)	Capacitance decreases (1) Energy stored increases (1)	2
		(ii)	Work done by separating plates or work done against field or increase in potential energy (1) (accept energy used instead of work done) Equal to increase in stored energy (1)	2
Question 3 Total			[10]	

Question		Marking details	Marks Available
4	(a)	$n = \frac{9560}{1.45} \quad (1)$ <p>Correct answer = 2.65×10^{-3} T UNIT mark (1)</p>	2
	(b)	$B = \frac{\mu_0 I}{2\pi a}$ used (e.g. 2.82×10^{-6} [T] or 4.35×10^{-6} [T] or 10^x slips) (1) <p>Subtracting or adding fields (1)</p> <p>1.53×10^{-6} [T] no ecf (1)</p> <p>Out of paper (1)</p>	4
	(c)	<p>Equating fields e.g. $\frac{\mu_0 \times 0.24}{2\pi \times a} = \frac{\mu_0 \times 0.37}{2\pi \times (0.034 - a)}$ or $\frac{\mu_0 \times I_1}{2\pi \times a} = \frac{\mu_0 \times I_2}{2\pi \times b}$ (1)</p> <p>(Accept $\frac{a_1}{a_2} = \frac{0.37}{0.24}$ (= 1.54 or 0.65 reciprocal))</p> <p>Algebra $\frac{0.24 \times 0.034}{(0.37 + 0.24)} = a$ or $\frac{0.37 \times 0.034}{(0.37 + 0.24)} = a$ or $a = \frac{1.54}{2.54} \times 0.034$ etc (1)</p> <p>Answer = 0.0134 [m] or 0.0206 [m] but must be clear from algebra, working or statement that the point is nearer the upper wire (1)</p> <p>Question 4 Total</p>	3
			[9]

Question		Marking details	Marks Available
5	(a)	$F = Eq$ (or eE) used or implied (1) $E = \frac{V}{d}$ quoted or implied (1) $a = \frac{F}{m}$ used or implied (1) $a = \frac{11.2 \times 1.6 \times 10^{-19}}{9.11 \times 10^{-31} \times 7.6 \times 10^{-3}} [= 2.588 \times 10^{14}]$ (1) N.B. Use of $a = \frac{Eq}{m}$ or $F = \frac{Vq}{d}$ award 2 marks or $a = \frac{Vq}{md}$ award 3 marks	4
	(b)	(i) No horizontal forces (don't accept no horizontal acceleration or because it's in a vacuum)	1
		(ii) Constant vertical force or uniform electric field	1
	(c)	Valid method for obtaining time e.g. $s = ut + \frac{1}{2}at^2$ (1) Time correct = 5.4×10^{-9} [s] (1) Answer = $8.00 \times 10^7 \times 5.4 \text{ ns} = 43$ [cm] (ecf) (1) (factors of 10 or $\sqrt{10}$ slips only penalised 1 mark)	3
	(d)	Valid method e.g. definition of eV, force \times distance, getting resultant velocity and finding change in $\frac{1}{2}mv^2$ (1) Answer = 5.6 [eV] (which can simply be written for full marks) or 8.96×10^{-19} [J] (ecf) (1) (answer of 11.2 eV gets 1/2 marks)	2
		Question 5 Total	[11]

Question		Marking details	Marks Available
6	(a)	(i) Flux linkage = $NBA\cos\theta$ used (1) 0.251 [Wb] [and 0.251 Wb] (1)	2
		(ii) No change in flux [linkage] or field lines cut in one direction and then the opposite direction Don't accept rate of change of flux is 0	1
	(b)	Flux linkage = 0.0443 or -0.0443 (1) Time = $\frac{20}{360} \times 0.1$ (1) Attempt at change of flux (linkage) divided by time (1) Answer = [-] 15.9 [V] (1)	4
	(c)	Peak emf = 17 [V] Sinusoid with peak of 3.4 squares high (ecf) (1) Sinusoid with period of 4 squares (1)	3
		Question 6 total	[10]


Question		Marking details	Marks Available
7	(a)	<p>Any 2 (×1) from:</p> <ul style="list-style-type: none"> Near stars move relative to distant stars [due to Earth orbit] More movement (or larger angle) means stars nearer (inversely proportional etc.) or accept parsec = 1/arcsec Parallax (or distance) can be measured from readings 6 months apart (or accept readings where Earth movement is known etc.) 	2
	(b)	<p>4 parsec or angle = $1.5 \times 10^{11}/d$ (1)</p> <p>$4 \times 3.25 = 13$ [light year] (1)</p>	2
	(c)	<p>10× distance gives <u>100 times</u> less intensity (1)</p> <p>Substituting 1 and 0.1 into equation accept 1 and 10 (1)</p> <p>$m = M - 5$ and $m = M$ shown (1)</p> <p>Alternative: 2.5^5 roughly equal to 100 award 2 marks</p>	3
	(d)	<p>1[%] Accept 0.01 but not 0.01 %</p>	1
	(e)	<p>Electrons need to be in the high energy levels (1)</p> <p>They need to be in $n = 3$ (1st mark can be implied in the 2nd mark) (1)</p> <p>Not possible because no ultraviolet to absorb or collisions don't have enough KE (1)</p>	3
	(f)	<p>Comparison with $4\pi r^2 \sigma T^4$ or $b = 4\pi \sigma$ (1)</p> <p>Answer $b = 7.13 \times 10^{-7}$ (1)</p> <p>Unit = $\text{W m}^{-2} \text{K}^{-4}$ or equivalent (1)</p>	3

Question		Marking details	Marks Available
(g)	(i)	$T = 1$ (year) and $a = 1$ (AU) and $M = 1$ Accept because everything = 1	1
	(ii)	Assuming $M + m \approx 0.32 M_{\text{Sun}}$ (1) $a = \sqrt[3]{0.32 \times 0.46^2} = 0.41$ [AU] (1)	2
(h)		Drop when large eclipses small (1) And small eclipses large (1) Bigger drop when the <u>hotter/brighter</u> star is blocked (1) Award 3 marks for bigger drop when small in front of large	3
		Question 7 Total	[20]

Question		Marking details	Marks Available
8	(a)	(i) Sinusoidal reading on voltmeter @ 0.9 Hz (or across resistor) (1) Sinusoidal (or changing) B -field in primary (1) Leads to B -field cutting secondary or flux changing in secondary (1) emf induced in secondary due to Faraday's (1) (ii) Lost flux or no iron core or low frequency or low turns	4 1
	(b)	(i) $\omega L = \frac{1}{\omega C}$ or $f = \frac{1}{2\pi\sqrt{LC}}$ (1) Answer = 4 490 [Hz] (1) (ii) $V_R = 12$ [V] (1) $I = 0.067$ [A] (1) $V_L = I \times \omega L$ or $V_C = I \times \frac{1}{\omega C}$ (1) $V_L = 71.5$ [V] and $V_C = 71.5$ [V] or implied e.g. $V_C =$ same (1)	2 4
8	(c)	(i) $Z = \sqrt{(X_L - X_C)^2 + R^2}$ (1) $Z = 581$ [Ω] or implied (1) Current = $\frac{12}{581} = 21$ [mA] (1) (ii) Phasor diagram (1) $\tan\theta = \frac{X_L - X_C}{R}$ (this step implies vector diagram if omitted) (1) Answer = 72° (ecf) (1) (18° and similar slips gain 1/2)	3 3
	(d)	$\frac{R}{X_C} = \frac{3}{4}$ (1) $X_C = \frac{1}{2\pi f C}$ or $X_C = \frac{1}{\omega C}$ and $\omega = 2\pi f$ used (1) Answer = 20 [kHz] (1)	3
Question 8 Total			[20]

Question		Marking details	Marks Available
9	(a)	(i) Ørsted or Oersted (accept Orsted)	1
		(ii) Battery (not cell)	1
		(iii) Any 3 (×1) from:	3
		<ul style="list-style-type: none"> • Current passed through wire or pile connected across wire • Compass turned [nearly] at right angles to wire • When compass above wire points in opposite direction • Compass points according to rh grip (or screw) rule • Field lines circle around wire 	
	(iv) Electric effect arising from magnetism (or from magnet)	1	
	(b)	(i) Vortices shown separated by (smaller) idlers (1)	3
		Vortices and/or idlers labelled (1)	
Rotation shown or stated (1)			
(ii) Any 2 (×1) from:	2		
<ul style="list-style-type: none"> • Maxwell used it to predict e-m waves • Maxwell used it to explain magnetic field due to a wire • Maxwell used it to explain [any other e-m effect!] • Its existence is irrelevant / Maxwell didn't suppose it existed 			

Question		Marking details	Marks Available
9	(c)	(i) Produced when sparks occurred between [ball-ended] rods [connected to an induction coil or high voltage]. (1) Detected by sparks occurring across spark-gap between rods or across break in ring. (1)	2
		(ii) He found spark intensity varied according to orientation of detector rods [relative to transmitter rods]. or he interposed metal grille between transmitter and detector, finding spark intensity varied with grille orientation.	1
		(iii) He used metal reflector to produce stationary wave. (1) He measured distance between nodes [and doubled it]. (1)	2
	(d)	(i) Time between events in a frame in which the events occur at the same place. or time between events as measured by a clock present at both events.	1
		(ii) $\gamma = 1.01$ (1) $t_B - t_A = 0.5000 \gamma$ (1) despite mistakes in γ $t_B - t_A = 0.5050$ [s] (1) allow ecf on γ arising from slips.	3
Question 9 total			[20]

Question		Marking details	Marks Available
10	(a)	<p>Elastic, straight line (1) yield point (1) curve (1) <u>region</u> of plastic deformation(1)</p> 	6
	(b)	<p>(i) DE broken or E bonds with B (1) HG broken or H bonds with D (1) Movement of dislocations stated (1) Or all clearly seen from diagrams</p> <p>(ii) No dislocations (or equivalent) or no grain boundaries (1) Don't accept addition of foreign atoms Don't accept single crystal (stated in question)</p> <p>(iii) Any valid use e.g. (1) Turbine blades (don't accept wind turbines), combustion chambers, nuclear reactors, wear resistant materials, rocket engines etc.</p>	3 1 1
	(c)	<p>(i) $\frac{Fl}{2 \times 10^{11} A_{steel}} = \frac{Fl}{1 \times 10^{11} A_{brass}}$ (1) Convincing algebra to show $A_{brass} = 2 \times A_{steel}$ (1) (alternative: force, length and extension all the same 1 mark so brass must have twice the CSA 1 mark only – not fully shown as required)</p> <p>(ii) 50 [N] (1)</p> <p>(iii) $\Delta x = \frac{50 \times 2}{(2.8 \times 10^{-7}) \times 2 \times 10^{11}}$ (1) – substitution (ecf on 50 N) $\Delta x = 1.8$ [mm] (1) (correct unit required m or mm)</p> <p>(iv) $E = \frac{1}{2} Fx$ (1) (accept $E = \frac{1}{2} \sigma \epsilon V$) (1) $E = 0.044$ [J] (1) (ecf on Δx only)</p> <p>(v) Same (1) F and Δx same (1)</p> <p>Question 10 total</p>	2 1 2 2
Question 10 total			[20]

Question			Marking details	Marks Available	
11	(a)	(i)	Both background and line spectra labelled clearly	1	
		(ii)	I	[Inner] electrons [of target element] knocked out / ionised (1) Electrons from higher energy levels fall to take their place (1)	4
			II	Rapid deceleration of <u>electrons</u> (1) On collision with target element / nucleus (1)	
	(iii)		$\lambda = \frac{hc}{eV}$ (or rearrangement in figures) (1) $\lambda = 2.07 \times 10^{-11}$ [m] (1) Accept 2.1×10^{-11} [m]	2	
	(b)	(i)	Ultrasound B-scan (1) Moving pictures/ see organ development not 'give a 2D image' (1)	6	
		(ii)	CT scan (1) Distinguishes soft tissue well (1) Accept MRI cannot be used because of pacemaker		
		(iii)	MRI scan (1) Gives high quality images of soft tissue (1)		
	(c)		Time taken from scale 5 ± 1 [μ s] (1) Distance = 8.2×10^{-3} [m] (1) (ecf) Thickness = $\frac{8.2 \times 10^{-3}}{2} = 4.1 \times 10^{-3}$ [m] (1)	3	


Question			Marking details	Marks Available
11	(d)	(i)	QRS wave / R / central spike (1) Bigger / higher /more spiked (1)	2
		(ii)	P wave flatter / P wave extended/ prolonged PR interval / no P wave / smaller P wave / P wider / P lower amplitude	1
		(iii)	Deepening of Q wave / T wave inversion / ST elevation / ST depression { Irregular interval / inverted waves / bigger distance P to QRS to T wave} N.B. any incorrect statement negates the mark	1
			Question 11 total	[20]


Question		Marking details	Marks Available
12	(a)	(i) Any 2 ×(1) from: <ul style="list-style-type: none"> • Possible second use as a bridge • Cheap electricity after build • Zero or low CO₂ after built • High output • Predictable output • Sustainable/renewable/reliable energy source that will not run out 	2
		(ii) Any 2 ×(1) from: <ul style="list-style-type: none"> • Only available twice a day (i.e. not a constant output) • Possible huge impact on Severn estuary wildlife • High CO₂ costs to build • Expensive to build (£3k per kW as opposed to £1k per kW coal) 	2
	(b)	GPE (PE not good enough) to KE or GPE to electrical (1) KE / mechanical to electrical or const KE when running (1)	2
	(c)	Mean height increase = 0.5 <i>h</i> must be stated not implied (1) Either volume = <i>Ah</i> or mass = <i>Ahρ</i> (1) Correct substitution into <i>mgh</i> (ecf) (1)	3
	(d)	Values substituted into equation (1.38×10^{14} J) (1) ×2 (or using time as 12 hrs) and × 0.75 (1) Dividing by time or $P = E/t$ etc. (1) Answer = 2.4 [GW] (no ecf) (1)	4

Question			Marking details	Marks Available
12	(e)	(i)	Any 2 ×(1) from: <ul style="list-style-type: none"> • Continuous electricity not twice a day • Less damage to environment • Less impact on shipping 	2
		(ii)	Mass per second = $A\rho v$ (or implied) (1) KE per second = $0.5 A\rho v v^2$ (or good attempt at $0.5mv^2$) (1) $\times 0.75$ (1) Answer = 3.1 [GW] (no ecf) (1)	4
		(iii)	Because $\propto v^3 $ (accept v^2, v^3 etc.) and we need $ \overline{v^3} $ not $ \overline{v} $ Wordy answers also valid e.g. 'if speed is 1.5 and 4.5 say, 4.5^2 will be far more significant than 1.5^2 or 'the power output for high speeds will be far greater than for low speeds increasing the mean power' etc.	1
Question 12 total				[20]

Question		Marking details	Marks Available
(d)	(i)	<p>Large triangles used (should be close to the extremities of the lines) or 2 equivalent suitable points clearly indicated on each line. (1) Both gradients calculated correctly (ignore unit and significant figures) (1 + 1) Allow ecf for incorrect max/min lines. Exemplar values – values must be in agreement with candidate’s graph.</p> $\text{Max gradient} = \frac{6.33 - 4.69}{80.0 - 0.0} = 0.0205 \text{ } [\Omega^{\circ}\text{C}^{-1}]$ $\text{Min gradient} = \frac{6.17 - 4.86}{80.0 - 0.0} = 0.0164 \text{ } [\Omega^{\circ}\text{C}^{-1}]$ <p>Marking tips: First check: The value of m_{max} should be $\sim 0.021 \text{ } [\Omega^{\circ}\text{C}^{-1}]$ and the value of m_{min} should be about $0.016 \text{ } [\Omega^{\circ}\text{C}^{-1}]$. Candidates who have drawn lines which do not take full advantage of the error bars may get <0.020 and >0.017 respectively. This is penalised in (c), so apply ecf. Candidates who have drawn ‘tram lines’ will have two nearly identical values of ~ 0.018. Again ecf should be applied.</p>	3
	(ii)	<p>Mean gradient correct (1) [Exemplar value $\sim 0.0184[5] \text{ } [\Omega^{\circ}\text{C}^{-1}]$ but apply ecf from (b) and (d)(i)]. No unit penalty. Percentage uncertainty correct (1) [Exemplar value $\sim 11\%$. Allow 1 or 2 sig figs. Apply ecf from (b) and (d)(i)].</p>	2
(e)	(i)	<p>Mean value correct [Exemplar value $4.78 \text{ } [\Omega]$]. (1) Percentage uncertainty correct [Exemplar value $\sim 2\%$ - allow 1 or 2 sig figs]. (1) Allow ecf and for 3 sig figs in % uncertainty Intercept = Resistance (of copper) at 0°C. (1)</p>	3
	(ii)	<p>$\alpha = \frac{\text{gradient}}{R_0}$ or $\alpha = \frac{\text{gradient}}{\text{intercept}}$ stated or implied by calc. (1) Correct calculation, i.e. $\alpha = \frac{\text{answer to (d)(ii)}}{\text{answer to (e)(i)}} \text{ (1)}$</p>	4
	(iii)	<p>$\alpha = 3.9 \times 10^{-3} \text{ } [\text{Accept answer in range } 3.8 \text{ to } 4.0 \times 10^{-3}]. \text{ (1)}$ Note: This mark is for accuracy. Do not apply ecf. No sig figs penalty. Unit given as $^{\circ}\text{C}^{-1}$ (or K^{-1}). (1) N.B. If data points selected from the graph or table (1), calculation of α (1), correct unit (1) i.e. maximum of 3 marks awarded.</p> <p>Total % uncertainty = % in (d)(ii) + % in (e)(i). (1) [Exemplar value $\sim 13\%$. Apply ecf] Absolute uncertainty correct and given to 1 or 2 sig figs. (1) apply ecf [Exemplar value $\sim 0.5 \times 10^{-3}$] Temperature coefficient of resistance written correctly with its uncertainty, ignore unit, the value given to number of sig figs consistent with uncertainty [e.g. $0.0039 \pm 0.0005 \text{ } ^{\circ}\text{C}^{-1}$; $(3.9 \pm 0.5) \times 10^{-3} \text{ } ^{\circ}\text{C}^{-1}$]. (1) Award the mark if α and absolute uncertainty calculated correctly but written separately.</p>	3
Question total			25

Resistance / Ω





PH6
EXPERIMENTAL TASK
MARK SCHEME – TEST 1

Question	Marking details	Marks Available
(a)	$\ln T = n \ln d + \ln k$ (1) Graph of $\ln T$ (y-axis) against $\ln d$ (x-axis) stated to be plotted or explicit comparison with $y = mx + c$ (1) (accept any logarithm) <i>Remember not to award the marks if the information sheet was issued.</i>	2
(b)	Explain or state a method to ensure the threads are vertical or explanation of thread movement to ensure symmetry implied. (1) accept reference to paired readings or ruler being horizontal Minimum of 5 readings between and including the range 30 cm to 60 cm. (1) Minimum of 5 oscillations timed for each value of d . (1) Repeat readings taken. (1) All of the above points cannot be awarded from the table they must be stated in their plan.	4
(c)	Single, clear, main table with titles and units on each column. (1) Accept if candidates have recorded d in m or cm. N.B. log values should have no unit e.g. $\ln(T/s)$ is acceptable, however $\ln T/s$ and $\ln T/\ln s$ are both not acceptable. Repeat readings given and mean values correct. (1) Period of 1 oscillation calculated correctly and to 3/4 sig figs. ecf (1) The resolution of the ruler given as ± 1 mm and the stopwatch as ± 0.01 s: can be awarded from plan (1) All log values calculated correctly to 2 decimal places (accept 3 dp) and d to nearest mm (1)	5
(d)	Graph of $\ln T$ against $\ln d$ plotted with axis labelled and no units (ecf units from table). (1) Suitable scale so that data points occupy at least half of each axis. (1) All points plotted correctly to within $\frac{1}{2}$ small square division. (1) Good line of best fit consistent with the data. (1)	4
(e)	Large triangle used (should be close to the extremities of the line of best fit i.e. over half the line used) [or 2 equivalent suitable points clearly indicated on the graph]. (1) Gradient calculated correctly including negative sign. (1) Gradient = n clearly stated. (1)	3
(f)	(i) Yes - with 2 × (1) of the following points: (N.B. gradient in (e) must be between 0.70 to 1.30) <ul style="list-style-type: none"> • Gradient is negative; (1) • Gradient or n is approximately / equal to 1; (1) • Inversely proportional or T is proportional to d^{-1}. (1) Alternative: Yes because $n / \text{gradient} = -1$. (2) N.B. no ecf allowed from the graph or part (e) i.e. if graph is wrong and value of n is wrong they lose these 2 marks. (Yes on its own = 0 marks.) Award a maximum of 2 marks only.	2
	(ii) Rearrangement $b = \frac{T^2 d^2}{4\pi^2 l}$ (1) Correct calculation of b using points taken from their table. (1) Units cm s^2 (accept m s^2 if consistent with calculation). (1)	3
	(iii) Increase / become longer. (1) by $\sqrt{2}$. (1)	2
	Question total	[25]

MARK SCHEME – TEST 2
Same as for TEST 1



WJEC
245 Western Avenue
Cardiff CF5 2YX
Tel No 029 2026 5000
Fax 029 2057 5994
E-mail: exams@wjec.co.uk
website: www.wjec.co.uk